PID No.
 : MED120843380
 Register On
 : 26/11/2022 7:51 AM

 SID No.
 : 522229076
 Collection On
 : 26/11/2022 9:42 AM

 Age / Sex
 : 54 Year(s) / Male
 Report On
 : 26/11/2022 2:33 PM

Printed On



Ref. Dr : MediWheel

: OP

Type

Investigation IIA ENA TOLOGY	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HAEMATOLOGY Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood/Spectrophotometry)	15.8	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	49.6	%	42 - 52
RBC Count (EDTA Blood)	5.37	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (EDTA Blood)	92.5	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood)	29.5	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood)	31.9	g/dL	32 - 36
RDW-CV	14.0	%	11.5 - 16.0
RDW-SD	45.33	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood)	7400	cells/cu.mm	4000 - 11000
Neutrophils (Blood)	52.6	%	40 - 75
Lymphocytes (Blood)	38.8	%	20 - 45
Eosinophils (Blood)	1.8	%	01 - 06
Monocytes (Blood)	6.5	%	01 - 10

: 06/12/2022 11:14 AM



 PID No.
 : MED120843380
 Register On
 : 26/11/2022 7:51 AM

 SID No.
 : 522229076
 Collection On
 : 26/11/2022 9:42 AM

 Age / Sex
 : 54 Year(s) / Male
 Report On
 : 26/11/2022 2:33 PM

Printed On



Ref. Dr : MediWheel

: OP

Type

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Basophils (Blood)	0.3	%	00 - 02
INTERPRETATION: Tests done on Automated Five P	art cell counter. All	abnormal results are re	viewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood)	3.89	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood)	2.87	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood)	0.13	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood)	0.48	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood)	0.02	10^3 / μl	< 0.2
Platelet Count (EDTA Blood)	163	10^3 / μl	150 - 450
MPV (Blood)	10.6	fL	7.9 - 13.7
PCT (Automated Blood cell Counter)	0.17	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood)	4	mm/hr	< 20

: 06/12/2022 11:14 AM



 PID No.
 : MED120843380
 Register On
 : 26/11/2022 7:51 AM

 SID No.
 : 522229076
 Collection On
 : 26/11/2022 9:42 AM

 Age / Sex
 : 54 Year(s) / Male
 Report On
 : 26/11/2022 2:33 PM

Printed On



Ref. Dr : MediWheel

: OP

Type

<u>Investigation</u>	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	0.45	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.15	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.30	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	35.37	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	48.57	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	28.63	U/L	< 55
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	72.8	U/L	56 - 119
Total Protein (Serum/Biuret)	6.89	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.33	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	2.56	gm/dL	2.3 - 3.6
A: GRATIO (Serum/Derived)	1.69		1.1 - 2.2

: 06/12/2022 11:14 AM



PID No. : MED120843380 Register On : 26/11/2022 7:51 AM : 522229076 Collection On : 26/11/2022 9:42 AM SID No. Age / Sex : 54 Year(s) / Male Report On : 26/11/2022 2:33 PM



Type : OP

: 06/12/2022 11:14 AM **Printed On**

Ref. Dr : MediWheel

Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	232.82	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	144.61	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual_ circulating level of triglycerides during most part of the day.

F			
HDL Cholesterol (Serum/Immunoinhibition)	44.61	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	159.3	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	28.9	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	188.2	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220



PID No. : MED120843380 Register On : 26/11/2022 7:51 AM : 522229076 SID No. Collection On : 26/11/2022 9:42 AM Age / Sex : 54 Year(s) / Male Report On : 26/11/2022 2:33 PM



Type : OP

: 06/12/2022 11:14 AM **Printed On**

Ref. Dr : MediWheel

Investigation <u>Observed</u> **Unit Biological** <u>Value</u> Reference Interval

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio 5.2 Optimal: < 3.3Low Risk: 3.4 - 4.4 (Serum/Calculated) Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0 Triglyceride/HDL Cholesterol Ratio 3.2 Optimal: < 2.5Mild to moderate risk: 2.5 - 5.0 (TG/HDL)

High Risk: > 5.0(Serum/Calculated)

LDL/HDL Cholesterol Ratio 3.6 Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 (Serum/Calculated) High Risk: > 6.0



PID No. : MED120843380 Register On : 26/11/2022 7:51 AM : 522229076 SID No. Collection On : 26/11/2022 9:42 AM

Age / Sex : 54 Year(s) / Male Report On : 26/11/2022 2:33 PM

Printed On

Type : OP

Ref. Dr : MediWheel

: 06/12/2022 11:14 AM



Investigation Observed Unit Biological Reference Interval <u>Value</u> Glycosylated Haemoglobin (HbA1c) Normal: 4.5 - 5.6 HbA1C 6.0 % Prediabetes: 5.7 - 6.4 (Whole Blood/HPLC) Diabetic: \geq 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 125.5 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.



: MED120843380 **Register On** : 26/11/2022 7:51 AM

Printed On

SID No. : 522229076 Collection On : 26/11/2022 9:42 AM

Type : OP

PID No.

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u> <u>Ur</u>	<u>it</u> <u>Biological</u>
	Value	Reference Interval

: 06/12/2022 11:14 AM

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.34 ng/ml 0.4 - 1.81

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is

Metabolically active.

T4 (Tyroxine) - Total 10.56 μg/dl 4.2 - 12.0

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 3.09 µIU/mL 0.35 - 5.50

(Serum/ECLIA)

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



Printed On

Type : OP

Ref. Dr : MediWheel

PID No.



InvestigationObservedUnitBiologicalValueReference Interval

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING 'A' 'Positive'

(EDTA Blood/Agglutination)

INTERPRETATION: Note: Slide method is screening method. Kindly confirm with Tube method for transfusion.



APPROVED BY

: 06/12/2022 11:14 AM

 PID No.
 : MED120843380
 Register On
 : 26/11/2022 7:51 AM

 SID No.
 : 522229076
 Collection On
 : 26/11/2022 9:42 AM

Report On



Type : OP

Ref. Dr : MediWheel

Age / Sex : 54 Year(s) / Male

Printed On : 06/12/2022 11:14 AM

26/11/2022 2:33 PM

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	12.17		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	97.46	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine)	Negative		Negative
(Urine - F/GOD - POD)			
Glucose Postprandial (PPBS)	91.46	mg/dL	70 - 140
(Dlogmo DD/COD DAD)			

(Plasma - PP/GOD-PAP)

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	12.3	mg/dL	7.0 - 21
Creatinine (Serum/Modified Jaffe)	1.01	mg/dL	0.9 - 1.3

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetylcyteine, chemotherapeutic agent such as flucytosine etc.

Uric Acid	5.24	mg/dL	3.5 - 7.2
-----------	------	-------	-----------





 PID No.
 : MED120843380
 Register On
 : 26/11/2022 7:51 AM

 SID No.
 : 522229076
 Collection On
 : 26/11/2022 9:42 AM

 SID No.
 : 522229076
 Collection On : 26/11/2022 9:42 AM

 Age / Sex
 : 54 Year(s) / Male
 Report On : 26/11/2022 2:33 PM

Type : OP **Printed On** : 06/12/2022 11:14 AM

Ref. Dr : MediWheel



<u>Investigation</u>	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>IMMUNOASSAY</u>			
Prostate specific antigen - Total(PSA) (Serum/Manometric method)	0.300	ng/ml	Normal: 0.0 - 4.0 Inflammatory & Non Malignant conditions of Prostate & genitourinary system: 4.01 - 10.0 Suspicious of Malignant disease of Prostate: > 10.0



APPROVED BY

-- End of Report --



Name	Mr.SINGH SUSHIL KUMAR	ID	MED120843380
Age & Gender	54/MALE	Visit Date	26/11/2022
Ref Doctor Name	MediWheel		

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in shape, size (12.5cms) and has uniform echopattern. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER shows normal shape and has clear contents. Wall is of normal thickness. CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN shows normal shape, size and echopattern.

No demonstrable Para-aortic lymphadenopathy.

BOTH KIDNEYS

Right kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

Left kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

·	Bipolar length (cms)	Parenchymal thickness (cms)	
Right Kidney	8.7	1.3	
Left Kidney	9.1	1.2	

URINARY BLADDER shows normal shape and wall thickness. It has clear contents. No evidence of diverticula.

PROSTATE shows normal shape, size and echopattern.

No evidence of ascites.

IMPRESSION:

• No significant abnormality detected.

REPORT DISCLAIMER

- 1. This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7. Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11.Disputes, if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



Name	Mr.SINGH SUSHIL KUMAR	ID	MED120843380
Age & Gender	54/MALE	Visit Date	26/11/2022
Ref Doctor Name	MediWheel		

DR. S K SOMU ELANGOVAN CONSULTANT RADIOLOGIST SKS/an

REPORT DISCLAIMER

- 1.This is only a radiological imperssion.Like other investigations, radiological investication also have limitation. Therefore radiological reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11. Disputes, if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.