

CLIENT'S NAME AND ADDRESS:
ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI

DUVCTON EVANINATION LIDINE

SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL Ltd

S.K. Tower,Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: SUNIL AJAGEKAR PATIENT ID: SUNIM100678181A

AGE: 44 Years ACCESSION NO: 0181VI000857 SEX: Male

DRAWN: RECEIVED: 24/09/2022 08:11 REPORTED: 26/09/2022 15:43

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units **Final**

MEDI WHEEL FULL BODY HEALTH CHECK UP ABOVE 40 MALE

PHYSICAL EXAMINATION, URINE				
COLOR	PALE YELLOW			
METHOD: VISUAL INSPECTION				
APPEARANCE	CLEAR			
METHOD: VISUAL INSPECTION				
SPECIFIC GRAVITY	1.005	1.003 - 1.035		
METHOD: IONIC CONCENTRATION METHOD				
BLOOD COUNTS,EDTA WHOLE BLOOD				
HEMOGLOBIN	16.9	13.0 - 17.0	g/dL	
METHOD: SLS-HEMOGLOBIN DETECTION METHOD				
RED BLOOD CELL COUNT	5.31	4.5 - 5.5	mil/µL	
METHOD: HYDRODYNAMIC FOCUSING BY DC DETECTION				
WHITE BLOOD CELL COUNT	5.86	4.0 - 10.0	thou/µL	
METHOD: FLUORESCENCE FLOW CYTOMETRY				
PLATELET COUNT	1/4	150 - 410	thou/µL	
METHOD: HYDRODYNAMIC FOCUSING BY DC DETECTION				
RBC AND PLATELET INDICES				
HEMATOCRIT	49.9	40.0 - 50.0	%	
METHOD: CUMULATIVE PULSE HEIGHT DETECTION METHOD				
MEAN CORPUSCULAR VOL	94.0	83.0 - 101.0	†L	
METHOD: CALCULATED FROM RBC & HCT				
MEAN CORPUSCULAR HGB.	31.8	27.0 - 32.0	pg	
METHOD: CALCULATED FROM THE RBC & HGB				
MEAN CORPUSCULAR HEMOGLOBIN	33.9	31.5 - 34.5	g/dL	
CONCENTRATION METHOD: CALCULATED FROM THE HGB & HCT				
MENTZER INDEX	17.7			
RED CELL DISTRIBUTION WIDTH	12.9	11.6 - 14.0	%	
METHOD : CALCULATED FROM RBC SIZE DISTRIBUTION CURVE	12.3	11.0 1 1.0	,0	
MEAN PLATELET VOLUME	10.3	6.8 - 10.9	fL	
METHOD : CALCULATED FROM PLATELET COUNT & PLATELET HEMA		5.5		
CHEMICAL EXAMINATION, URINE				
P -	6.0	4.7 - 7.5		
METHOD : DOUBLE INDICATOR PRINCIPLE	-·-			
PROTEIN	NOT DETECTED	NOT DETECTED		



METHOD: TETRA BROMOPHENOL BLUE/SULFOSALICYLIC ACID

Page 1 Ot 12 Scan to View Report



CLIENT'S NAME AND ADDRESS:
ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI

NEW DELHI 110030 DELHI INDIA 8800465156

SRL Ltd

S.K. Tower,Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: SUNIL AJAGEKAR

PATIENT ID: SUNIM100678181A

ACCESSION NO: 0181VI000857 AGE: 44 Years SEX: Male

DRAWN: RECEIVED: 24/09/2022 08:11 REPORTED: 26/09/2022 15:43

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Referen	ce Interval Units
GLUCOSE	NOT DETECTED	NOT DETECTED	
METHOD : GLUCOSE OXIDASE PEROXIDASE			
KETONES	NOT DETECTED	NOT DETECTED	
METHOD: NITROPRUSSIDE REACTION			
BLOOD	NOT DETECTED	NOT DETECTED	
METHOD : PEROXIDASE			
BILIRUBIN	NOT DETECTED	NOT DETECTED	
UROBILINOGEN	NORMAL	NORMAL	
METHOD: MODIFIED EHRLICH REACTION			
NITRITE	NOT DETECTED	NOT DETECTED	
METHOD: 1,2,3,4-TETRAHYDROBENZO(H)QUINOLIN			
LEUKOCYTE ESTERASE	NOT DETECTED	NOT DETECTED	
WBC DIFFERENTIAL COUNT - NLR			
SEGMENTED NEUTROPHILS	58	40 - 80	%
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERI	NG		
ABSOLUTE NEUTROPHIL COUNT	3.40	2.0 - 7.0	thou/µL
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERI			
LYMPHOCYTES	33	20 - 40	%
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERI			
ABSOLUTE LYMPHOCYTE COUNT	1.94	1.0 - 3.0	thou/µL
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERI			
NEUTROPHIL LYMPHOCYTE RATIC (NLR)	1.8		
EOSINOPHILS	6	1 - 6	%
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERI			
ABSOLUTE EOSINOPHIL COUNT	0.33	0.02 - 0.50	thou/µL
METHOD : FLOW CYTOMETRY WITH LIGHT SCATTERI			
MONOCYTES	3	2 - 10	%
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERI			
ABSOLUTE MONOCYTE COUNT	0.19	Low 0.2 - 1.0	thou/µL
METHOD : FLOW CYTOMETRY WITH LIGHT SCATTERI			
DIFFERENTIAL COUNT PERFORMED ON:	EDTA SMEAR		
MICROSCOPIC EXAMINATION, URINE			
PUS CELL (WBC'S)	1-2	0-5	/HPF
METHOD: MICROSCOPIC EXAMINATION			
EPITHELIAL CELLS	1-2	0-5	/HPF
METHOD: MICROSCOPIC EXAMINATION			
ERYTHROCYTES (RBC'S)	NOT DETECTED	NOT DETECTED	/HPF
METHOD: MICROSCOPIC EXAMINATION			



Page 2 Ot 12 Scan to View Report



CLIENT'S NAME AND ADDRESS:
ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI

NEW DELHI 110030 DELHI INDIA 8800465156

S.K. Tower,Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: SUNIL AJAGEKAR PATIENT ID: SUNIM100678181A

SRL Ltd

ACCESSION NO: 0181VI000857 AGE: 44 Years SEX: Male

DRAWN: RECEIVED: 24/09/2022 08:11 REPORTED: 26/09/2022 15:43

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status	<u>Final</u>	Results		Biological Reference Interva	al Units
CASTS		NOT DETECTED			
METHOD : MICROSCOPIC EX	AMINATION	NOT DETECTED			
CRYSTALS	ANTHIATTON	NOT DETECTED			
METHOD : MICROSCOPIC EX	AMINATION	NOT DETECTED		NOT DETECTED	
BACTERIA	AMAINIATION	NOT DETECTED		NOT DETECTED	
METHOD: MICROSCOPIC EX YEAST	AMINATION	NOT DETECTED		NOT DETECTED	
		NOT DETECTED		NOT DETECTED	
MORPHOLOGY					
RBC		NORMOCYTIC NORMOCHROMIC			
WBC		NORMAL MORPHOLO	GΥ		
METHOD: MICROSCOPIC EX	AMINATION				
PLATELETS		ADEQUATE			
ERYTHRO SEDIMENTA	ATION RATE, BLOOD				
SEDIMENTATION RATE	(ESR)	01		0 - 14	mm at 1 hr
METHOD: WESTERGREN ME					
GLYCOSYLATED HEM	OGLOBIN, EDTA WHOLE BL	.OOD			
GLYCOSYLATED HEMOC	GLOBIN (HBA1C)	5.3		Non-diabetic: < 5.7 Pre-diabetics: 5.7 - 6.4 Diabetics: > or = 6.5 ADA Target: 7.0 Action suggested: > 8.0	%
METHOD: HPLC					
MEAN PLASMA GLUCOS		105.4		< 116.0	mg/dL
METHOD : CALCULATED PARA					
GLUCOSE, FASTING, I					
GLUCOSE, FASTING, PL	.ASMA	102	High	Normal 75 - 99 Pre-diabetics: 100 - 125 Diabetic: > or = 126	mg/dL
	RENCE METHOD WITH HEXOKINASE				
GLUCOSE, POST-PRA	•				
GLUCOSE, POST-PRAND	DIAL, PLASMA	98		70 - 139	mg/dL
	RENCE METHOD WITH HEXOKINASE				
CORONARY RISK PRO	OFILE, SERUM				
CHOLESTEROL		257	High	Desirable cholesterol level < 200 Borderline high cholesterol 200 - 239 High cholesterol > / = 240	mg/dL
METHOD - ENTERMATIC COLO	DIMETRIC ACCA)(- , - 2.10	

METHOD: ENZYMATIC COLORIMETRIC ASSAY







CLIENT'S NAME AND ADDRESS:
ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI

NEW DELHI 110030 DELHI INDIA 8800465156

SRL Ltd

S.K. Tower,Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: SUNIL AJAGEKAR

PATIENT ID: SUNIM100678181A

ACCESSION NO: 0181VI000857 AGE: 44 Years SEX: Male

DRAWN: RECEIVED: 24/09/2022 08:11 REPORTED: 26/09/2022 15:43

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results		Biological Reference Interv	al Units
TRIGLYCERIDES METHOD: ENZYMATIC COLORIMETRIC ASSAY	223	High	Normal: < 150 Borderline high: 150 - 199 High: 200 - 499 Very High: >/= 500	mg/dL
HDL CHOLESTEROL	41		Low HDL Cholesterol <40	mg/dL
METHOD: ENZYMATIC, COLORIMETRIC			High HDL Cholesterol >/= 60	1
CHOLESTEROL LDL	171	High	Adult levels: Optimal < 100 Near optimal/above optimal: 1 129 Borderline high: 130-159 High: 160-189 Very high: = 190	mg/dL 00-
METHOD : ENZYMATIC COLORIMETRIC ASSAY NON HDL CHOLESTEROL	216	High	Desirable: < 130 Above Desirable: 130 -159 Borderline High: 160 - 189 High: 190 - 219 Very high: > / = 220	mg/dL
CHOL/HDL RATIO	6.3	High	Low Risk: 3.3 - 4.4 Average Risk: 4.5 - 7.0 Moderate Risk: 7.1 - 11.0 High Risk: > 11.0	
LDL/HDL RATIO	4.2	High	0.5 - 3.0 Desirable/Low Risk 3.1 - 6.0 Borderline/Moderate Risk >6.0 High Risk	
VERY LOW DENSITY LIPOPROTEIN	44.6	High	< OR = 30.0	mg/dL
LIVER FUNCTION PROFILE, SERUM				
BILIRUBIN, TOTAL	1.29	High	Upto 1.2	mg/dL
METHOD: COLORIMETRIC DIAZO	0.44		. 0.00	6.10
BILIRUBIN, DIRECT	0.44	High	< 0.30	mg/dL
BILIRUBIN, INDIRECT	0.85		0.1 - 1.0	mg/dL
TOTAL PROTEIN METHOD: COLORIMETRIC	7.5		6.0 - 8.0	g/dL
ALBUMIN METHOD: COLORIMETRIC	4.9		3.97 - 4.94	g/dL
GLOBULIN	2.6		2.0 - 3.5	g/dL
ALBUMIN/GLOBULIN RATIO	1.9		1.0 - 2.1	RATIO
ASPARTATE AMINOTRANSFERASE (AST/SGOT) METHOD: UV ABSORBANCE	25		< OR = 50	U/L



Page 4 Ot 12 Scan to View Report



CLIENT'S NAME AND ADDRESS:
ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI

ACCESSION NO: 0181VI000857

NEW DELHI 110030 DELHI INDIA 8800465156

SRL Ltd S.K. Tower,Hari Niwas, LBS Marg
THANE, 400602
MAHARASHTRA, INDIA
Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956
Email: customercare.thane@srl.in

PATIENT ID: SUNIM100678181A

PATIENT NAME: SUNIL AJAGEKAR

DRAWN: RECEIVED: 24/09/2022 08:11 REPORTED: 26/09/2022 15:43

SEX: Male

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

AGE: 44 Years

Test Report Status <u>Final</u>	Results		Biological Reference	Interval Units
ALANINE AMINISTRANICEDADE (ALTICODE)	27		4 OD _ FO	1.171
ALANINE AMINOTRANSFERASE (ALT/SGPT) METHOD: UV ABSORBANCE	37		< OR = 50	U/L
ALKALINE PHOSPHATASE	74		40 - 129	U/L
METHOD: COLORIMETRIC				-, -
GAMMA GLUTAMYL TRANSFERASE (GGT)	21		0 - 60	U/L
METHOD : ENZYMATIC, COLORIMETRIC				
LACTATE DEHYDROGENASE	232	High	125 - 220	U/L
METHOD: UV ABSORBANCE				
SERUM BLOOD UREA NITROGEN				
BLOOD UREA NITROGEN	11		6 - 20	mg/dL
METHOD : ENZYMATIC ASSAY CREATININE, SERUM				
CREATININE CREATININE	0.89		0.7 - 1.2	mg/dL
METHOD: COLORIMETRIC	0.09		0.7 1.2	mg/ac
BUN/CREAT RATIO				
BUN/CREAT RATIO	12.36		8.0 - 15.0	
URIC ACID, SERUM				
URIC ACID	7.6	High	3.4 - 7.0	mg/dL
METHOD: ENZYMATIC COLORIMETRIC ASSAY				
TOTAL PROTEIN, SERUM				
TOTAL PROTEIN	7.5		6.0 - 8.0	g/dL
METHOD: COLORIMETRIC				
ALBUMIN, SERUM				
ALBUMIN	4.9		3.97 - 4.94	g/dL
METHOD: COLORIMETRIC GLOBULIN				
GLOBULIN	2.6		2.0 - 3.5	g/dL
ELECTROLYTES (NA/K/CL), SERUM	2.0		2.0 - 3.3	g/uL
SODIUM	135	Low	136 - 145	mmol/L
POTASSIUM	4.06	2017	3.5 - 5.1	mmol/L
CHLORIDE	99		98 - 107	mmol/L
THYROID PANEL, SERUM	22		50 107	TTITTOI) L
T3	103.0		80 - 200	ng/dL
METHOD: ELECTROCHEMILUMINESCENCE	200.0			
T4	8.48		5.1 - 14.1	µg/dL
METHOD: ELECTROCHEMILUMINESCENCE				
TSH 3RD GENERATION	2.050		0.27 - 4.2	μIU/mL







ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI NEW DELHI 110030

DELHI INDIA 8800465156

SRL Ltd S.K. Tower, Hari Niwas, LBS Marg THANE, 400602

MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT ID:

SUNIM100678181A

Email: customercare.thane@srl.in

PATIENT NAME: SUNIL AJAGEKAR

DRAWN: RECEIVED: 24/09/2022 08:11 REPORTED: 26/09/2022 15:43

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

AGE: 44 Years

Test Report Status Results Biological Reference Interval Units <u>Final</u>

SEX: Male

METHOD: ELECTROCHEMILUMINESCENCE STOOL: OVA & PARASITE

ACCESSION NO: 0181VI000857

REMARK SAMPLE NOT RECEIVED

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

ABO GROUP TYPE B

METHOD: GEL COLUMN AGGLUTINATION METHOD.

RH TYPE **POSITIVE**

METHOD: GEL COLUMN AGGLUTINATION METHOD.

XRAY-CHEST

IMPRESSION NO ABNORMALITY DETECTED

TMT OR ECHO

TMT OR ECHO 2D ECHO: - MILD CONCENTRIC LVH.

ECG

ECG T ABNORMALITY IN INFERIOR LEADS.

MEDICAL HISTORY

RELEVANT PRESENT HISTORY HYPERTENSIV SINCE 1 YEAR.

RELEVANT PAST HISTORY PAST H/O DYSLIPIDEMIA NOT ON ANY TREATMENT LEFT CLAVICLE

FRACTURE TREATED CONSERVATIVELY.

RELEVANT PERSONAL HISTORY MARRIED / 2 CHILD / MIXED DIET / NO ALLERGIES / NO SMOKING /

OCC ALCÓHOL.

RELEVANT FAMILY HISTORY NOT SIGNIFICANT HISTORY OF MEDICATIONS NOT SIGNIFICANT

ANTHROPOMETRIC DATA & BMI

HEIGHT IN METERS 1.72 mts WEIGHT IN KGS. 89 Kas BMI 30 BMI & Weight Status as follows: kg/sqmts

Below 18.5: Underweight 18.5 - 24.9: Normal

25.0 - 29.9: Overweight 30.0 and Above: Obese

GENERAL EXAMINATION

MENTAL / EMOTIONAL STATE NORMAL PHYSICAL ATTITUDE NORMAL GENERAL APPEARANCE / NUTRITIONAL STATUS OBESE BUILT / SKELETAL FRAMEWORK **AVERAGE** FACIAL APPEARANCE NORMAL SKIN NORMAL



Page 6 Ot 12 Scan to View Report



ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

S.K. Tower, Hari Niwas, LBS Marg

PATIENT NAME: SUNIL AJAGEKAR PATIENT ID: SUNIM100678181A

SRL Ltd

THANE, 400602

ACCESSION NO: 0181VI000857 AGE: 44 Years SEX: Male

DRAWN: RECEIVED: 24/09/2022 08:11 REPORTED: 26/09/2022 15:43

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units **Final** UPPER LIMB NORMAL LOWER LIMB NORMAL NECK NORMAL NECK LYMPHATICS / SALIVARY GLANDS NOT ENLARGED OR TENDER THYROID GLAND NOT ENLARGED CAROTID PULSATION NORMAL **TEMPERATURE** NORMAL

PULSE 82/MIN.REGULAR, ALL PERIPHERAL PULSES WELL FELT, NO CAROTID

BRUIT

RESPIRATORY RATE NORMAL

CARDIOVASCULAR SYSTEM

BP 150/90 MM HG mm/Hg

(SUPINE) NORMAL NORMAL

HEART SOUNDS NORMAL **MURMURS ABSENT**

RESPIRATORY SYSTEM

SIZE AND SHAPE OF CHEST NORMAL MOVEMENTS OF CHEST SYMMETRICAL BREATH SOUNDS INTENSITY NORMAL

BREATH SOUNDS QUALITY VESICULAR (NORMAL)

ADDED SOUNDS **ABSENT**

PER ABDOMEN

PERICARDIUM

APEX BEAT

APPEARANCE NORMAL VENOUS PROMINENCE ABSENT NOT PALPABLE **LIVER SPLEEN** NOT PALPABLE HERNIA **ABSENT**

CENTRAL NERVOUS SYSTEM

HIGHER FUNCTIONS NORMAL CRANIAL NERVES NORMAL CEREBELLAR FUNCTIONS NORMAL SENSORY SYSTEM NORMAL MOTOR SYSTEM NORMAL







CLIENT'S NAME AND ADDRESS:
ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI

ACCESSION NO: 0181VI000857

SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT ID:

SUNIM100678181A

Email: customercare.thane@srl.in

PATIENT NAME: SUNIL AJAGEKAR AGE: 44 Years

DRAWN: RECEIVED: 24/09/2022 08:11 REPORTED: 26/09/2022 15:43

SEX: Male

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Reference Interval Units	
REFLEXES	NORMAL		
MUSCULOSKELETAL SYSTEM			
SPINE	NORMAL		
JOINTS	NORMAL		
BASIC EYE EXAMINATION			
CONJUNCTIVA	NORMAL		
EYELIDS	NORMAL		
EYE MOVEMENTS	NORMAL		
CORNEA	NORMAL		
DISTANT VISION RIGHT EYE WITHOUT GLASSES	REDUCED VISUAL ACUITY 6/9		
DISTANT VISION LEFT EYE WITHOUT GLASSES	WITHIN NORMAL LIMIT		
NEAR VISION RIGHT EYE WITHOUT GLASSES	REDUCED VISUAL ACUITY N/8		
NEAR VISION LEFT EYE WITHOUT GLASSES	REDUCED VISUAL ACUITY N/8		
COLOUR VISION	NORMAL		
SUMMARY			
RELEVANT HISTORY	NOT SIGNIFICANT		
RELEVANT GP EXAMINATION FINDINGS	OBESE: BMI 30		
REMARKS / RECOMMENDATIONS	 WEIGHT LOSS:-STRICT DIET. 	SICIAN FOR PP CONTROL & DYSLIPIDEMIA. LOW FAT, LOW CARBOHYDRATE, HIGH FIBRE	
		GULAR WALK FOR 30-40 MIN DAILY. UROC ACID AFTER 3 MONTHS OF DIET AND	
	6) AVOID HIGH QUALITY F	PROTEIN DIET.	
	7) UROLOGY CONSULT SC 8) CARDIOLOGY CONSULT	OS FOR RENAL CALCULUS. TIN VIEW OF ECG FINDINGS.	

Interpretation(s)
BLOOD COUNTS,EDTA WHOLE BLOOD-

The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

WBC DIFFERENTIAL COUNT - NLRThe optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to

show mild disease.
(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients; A.-P. Yang, et al.; International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope.

MICROSCOPIC EXAMINATION, URINERoutine urine analysis assists in screening and diagnosis of various metabolic, urological, kidney and liver disorders







ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI **NEW DELHI 110030** DELHI INDIA 8800465156

SRL Ltd S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT ID:

SUNIM100678181A

Email: customercare.thane@srl.in

PATIENT NAME: SUNIL AJAGEKAR

ACCESSION NO: 0181VI000857 AGE: 44 Years SEX: Male

DRAWN: RECEIVED: 24/09/2022 08:11 REPORTED: 26/09/2022 15:43

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Results Biological Reference Interval Test Report Status Units <u>Final</u>

Protein: Elevated proteins car be an early sign of kidney disease. Urinary protein excretion can also be temporarily elevated by strenuous exercise, orthostatic proteinuria,

dehydration, urnary tract infections and acute illness with fever
Glucose: Uncontrolled diabetes mellitus can lead to presence of glucose in urine. Other causes include pregnancy, hormonal disturbances, liver disease and certain miedications.

Ketones: Uncontrolled diabetes mellitus car lead to presence of ketones in urine. Ketones can also be seen in starvation, frequent vomiting, pregnancy and strenuous

Blood: Occult blood can occur in urine as intact erythrocytes or haemoglobin, which can occur in various urological, nephrological and bleeding disorders.

Leukocytes: An increase in leukocytes is an indication of inflammation in urinary tract or kidneys. Most common cause is bacterial urinary tract infection. Nitrite: Many bacteria give positive results when their number is high. Nitrite concentration during infection increases with length of time the urine specimen is retained in bladder prior to collection.

pH: The kidneys play an important role in maintaining acid base balance of the body. Conditions of the body producing acidosis/ alkalosis or ingestion of certain type of food can affect the nH of urine.

Specific gravity: Specific gravity gives an indication of how concentratec the urine is. Increasec specific gravity is seen in conditions like dehydration, glycosuria and proteinum while decreased specific gravity is seen in excessive fluid intake, renal failure and diabetes insipidus. Bilirubin: In certain liver diseases such as biliary obstruction or hepatitis, bilirubin gets excreted in urine.

Urobilinogen: Positive results are seen in liver diseases like hepatitis and cirrhosis and in cases of hemolytic anemia ERYTHRO SEDIMENTATION RATE, BLOOD-

Erythrocyte sedimentation rate (ESR) is a non - specific phenomena and is clinically useful in the diagnosis and monitoring of disorders associated with an increased production of acute phase reactants. The ESR is increased in pregnancy from about the 3rd month and returns to normal by the 4th week post partum. ESR is influenced by age, sex, menstrual cycle and drugs (eg. corticosteroids, contraceptives). It is especially low (0-1mm) in polycythaemia, hypofibrinogenemia or congestive cardiac failure and when there are abnormalities of the redicells such as polkilocytosis, spherocytosis or sickle cells.

- Nathan and Oski's Haematology of Infancy and Childhood, 5th edition
 Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin
- 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th Edition" GLYCOSYLATED HEMOGLOBIN, EDTA WHOLE BLOOD-

Glycosylatec hemoglobin (GHb) has been firmly established as an index of long-term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. Formation of GHb is essentially irreversible, and the concentration in the blood depends on both the life span of the rec blood cell (average 120 days) and the blood glucose concentration. Because the rate of formation of GHb is directly proportional to the concentration of glucose in the blood, the GHb concentration represents the integrated values for glucose over the preceding 6-8 weeks.

Any condition that alters the life span of the red blood cells has the potential to alter the GHb level. Samples from patients with hemolytic anemias will exhibit decreased

glycated hemoglobin values due to the shortened life span of the red cells. This effect will depend upon the severity of the anemia. Samples from patients with polycythemia or post-spienectomy may exhibit increased glycated hemoglobin values due to a somewhat longer life span of the red cells.

Glycosylated hemoglobins results from patients with HbSS, HbCC, and HbSC and HbD must be interpreted with caution, given the pathological processes, including anemia,

increased rec cell turnover, transfusion requirements, that adversely impact HbA1c as a marker of long-term glycemic control. Ir these conditions, alternative forms of

testing such as glycated serum protein (fructosamine) should be considered.

'Targets should be individualized; More or less stringent glycemic goals may be appropriate for individual patients. Goals should be individualized based on duration of diabetes, age/life expectancy, comorbid conditions, known CVD or advanced microvascular complications, hypoglycemia unawareness, and individual patient considerations."

References

- 1. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics, edited by Carl A Burtis, Edward R. Ashwood, David E Bruns, 4th Edition, Elsevier publication, 2006, 879-884.
- Forsham PH. Diabetes Mellitus: A rational plan for management. Postgrad Med 1982, 71,139-154.
- 3. Mayer TK, Freedman ZR: Protein glycosylation in Diabetes Mellitus: A review of laboratory measurements and their clinical utility. Clin Chim Acta 1983, 127, 147-184. GLUCOSE, FASTING, PLASMA-

ADA 2021 guidelines for adults, after 8 hrs fasting is as follows:

Pre-diabetics: 100 - 125 mg/dL

Diabetic: > or = 126 mg/dL

Diabetic: > or = 126 mg/dL

GLUCOSE, POST-FRANDIAL, PLASMA-ADA Guidelines for 2hr post prandial glucose levels is only after ingestion of 75grams of glucose in 300 ml water, over a period of 5 minutes.

LIVER FUNCTION PROFILE, SERUM-LIVER FUNCTION PROFILE

Bilirubin is a yellowish pigment found in bile and is a breakdowr product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give sellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.
AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured

As its an enzymeround in various parts of the body. As its round in the liver, skelect missies, klurleys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepaticsellular injury, to determine liver health. AST levels increase during acute hepatitis, sometimes due to a viral infection, is chemia to the liver, chronic hepatitis.obstruction of bile ducts.cirrhosis.







ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI **NEW DELHI 110030** DELHI INDIA

8800465156

SRL Ltd S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: SUNIL AJAGEKAR PATIENT ID: SUNIM100678181A

ACCESSION NO: 0181VI000857 AGE: 44 Years SEX: Male

DRAWN: RECEIVED: 24/09/2022 08:11 REPORTED: 26/09/2022 15:43

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Results Biological Reference Interval Test Report Status Units <u>Final</u>

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Paget's disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Mainutrition, Protein deficiency, Wilson's disease. GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, billiary system normal ertyffile activity, setum of i has been widely used as affilined to fill the distriction. He would be activity car be round in a seeses in liver, billiary system and pancereas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc. Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myelome, a, Waldenstrom's disease. Lower-than-normal levels may be due to: Agamm aglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver iseases, Malabsorption, Mahrutrition, Nephrotic is syndrome, Protein-losing enteropathy etc. Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver diseases like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular negrous blood and appropriate the cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular negrous blood and the protein permeability or decreased lymphatic clearance, mainutrition and wasting etc SERUM BLOOD UREA NITROGEN-

Causes of Increased levels

Pre renal

• High protein diet, Increased protein databolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal

Renal Failure

• Malignancy, Nephrolithiasis, Prostatism

Causes of decreased levels

Liver disease

SIADH.
CREATININE, SERUMHigher than normal level may be due to:

Blockage in the urinary tract

- Kidney problems, such as kidney damage or failure, infection, or reduced blood flow
- Loss of body fluid (dehydration)
- Muscle problems, such as breakdown of muscle fibers
 Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:

- Myasthenia Gravis

Múscular dystrophy URIC ACID, SERUM-

Causes of Increasec levels

Dietary

- High Protein Intake.
- Prolonged Fasting,
 Rapid weight loss.

Gout

Lesch nyhan syndrome. Type 2 ĎM.

Metabolic syndrome.

Causes of decreased levels

- . Low Zinc Intake
- Multiple Sclerosis

Nutritional tips to manage increased Uric acid levels

- Drink plenty of fluids
- Limit animal proteins
- High Fibre foods
 Vit C Intake
- · Antioxidant rich foods

TOTAL PROTEIN, SERUM-

Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome,Protein-losing enteropathy etc. ALBUMIN, SERUM-

Human serum albumir is the most abundant protein in human blooc plasma. It is produced in the liver. Albumin constitutes about half of the blooc serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.



Page 10 Ot 12 Scan to View Report



ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI **NEW DELHI 110030** DELHI INDIA 8800465156

SRL Ltd S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT ID:

SUNIM100678181A

Email: customercare.thane@srl.in

PATIENT NAME: SUNIL AJAGEKAR

ACCESSION NO: 0181VI000857 AGE: 44 Years SEX: Male

DRAWN: RECEIVED: 24/09/2022 08:11 REPORTED: 26/09/2022 15:43

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Results Biological Reference Interval Test Report Status Units <u>Final</u>

ELECTROLYTES (NA/K/CL), SERUM-

Sodium levels are Increased in dehydration, cushing's syndrome, aldosteronism & decreased in Addison's disease, hypopituitarism, liver disease. Hypokalemia (low K) is common in vomiting, diarrhea, alcoholism, folic acid deficiency and primary aldosteronism. Hyperkalemia may be seen in end-stage renal failure, hemolysis, trauma, Addison's disease, metabolic acidosis, acute starvation, dehydration, and with rapid K infusion. Chloride is increased in dehydration, renal tubular acidosis (hyperchloremia metabolic acidosis), acute renal failure, metabolic acidosis associated with prolonged diarrhea and loss of sodium bidarbonate, diabetes insipidus, adrenocortical hyperfuction, salicylate intoxication and with excessive infusion of isotonic saline or extremely high dietary intake of salt. Chloride is decreased in overhydration, chronic respiratory acidosis, salt-losing nephritis, metabolic alkalosis, congestive heart failure, Addisonian crisis, certain types of metabolic acidosis, persistent gastric secretion and

respiratory actions, satisfying reprints, inetabolic arkaiosis, congestive heartrailure, Audisoniar crisis, certain types of metabolic actions, persistent gastric secretion and prolonged vomiting.

THYROID PANEL, SERUMTriodothyronine T3 , is a thyroid hormone. It affects almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate. Production of T3 and ts prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Thyroxine T4, Thyroxine's principal function is to stimulate the metabolism of all cells and tissues in the body. Excessive secretion of thyroxine in the body is hyperthyroidism, and detirent secretion is called by activity small traction of the

hyperthyroidism, and deficient secretion is called hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low. Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3

Levels in TOTAL T4 TSH3G TOTAL T3

(μIU/mL) 0.1 - 2.5 0.2 - 3.0 0.3 - 3.0 Pregnancy First Trimester (µg/dL) 6.6 - 12.4 (ng/dL) 81 - 190 6.6 - 15.5 100 - 260 100 - 260 2nd Trimester 6.6 - 15.5 3rc Trimester

Below mentioned are the guidelines for age related reference ranges for T3 and T4.

(ng/dL) (µg/dL) New Born: 75 - 260 1-3 day: 8.2 - 19.9 1 Week: 6.0 - 15.9

NOTE: TSH concentrations in apparently normal euthyroid subjects are known to be highly skewed, with a strong tailed distribution towards higher TSH values. This is well documented in the pediatric population including the infant age group.

Kindly note: Method specific reference ranges are appearing on the report under biological reference range.

- Reference:

 1. Burtis C.A., Ashwood E. R. Bruns D.E. Teitz textbook of Clinical Chemistry and Molecular Diagnostics, 4th Edition.

 2. Gowenlock A.H. Varley's Practical Clinical Biochemistry, 6th Edition.

 1. Burling Tayl Book of Padiatrics. 17th Edition
- 3. Behrman R.E. Kilegman R.M., Jenson H. B. Nelsor Text Book of Pediatrics, 17th Edition STOOL: OVA & PARASITE-

Acute infective diarrhoea and gastroenteritis (diarrhoea with vomiting) are major causes of ill health and premature death in developing countries. Loss of water and electrolytes from the body can lead to severe dehydration which if untreated, can be rapidly fatal in young children, especially that are malnourished, hypoglycaemic, and generally in poor health.

Laboratory diagnosis of parasitic infection is mainly based on microscopic examination and the gross examination of the stool specimen. Depending on the nature of the parasite, the microscopic observations include the identification of cysts, ova, trophozoites, larvae or portions of adult structure. The two classes of parasites that cause human infection are the Protozoa and Helminths. The protozoan infections include amoebiasis mainly caused by Entamoeba histolytica and giardiasis caused by Giardia lambila. The common helminthic parasites are Trichurs trichiura, Ascaris lumbricoides, Strongyloides stercoralis, Taenia sp. etc.

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD
Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in

plasma. To determine blook group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Flease note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for

The test is performed by both forward as well as reverse grouping methods.

THIS REPORT CARRIES THE SIGNATURE OF OUR LABORATORY DIRECTOR, THIS IS AN INVIOLABLE FEATURE OF OUR LAB MANAGEMENT SOFTWARE, HOWEVER, ALL EXAMINATIONS AND INVESTIGATIONS HAVE BEEN CONDUCTED BY OUR PANEL OF DOCTORS.



Page 11 Ot 12 Scan to View Report



CLIENT'S NAME AND ADDRESS:
ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL Ltd

S.K. Tower,Hari Niwas, LBS Marg THANE, 400602

MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: SUNIL AJAGEKAR

PATIENT ID: SUNIM100678181A

AGE: 44 Years ACCESSION NO: 0181VI000857 SEX: Male

RECEIVED: 24/09/2022 08:11 DRAWN: REPORTED: 26/09/2022 15:43

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units **Final**

MEDI WHEEL FULL BODY HEALTH CHECK UP ABOVE 40 MALE

ULTRASOUND ABDOMEN ULTRASOUND ABDOMEN GRADE | FATTY LIVER. RIGHT RENAL NON-OBSTRUCTING CALCULUS.

> **End Of Report** Please visit www.srlworld.com for related Test Information for this accession

Dr. Sheetal Sawant Consultant Microbiologist

Dr. Ushma Wartikar Consultant Pathologist Dr.(Mrs)Neelu K Bhojani





