

Name : MR. VARADA ABHINAY V M

Age / Gender : 31 Years / Male

Consulting Dr. :

Reg. Location

: Malad West (Main Centre)

Authenticity Check

R

E

Use a QR Code Scanner Application To Scan the Code

:30-Mar-2023 / 10:37 :30-Mar-2023 / 14:00

# AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE

Collected

Reported

CBC (Complete Blood Count), Blood			
<u>PARAMETER</u>	<u>RESULTS</u>	<b>BIOLOGICAL REF RANGE</b>	<u>METHOD</u>
RBC PARAMETERS			
Haemoglobin	13.1	13.0-17.0 g/dL	Spectrophotometric
RBC	6.41	4.5-5.5 mil/cmm	Elect. Impedance
PCV	39.6	40-50 %	Calculated
MCV	61.8	80-100 fl	Measured
MCH	20.5	27-32 pg	Calculated
MCHC	33.2	31.5-34.5 g/dL	Calculated
RDW	17.3	11.6-14.0 %	Calculated
WBC PARAMETERS			
WBC Total Count	6780	4000-10000 /cmm	Elect. Impedance
WBC DIFFERENTIAL AND ABSOLUTE COUNTS			
Lymphocytes	42.2	20-40 %	
Absolute Lymphocytes	2861.2	1000-3000 /cmm	Calculated
Monocytes	8.1	2-10 %	
Absolute Monocytes	549.2	200-1000 /cmm	Calculated
Neutrophils	41.7	40-80 %	
Absolute Neutrophils	2827.3	2000-7000 /cmm	Calculated
Eosinophils	7.5	1-6 %	
Absolute Eosinophils	508.5	20-500 /cmm	Calculated
Basophils	0.5	0.1-2 %	
Absolute Basophils	33.9	20-100 /cmm	Calculated

WBC Differential Count by Absorbance & Impedance method/Microscopy.

### **PLATELET PARAMETERS**

Platelet Count	254000	150000-400000 /cmm	Elect. Impedance
MPV	9.9	6-11 fl	Measured
PDW	21.2	11-18 %	Calculated

**RBC MORPHOLOGY** 

Immature Leukocytes



Name : MR. VARADA ABHINAY V M

Age / Gender : 31 Years / Male

Consulting Dr. : -

Reg. Location : Malad West (Main Centre)



Use a QR Code Scanner Application To Scan the Code

Collected

Reported

:30-Mar-2023 / 10:37

:30-Mar-2023 / 13:22

Hypochromia

Microcytosis ++

Macrocytosis - Anisocytosis N

Anisocytosis Mild Poikilocytosis Mild

Polychromasia -

Target Cells -

Basophilic Stippling Normoblasts -

Others Elliptocytes-occasional

+

WBC MORPHOLOGY -

PLATELET MORPHOLOGY -

COMMENT Eosinophilia

Note: Features are suggestive of thalassemia trait. Advice: Hemoglobin studies by HPLC, Reticulocyte count.

Result rechecked.

Specimen: EDTA Whole Blood

ESR, EDTA WB-ESR 5 2-15 mm at 1 hr. Sedimentation

\*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West
\*\*\* End Of Report \*\*\*







Dr.JYOT THAKKER
M.D. (PATH), DPB
Pathologist & AVP( Medical Services)

Page 2 of 11



Name : MR. VARADA ABHINAY V M

Age / Gender :31 Years / Male

Consulting Dr.

Reg. Location

: Malad West (Main Centre)

Authenticity Check

Use a QR Code Scanner Application To Scan the Code

:30-Mar-2023 / 10:37

Collected Reported :30-Mar-2023 / 14:11

# AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
GLUCOSE (SUGAR) FASTING, Fluoride Plasma	91.6	Non-Diabetic: < 100 mg/dl Impaired Fasting Glucose: 100-125 mg/dl Diabetic: >/= 126 mg/dl	Hexokinase
GLUCOSE (SUGAR) PP, Fluoride Plasma PP/R	98.7	Non-Diabetic: < 140 mg/dl Impaired Glucose Tolerance: 140-199 mg/dl Diabetic: >/= 200 mg/dl	Hexokinase
BILIRUBIN (TOTAL), Serum	0.68	0.1-1.2 mg/dl	Colorimetric
BILIRUBIN (DIRECT), Serum	0.26	0-0.3 mg/dl	Diazo
BILIRUBIN (INDIRECT), Serum	0.42	0.1-1.0 mg/dl	Calculated
TOTAL PROTEINS, Serum	7.1	6.4-8.3 g/dL	Biuret
ALBUMIN, Serum	4.7	3.5-5.2 g/dL	BCG
GLOBULIN, Serum	2.4	2.3-3.5 g/dL	Calculated
A/G RATIO, Serum	2	1 - 2	Calculated
SGOT (AST), Serum	12.9	5-40 U/L	NADH (w/o P-5-P)
SGPT (ALT), Serum	13.5	5-45 U/L	NADH (w/o P-5-P)
GAMMA GT, Serum	12.5	3-60 U/L	Enzymatic
ALKALINE PHOSPHATASE, Serum	75.7	40-130 U/L	Colorimetric
BLOOD UREA, Serum	24.7	12.8-42.8 mg/dl	Kinetic
BUN, Serum	11.5	6-20 mg/dl	Calculated
CREATININE, Serum	0.97	0.67-1.17 mg/dl	Enzymatic



Name : MR. VARADA ABHINAY V M

Age / Gender : 31 Years / Male

Consulting Dr. :

Reg. Location

: Malad West (Main Centre)

Authenticity Check

Use a QR Code Scanner Application To Scan the Code

:30-Mar-2023 / 10:37

:30-Mar-2023 / 19:17

eGFR, Serum 96 >60 ml/min/1.73sqm Calculated

Note: eGFR estimation is calculated using MDRD (Modification of diet in renal disease study group) equation

URIC ACID, Serum

3.5-7.2 mg/dl

Collected

Reported

Enzymatic

Urine Sugar (Fasting)AbsentAbsentUrine Ketones (Fasting)AbsentAbsent

5.6

Urine Sugar (PP)AbsentAbsentUrine Ketones (PP)AbsentAbsent

\*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West
\*\*\* End Of Report \*\*\*







Dr.ANUPA DIXIT

Dr.ANUPA DIXIT M.D.(PATH) Consultant Pathologist & Lab Director

Page 4 of 11



Name : MR. VARADA ABHINAY V M

Age / Gender :31 Years / Male

Consulting Dr.

Reg. Location : Malad West (Main Centre)



Authenticity Check

Use a OR Code Scanner Application To Scan the Code

: 30-Mar-2023 / 10:37

:30-Mar-2023 / 13:59

# AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE **GLYCOSYLATED HEMOGLOBIN (HbA1c)**

Collected

Reported

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
Glycosylated Hemoglobin (HbA1c), EDTA WB - CC	5.0	Non-Diabetic Level: < 5.7 % Prediabetic Level: 5.7-6.4 % Diabetic Level: >/= 6.5 %	HPLC
Estimated Average Glucose (eAG), EDTA WB - CC	96.8	mg/dl	Calculated

#### Intended use:

- In patients who are meeting treatment goals, HbA1c test should be performed at least 2 times a year
- In patients whose therapy has changed or who are not meeting glycemic goals, it should be performed quarterly
- For microvascular disease prevention, the HbA1C goal for non pregnant adults in general is Less than 7%.

#### Clinical Significance:

- HbA1c, Glycosylated hemoglobin or glycated hemoglobin, is hemoglobin with glucose molecule attached to it.
- The HbA1c test evaluates the average amount of glucose in the blood over the last 2 to 3 months by measuring the percentage of glycosylated hemoglobin in the blood.

#### Test Interpretation:

- The HbA1c test evaluates the average amount of glucose in the blood over the last 2 to 3 months by measuring the percentage of Glycosylated hemoglobin in the blood.
- HbA1c test may be used to screen for and diagnose diabetes or risk of developing diabetes.
- To monitor compliance and long term blood glucose level control in patients with diabetes.
- Index of diabetic control, predicting development and progression of diabetic micro vascular complications.

#### Factors affecting HbA1c results:

Increased in: High fetal hemoglobin, Chronic renal failure, Iron deficiency anemia, Splenectomy, Increased serum triglycerides, Alcohol ingestion, Lead/opiate poisoning and Salicylate treatment.

Decreased in: Shortened RBC lifespan (Hemolytic anemia, blood loss), following transfusions, pregnancy, ingestion of large amount of Vitamin E or Vitamin C and Hemoglobinopathies

Reflex tests: Blood glucose levels, CGM (Continuous Glucose monitoring)

References: ADA recommendations, AACC, Wallach's interpretation of diagnostic tests 10th edition.

\*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West \*\*\* End Of Report \*\*\*







Dr..JYOT THAKKER M.D. (PATH), DPB

Pathologist & AVP( Medical Services)

Page 5 of 11



Name : MR. VARADA ABHINAY V M

Age / Gender :31 Years / Male

Collected Consulting Dr. :30-Mar-2023 / 10:37 Reg. Location

:30-Mar-2023 / 15:19 : Malad West (Main Centre) Reported

# AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE **URINE EXAMINATION REPORT**

PARAMETER	RESULTS	BIOLOGICAL REF RANGE	METHOD
	<u></u>		<u></u>
PHYSICAL EXAMINATION			
Color	Pale yellow	Pale Yellow	-
Reaction (pH)	5.0	4.5 - 8.0	Chemical Indicator
Specific Gravity	1.025	1.001-1.030	Chemical Indicator
Transparency	Clear	Clear	-
Volume (ml)	20	-	-
<b>CHEMICAL EXAMINATION</b>			
Proteins	Absent	Absent	pH Indicator
Glucose	Absent	Absent	GOD-POD
Ketones	Absent	Absent	Legals Test
Blood	Trace	Absent	Peroxidase
Bilirubin	Absent	Absent	Diazonium Salt
Urobilinogen	Normal	Normal	Diazonium Salt
Nitrite	Absent	Absent	Griess Test
MICROSCOPIC EXAMINATION			
Leukocytes(Pus cells)/hpf	1-2	0-5/hpf	
Red Blood Cells / hpf	Occasional	0-2/hpf	
Epithelial Cells / hpf	0-1		
Casts	Absent	Absent	
Crystals	Absent	Absent	
Amorphous debris	Absent	Absent	
Bacteria / hpf	2-3	Less than 20/hpf	
Others	-		

Interpretation: The concentration values of Chemical analytes corresponding to the grading given in the report are as follows:

- Protein: (1+ ~25 mg/dl, 2+ ~75 mg/dl, 3+ ~ 150 mg/dl, 4+ ~ 500 mg/dl)
- Glucose:(1+ ~ 50 mg/dl, 2+ ~100 mg/dl, 3+ ~300 mg/dl,4+ ~1000 mg/dl)
- Ketone: (1 + ~5 mg/dl, 2 + ~15 mg/dl, 3 + ~50 mg/dl, 4 + ~150 mg/dl)

Reference: Pack insert

\*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West







Dr.JYOT THAKKER M.D. (PATH), DPB

Authenticity Check

Use a OR Code Scanner

Application To Scan the Code

Pathologist & AVP( Medical Services)

Page 6 of 11



Name : MR. VARADA ABHINAY V M

Age / Gender :31 Years / Male

Consulting Dr. Reg. Location : Malad West (Main Centre)



Authenticity Check

Use a QR Code Scanner Application To Scan the Code

**Collected** Reported

\*\*\* End Of Report \*\*\*



Name : MR. VARADA ABHINAY V M

Age / Gender :31 Years / Male

Consulting Dr.

Reg. Location : Malad West (Main Centre)



Use a OR Code Scanner Application To Scan the Code

: 30-Mar-2023 / 10:37 :30-Mar-2023 / 14:11

## AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE **BLOOD GROUPING & Rh TYPING**

Collected

Reported

**PARAMETER RESULTS** 

**ABO GROUP** AΒ

Rh TYPING **POSITIVE** 

NOTE: Test performed by automated column agglutination technology (CAT) which is more sensitive than conventional methods.

Specimen: EDTA Whole Blood and/or serum

ABO system is most important of all blood group in transfusion medicine

#### Limitations:

- ABO blood group of new born is performed only by cell (forward) grouping because allo antibodies in cord blood are of maternal origin.
- Since A & B antigens are not fully developed at birth, both Anti-A & Anti-B antibodies appear after the first 4 to 6 months of life. As a result, weaker reactions may occur with red cells of newborns than of adults.
- Confirmation of newborn's blood group is indicated when A & B antigen expression and the isoagglutinins are fully developed at 2 to 4 years of age & remains constant throughout life.
- Cord blood is contaminated with Wharton's jelly that causes red cell aggregation leading to false positive result
- The Hh blood group also known as Oh or Bombay blood group is rare blood group type. The term Bombay is used to refer the phenotype that lacks normal expression of ABH antigens because of inheritance of hh genotype.

### Refernces:

- Denise M Harmening, Modern Blood Banking and Transfusion Practices- 6th Edition 2012. F.A. Davis company. Philadelphia 1.
- AABB technical manual

\*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West \*\*\* End Of Report \*\*







Dr..JYOT THAKKER M.D. (PATH), DPB

Pathologist & AVP( Medical Services)

Page 8 of 11



Name : MR. VARADA ABHINAY V M

Age / Gender : 31 Years / Male

Consulting Dr. :

Reg. Location : Malad West (Main Centre)



Use a QR Code Scanner Application To Scan the Code

: 30-Mar-2023 / 10:37

:30-Mar-2023 / 14:11

# AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE LIPID PROFILE

Collected

Reported

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
CHOLESTEROL, Serum	167.1	Desirable: <200 mg/dl Borderline High: 200-239mg/dl High: >/=240 mg/dl	CHOD-POD
TRIGLYCERIDES, Serum	44.4	Normal: <150 mg/dl Borderline-high: 150 - 199 mg/dl High: 200 - 499 mg/dl Very high:>/=500 mg/dl	GPO-POD
HDL CHOLESTEROL, Serum	52.7	Desirable: >60 mg/dl Borderline: 40 - 60 mg/dl Low (High risk): <40 mg/dl	Homogeneous enzymatic colorimetric assay
NON HDL CHOLESTEROL, Serum	114.4	Desirable: <130 mg/dl Borderline-high:130 - 159 mg/dl High:160 - 189 mg/dl Very high: >/=190 mg/dl	Calculated
LDL CHOLESTEROL, Serum	105.0	Optimal: <100 mg/dl Near Optimal: 100 - 129 mg/dl Borderline High: 130 - 159 mg/dl High: 160 - 189 mg/dl Very High: >/= 190 mg/dl	Calculated
VLDL CHOLESTEROL, Serum	9.4	< /= 30 mg/dl	Calculated
CHOL / HDL CHOL RATIO, Serum	3.2	0-4.5 Ratio	Calculated
LDL CHOL / HDL CHOL RATIO,	2.0	0-3.5 Ratio	Calculated

<sup>\*</sup>Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West
\*\*\* End Of Report \*\*\*







Dr.JYOT THAKKER M.D. (PATH), DPB

Pathologist & AVP( Medical Services)

Page 9 of 11



Name : MR. VARADA ABHINAY V M

Age / Gender : 31 Years / Male

Consulting Dr. :

Reg. Location

: Malad West (Main Centre)

Authenticity Check

R

E

Use a QR Code Scanner Application To Scan the Code

:30-Mar-2023 / 10:37

**Reported** :30-Mar-2023 / 14:11

Collected

# AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE THYROID FUNCTION TESTS

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
Free T3, Serum	4.6	3.5-6.5 pmol/L	ECLIA
Free T4, Serum	12.9	11.5-22.7 pmol/L	ECLIA
sensitiveTSH, Serum	6.13	0.35-5.5 microIU/ml	ECLIA



Name : MR. VARADA ABHINAY V M

Age / Gender :31 Years / Male

Consulting Dr. Collected Reported

Reg. Location : Malad West (Main Centre)



Authenticity Check

Use a OR Code Scanner Application To Scan the Code

:30-Mar-2023 / 10:37 :30-Mar-2023 / 14:11

A thyroid panel is used to evaluate thyroid function and/or help diagnose various thyroid disorders.

#### Clinical Significance:

1)TSH Values between high abnormal upto15 microIU/ml should be correlated clinically or repeat the test with new sample as physiological

can give falsely high TSH.

2)TSH values may be trasiently altered becuase of non thyroidal illness like severe infections, liver disease, renal and heart severe burns, trauma and surgery etc.

TSH	FT4 / T4	FT3 / T3	Interpretation
High	Normal	Normal	Subclinical hypothyroidism, poor compliance with thyroxine, drugs like amiodarone, Recovery phase of non-thyroidal illness, TSH Resistance.
High	Low	Low	Hypothyroidism, Autoimmune thyroiditis, post radio iodine Rx, post thyroidectomy, Anti thyroid drugs, tyrosine kinase inhibitors & amiodarone, amyloid deposits in thyroid, thyroid tumors & congenital hypothyroidism.
Low	High	High	Hyperthyroidism, Graves disease, toxic multinodular goiter, toxic adenoma, excess iodine or thyroxine intake, pregnancy related (hyperemesis gravidarum, hydatiform mole)
Low	Normal	Normal	Subclinical Hyperthyroidism, recent Rx for Hyperthyroidism, drugs like steroids & dopamine), Non thyroidal illness.
Low	Low	Low	Central Hypothyroidism, Non Thyroidal Illness, Recent Rx for Hyperthyroidism.
High	High	High	Interfering anti TPO antibodies, Drug interference: Amiodarone, Heparin, Beta Blockers, steroids & anti epileptics.

Diurnal Variation: TSH follows a diurnal rhythm and is at maximum between 2 am and 4 am, and is at a minimum between 6 pm and 10 pm. The variation is on the order of 50 to 206%. Biological variation:19.7% (with in subject variation)

Reflex Tests:Anti thyroid Antibodies, USG Thyroid ,TSH receptor Antibody. Thyroglobulin, Calcitonin

#### Limitations:

- 1. Samples should not be taken from patients receiving therapy with high biotin doses (i.e. >5 mg/day) until atleast 8 hours following the last biotin administration.
- 2. Patient samples may contain heterophilic antibodies that could react in immunoassays to give falsely elevated or depressed results. this assay is designed to minimize interference from heterophilic antibodies.

#### Reference:

- 1.O.koulouri et al. / Best Practice and Research clinical Endocrinology and Metabolism 27(2013)
- 2.Interpretation of the thyroid function tests, Dayan et al. THE LANCET . Vol 357
- 3. Tietz , Text Book of Clinical Chemistry and Molecular Biology -5th Edition
- 4. Biological Variation: From principles to Practice-Callum G Fraser (AACC Press)

\*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West \*\*\* End Of Report \*\*\*







Dr..JYOT THAKKER M.D. (PATH), DPB

Pathologist & AVP( Medical Services)

Page 11 of 11



CID

: 2308913174

Name

: Mr VARADA ABHINAY V M

Age / Sex

: 31 Years/Male

Ref. Dr

.

Reg. Location

: Malad West Main Centre

Reg. Date

: 30-Mar-2023

Reported

: 01-Apr-2023 / 11:51

# USG WHOLE ABDOMEN

### LIVER:

The liver is normal in size, shape and smooth margins. It shows normal parenchymal echo pattern. The intra hepatic biliary and portal radical appear normal. No evidence of any intra hepatic cystic or solid lesion seen. The main portal vein and CBD appears normal.

# GALL BLADDER:

The gall bladder is physiologically distended and appears normal. No evidence of gall stones or mass lesions seen.

### PANCREAS:

The pancreas is well visualised and appears normal. No evidence of solid or cystic mass lesion.

## **KIDNEYS:**

Both the kidneys are normal in size, shape and echotexture.

No evidence of any calculus, hydronephrosis or mass lesion seen.

### SPLEEN:

The spleen is normal in size and echotexture. No evidence of focal lesion is noted.

There is no evidence of any lymphadenopathy or ascites.

# **URINARY BLADDER:**

The urinary bladder is well distended and reveal no intraluminal abnormality.

## PROSTATE:

The prostate is normal in size.

Click here to view images << ImageLink>>

REGD. OFFICE: Suburban Diagnostics (India) Pvt. Ltd., Aston, 2<sup>nd</sup> Floor, Sundervan Complex



CID

: 2308913174

Name

: Mr VARADA ABHINAY V M

Age / Sex

: 31 Years/Male

Ref. Dr

Reg. Location

: Malad West Main Centre

Reg. Date

: 30-Mar-2023

Reported

: 01-Apr-2023 / 11:51

# **IMPRESSION:**

No significant abnormality is seen.

# Suggestion: Clinicopathological correlation.

Note: Investigations have their limitations. Solitary radiological investigations never confirm the final diagnosis. They only help in diagnosing the disease in correlation to clinical symptoms and other related tests. USG is known to have inter-observer variations. Further / Follow-up imaging may be needed in some case for confirmation of findings. Patient has been explained in detail about the USG findings including its limitations and need for further imaging if clinically indicated. Please interpret accordingly. All the possible precaution have been taken under covid-19 pandemic. -----End of Report-----

Dr. Sunil Bhutka

DMRD DNB

MMC REG NO:2011051101

REGD. OFFICE: Suburban Diagnostics (India) Pvt. Ltd., Aston, 2<sup>nd</sup> Floor, S

Reported



Authenticity Check



Use a QR Code Scanner

R

E

Application To Scan the Code

: 30-Mar-2023 / 14:00

Reg. Date : 30-Mar-2023

CID

: 2308913174

Name

: Mr VARADA ABHINAY V M

Age / Sex

: 31 Years/Male

Ref. Dr

Reg. Location

: Malad West Main Centre

# X-RAY CHEST PA VIEW

Both lung fields are clear.

Both costo-phrenic angles are clear.

The cardiac size and shape are within normal limits.

The domes of diaphragm are normal in position and outlines.

The skeleton under review appears normal.

# **IMPRESSION:**

NO SIGNIFICANT ABNORMALITY IS DETECTED.

Kindly correlate clinically.

Note: Investigations have their limitations. Solitary radiological investigations never confirm the final diagnosis. X- ray is known to have interobserver variations. FThey only help in diagnosing the disease in correlation to clinical symptoms and other related tests urther / Follow-up imaging may be needed in some case for confirmation of findings. Please interpret accordingly.

-----End of Report-----

DR. Akash Chhari MBBS. MD. Radio-Diagnosis Mumbai MMC REG NO - 2011/08/2862

Click here to view images http://3.111.232.119/iRISViewer/NeoradViewer?AccessionNo=2023033009350632