

DIAGNOSTICS REPORT

Patient Name	: Mr. RAJ BEER SINGH	Order Date	: 24/01/2023 10:08
Age/Sex	: 50 Year(s)/Male	Report Date	: 24/01/2023 11:17
UHID	: SHHM.57122	IP No	:
Ref. Doctor	: Self	Facility	: SEVENHILLS HOSPITAL, MUMBAI

2D ECHOCARDIOGRAPHY WITH COLOUR DOPPLER STUDY

Normal LV and RV systolic function.

Estimated LVEF = 60%

No LV regional wall motion abnormality at rest .

All valves are structurally and functionally normal.

Normal sized cardiac chambers.

No LV Diastolic dysfunction .

No pulmonary arterial hypertension.

No regurgitation across any other valves.


Normal forward flow velocities across all the cardiac valves.

Aorta and pulmonary artery dimensions: normal.

IAS / IVS: Intact.

No evidence of clot, vegetation, calcification, pericardial effusion.

COLOUR DOPPLER: NO MR/AR.



Dr. Jayashree Dash ,

(Junior Consultant NIC)

RegNo: 3393/09/2003

LABORATORY INVESTIGATION REPORT

Patient Name : Mr. RAJ BEER SINGH	Age/Sex : 50 Year(s) / Male
UHID : SHHM.57122	Order Date : 24/01/2023 10:08
Episode : OP	Mobile No : 9991905005
Ref. Doctor : Self	DOB : 01/01/1973
	Facility : SEVENHILLS HOSPITAL, MUMBAI

Blood Bank

Test Name

Result

Sample No : O0256818A

Collection Date : 24/01/23 10:17

Ack Date : 24/01/2023 10:49

Report Date : 24/01/23 12:23

BLOOD GROUPING (ABO+RH) BY COLUMN AGGLUTINATION METHOD

BLOOD GROUP (ABO)

' B '

Rh Type

POSITIVE

REMARK :- The reported results pertain to the sample received at the blood centre.

Interpretation :

Blood typing is used to determine an individual's blood group, to establish whether a person is blood group A, B, AB, or O and whether he or she is Rh positive or Rh negative. Blood typing has the following significance,

- *Ensure compatibility between the blood type of a person who requires a transfusion of blood or blood components and the ABO and Rh type of the unit of blood that will be transfused.*
- *Determine compatibility between a pregnant woman and her developing baby (fetus). Rh typing is especially important during pregnancy because a mother and her fetus could be incompatible.*
- *Determine the blood group of potential blood donors at a collection facility.*
- *Determine the blood group of potential donors and recipients of organs, tissues, or bone marrow, as part of a workup for a transplant procedure.*

End of Report

Dr.Ritesh Kharche
MD, PGD

HOD, Laboratory Medicine Dept.

RegNo: 2006/03/1680

LABORATORY INVESTIGATION REPORT

Patient Name : Mr. RAJ BEER SINGH	Age/Sex : 50 Year(s) / Male
UHID : SHHM.57122	Order Date : 24/01/2023 10:08
Episode : OP	Mobile No : 9991905005
Ref. Doctor : Self	DOB : 01/01/1973
	Facility : SEVENHILLS HOSPITAL, MUMBAI

Biochemistry

Test Name	Result	Unit	Ref. Range
Sample No : O0256818A	Collection Date : 24/01/23 10:17	Ack Date : 24/01/2023 10:30	Report Date : 24/01/23 12:15

GLYCOSYLATED

HAEMOGLOBIN (HBA1C)

HbA1c	7.91 ▲	%	4 to 6% Non-diabetic 6.0--7.0% Excellent control 7.0--8.0% Fair to good control 8.0--10% Unsatisfactory control ABOVE 10% Poor control
-------	---------------	---	--

Method - BIOCHEMISTRY

Estimated Average Glucose (eAG)	180.32 ▲	mg/dl	90 - 126
------------------------------------	-----------------	-------	----------

Method - Calculated

NOTES :-

1. HbA1c is used for monitoring diabetic control. It reflects the mean plasma glucose over three months
2. HbA1c may be falsely low in diabetics with hemolytic disease. In these individuals a plasma fructosamine level may be used which evaluates diabetes over 15 days.
3. Inappropriately low HbA1c values may be reported due to hemolysis, recent blood transfusion, acute blood loss, hypertriglyceridemia, chronic liver disease. Drugs like dapsone, ribavirin, antiretroviral drugs, trimethoprim, may also cause interference with estimation of HbA1c, causing falsely low values.
4. HbA1c may be increased in patients with polycythemia or post-splenectomy.
5. Inappropriately higher values of HbA1c may be caused due to iron deficiency, vitamin B12 deficiency, alcohol intake, uremia, hyperbilirubinemia and large doses of aspirin.
6. Trends in HbA1c are a better indicator of diabetic control than a solitary test.
7. Any sample with >15% HbA1c should be suspected of having a hemoglobin variant, especially in a non-diabetic patient. Similarly, below 4% should prompt additional studies to determine the possible presence of variant hemoglobin.
8. HbA1c target in pregnancy is to attain level <6 % .
9. HbA1c target in paediatric age group is to attain level < 7.5 %.

Method : turbidimetric inhibition immunoassay (TINIA) for hemolyzed whole blood

Reference : American Diabetes Associations. Standards of Medical Care in Diabetes 2015

Sample No : O0256818C	Collection Date : 24/01/23 10:17	Ack Date : 24/01/2023 11:38	Report Date : 24/01/23 12:54
-----------------------	----------------------------------	-----------------------------	------------------------------

Lipid Profile

LABORATORY INVESTIGATION REPORT

Patient Name : Mr. RAJ BEER SINGH UHID : SHHM.57122 Episode : OP Ref. Doctor : Self	Age/Sex : 50 Year(s) / Male Order Date : 24/01/2023 10:08 Mobile No : 9991905005 DOB : 01/01/1973 Facility : SEVENHILLS HOSPITAL, MUMBAI
--	---

Total Cholesterol	250	mg/dl	Reference Values : Up to 200 mg/dL - Desirable 200-239 mg/dL - Borderline High >240 mg/dL - High
Triglycerides	331	mg/dl	Reference Values: Up to 150 mg/dL - Normal 150-199 mg/dL - Borderline High 200-499 mg/dL - High >500 mg/dL - Very High
<i>Method - Enzymatic</i>			
HDL Cholesterol	43.8	mg/dl	0 - 60
<i>Method - Enzymatic immuno inhibition</i>			
LDL Cholesterol	140.00 ▲	mg/dl	0 - 130
<i>Method - Calculated</i>			
VLDL Cholesterol	66.20 ▲	mg/dl	0 - 40
<i>Method - Calculated</i>			
Total Cholesterol / HDL Cholesterol Ratio - Calculated	5.71 ▲	RATIO	0 - 5
<i>Method - Calculated</i>			
LDL / HDL Cholesterol Ratio - Calculated	3.20	RATIO	0 - 4.3
<i>Method - Calculated</i>			

LABORATORY INVESTIGATION REPORT

Patient Name : Mr. RAJ BEER SINGH	Age/Sex : 50 Year(s) / Male
UHID : SHHM.57122	Order Date : 24/01/2023 10:08
Episode : OP	
Ref. Doctor : Self	Mobile No : 9991905005
	DOB : 01/01/1973
	Facility : SEVENHILLS HOSPITAL, MUMBAI

References:

- 1) Pack Insert of Bio system
- 2) Tietz Textbook Of Clinical Chemistry And Molecular Diagnostics, 6th Ed, Editors: Rifai et al. 2018

Interpretation

1. Triglycerides: When triglycerides are very high greater than 1000 mg/dL, there is a risk of developing pancreatitis in children and adults. Triglycerides change dramatically in response to meals, increasing as much as 5 to 10 times higher than fasting levels just a few hours after eating. Even fasting levels vary considerably day to day. Therefore, modest changes in fasting triglycerides measured on different days are not considered to be abnormal.
2. HDL-Cholesterol: HDL- C is considered to be beneficial, the so-called "good" cholesterol, because it removes excess cholesterol from tissues and carries it to the liver for disposal. If HDL-C is less than 40 mg/dL for men and less than 50 mg/dL for women, there is an increased risk of heart disease that is independent of other risk factors, including the LDL-C level. The NCEP guidelines suggest that an HDL cholesterol value greater than 60 mg/dL is protective and should be treated as a negative risk factor.
3. LDL-Cholesterol: Desired goals for LDL-C levels change based on individual risk factors. For young adults, less than 120 mg/dL is acceptable. Values between 120-159 mg/dL are considered Borderline high. Values greater than 160 mg/dL are considered high. Low levels of LDL cholesterol may be seen in people with an inherited lipoprotein deficiency and in people with hyperthyroidism, infection, inflammation, or cirrhosis.

Uric Acid (Serum)

Uric Acid	5.3	mg/dl	3.5 - 7.2
-----------	-----	-------	-----------

Method - Uricase

References:

- 1) Pack Insert of Bio system
- 2) Tietz Textbook of Clinical chemistry and Molecular Diagnostics Edited by: Carl A. burtis, Edward R. Ashwood, David e. Bruns

Interpretation:-

Uric acid is produced by the breakdown of purines. Purines are nitrogen-containing compounds found in the cells of the body, including our DNA. Increased concentrations of uric acid can cause crystals to form in the joints, which can lead to the joint inflammation and pain characteristic of gout. Low values can be associated with some kinds of liver or kidney diseases, Fanconi syndrome, exposure to toxic compounds, and rarely as the result of an inherited metabolic defect (Wilson disease).

Liver Function Test (LFT)

SGOT (Aspartate Transaminase) - SERUM	23.5	U/L	0 - 35
<i>Method - IFCC</i>			
SGPT (Alanine Transaminase) - SERUM	36.0	U/L	0 - 45
<i>Method - IFCC</i>			
Total Bilirubin - SERUM	0.77	mg/dl	0 - 2
<i>Method - Diazo</i>			
Direct Bilirubin - - SERUM	0.31	mg/dl	0 - 0.4
<i>Method - Diazotization</i>			
Indirect Bilirubin - Calculated	0.46	mg/dl	0.1 - 0.8
<i>Method - Calculated</i>			
Alkaline Phosphatase - SERUM	72	U/L	0 - 115

LABORATORY INVESTIGATION REPORT

Patient Name : Mr. RAJ BEER SINGH	Age/Sex : 50 Year(s) / Male
UHID : SHHM.57122	Order Date : 24/01/2023 10:08
Episode : OP	
Ref. Doctor : Self	Mobile No : 9991905005
	DOB : 01/01/1973
	Facility : SEVENHILLS HOSPITAL, MUMBAI

Method - IFCC AMP Buffer

Total Protein - SERUM 6.76 gm/dl 6 - 7.8

Method - Biuret

Albumin - SERUM 4.46 gm/dl 3.5 - 5.2

Method - Bromo Cresol Green(BCG)

Globulin - Calculated 2.30 gm/dl 2 - 4

Method - Calculated

A:G Ratio 1.94 :1 1 - 3

Method - Calculated

Gamma Glutamyl 23.4 U/L 0 - 55

Transferase (GGT) -
Gglutamyl carboxy
nitroanilide - SERUM

Method - G glutamyl carboxy nitroanilide

References:

1) Pack Insert of Bio system

2) Tietz Textbook Of Clinical Chemistry And Molecular Diagnostics, 6th Ed, Editors: Rifai et al. 2018

Interperatation :-

Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Elevated levels results from increased bilirubin production (eg hemolysis and ineffective erythropoiesis); decreased bilirubin excretion (eg; obstruction and hepatitis); and abnormal bilirubin metabolism (eg; hereditary and neonatal jaundice). conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts tumors & Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of hemolytic or pernicious anemia, transfusion reaction & a common metabolic condition termed Gilbert syndrome.

AST levels increase in viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. Ast levels may also increase after a heart attack or strenuous activity. ALT is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health. Elevated ALP levels are seen in Biliary Obstruction, Osteoblastic Bone Tumors, Osteomalacia, Hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, paget's disease, Rickets, Sarcoidosis etc.

Elevated serum GGT activity can be found in diseases of the liver, Biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-including drugs etc.

Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum.. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease. Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic - Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.

Renal Function Test (

RFT)

Urea - SERUM 25 mg/dl 15 - 39

Method - Urease

BUN - SERUM 11.68 mg/dl 4 - 18

Method - Urease-GLDH

Creatinine - SERUM 0.87 mg/dl 0.5 - 1.3

Method - Jaffes Kinetic

LABORATORY INVESTIGATION REPORT

Patient Name : Mr. RAJ BEER SINGH

UHID : SHHM.57122

Episode : OP

Ref. Doctor : Self

Age/Sex : 50 Year(s) / Male

Order Date : 24/01/2023 10:08

Mobile No : 9991905005

DOB : 01/01/1973

Facility : SEVENHILLS HOSPITAL, MUMBAI

American Diabetes Association Reference Range :

Post-Prandial Blood Glucose:

Non- Diabetic: Up to 140mg/dL

Pre-Diabetic: 140-199 mg/dL

Diabetic : >200 mg/dL

References:

1)Pack Insert of Bio system

2) Tietz Textbook Of Clinical Chemistry And Molecular Diagnostics, 6th Ed, Editors: Rifai et al. 2018

Interpretation :-

Conditions that can result in an elevated blood glucose level include: Acromegaly, Acute stress (response to trauma, heart attack, and stroke for instance), Chronic kidney disease, Cushing syndrome, Excessive consumption of food, Hyperthyroidism, Pancreatitis.

A low level of glucose may indicate hypoglycemia, a condition characterized by a drop in blood glucose to a level where first it causes nervous system symptoms (sweating, palpitations, hunger, trembling, and anxiety), then begins to affect the brain (causing confusion, hallucinations, blurred vision, and sometimes even coma and death). A low blood glucose level (hypoglycemia) may be seen with: Adrenal insufficiency, Drinking excessive alcohol, Severe liver disease, Hypopituitarism, Hypothyroidism, Severe infections, Severe heart failure, Chronic kidney (renal) failure, Insulin overdose, Tumors that produce insulin (insulinomas), Starvation.

End of Report



Dr. Ritesh Kharche

MD, PGD

HOD, Laboratory Medicine Dept.

RegNo: 2006/03/1680

LABORATORY INVESTIGATION REPORT

Patient Name : Mr. RAJ BEER SINGH	Age/Sex : 50 Year(s) / Male
UHID : SHHM.57122	Order Date : 24/01/2023 10:08
Episode : OP	Mobile No : 9991905005
Ref. Doctor : Self	DOB : 01/01/1973
	Facility : SEVENHILLS HOSPITAL, MUMBAI

HAEMATOLOGY

Test Name	Result	Unit	Ref. Range
Sample No : O0256818A	Collection Date : 24/01/23 10:17	Ack Date : 24/01/2023 10:30	Report Date : 24/01/23 10:50

COMPLETE BLOOD COUNT (CBC) - EDTA WHOLE BLOOD

Total WBC Count	5.28	x10 ³ /ul	4.00 - 10.00
Neutrophils	59.9	%	40.00 - 80.00
Lymphocytes	32.4	%	20.00 - 40.00
Eosinophils	1.3	%	1.00 - 6.00
Monocytes	6.0	%	2.00 - 10.00
Basophils	0.4 ▼	%	1.00 - 2.00
Absolute Neutrophils Count	3.16	x10 ³ /ul	2.00 - 7.00
Absolute Lymphocytes Count	1.71	x10 ³ /ul	0.80 - 4.00
Absolute Eosinophils Count	0.07	x10 ³ /ul	0.02 - 0.50
Absolute Monocytes Count	0.32	x10 ³ /ul	0.12 - 1.20
Absolute Basophils Count	0.02	x10 ³ /ul	0.00 - 0.10
RBCs	5.32	x10 ⁶ /ul	4.50 - 5.50
Haemoglobin	15.8	gm/dl	13.00 - 17.00
Hematocrit	49.0	%	40.00 - 50.00
MCV	92.0	fl	83.00 - 101.00
MCH	29.7	pg	27.00 - 32.00
MCHC	32.3	gm/dl	31.50 - 34.50
RED CELL DISTRIBUTION WIDTH-CV (RDW-CV)	12.1	%	11.00 - 16.00
RED CELL DISTRIBUTION WIDTH-SD (RDW-SD)	40.9	fl	35.00 - 56.00
Platelet	200	x10 ³ /ul	150.00 - 410.00
MPV	11.9	fl	6.78 - 13.46
PLATELET DISTRIBUTION WIDTH (PDW)	16.4	%	9.00 - 17.00
PLATELETCRIT (PCT)	0.238	%	0.11 - 0.28

LABORATORY INVESTIGATION REPORT

Patient Name : Mr. RAJ BEER SINGH	Age/Sex : 50 Year(s) / Male
UHID : SHHM.57122	Order Date : 24/01/2023 10:08
Episode : OP	Mobile No : 9991905005
Ref. Doctor : Self	DOB : 01/01/1973
	Facility : SEVENHILLS HOSPITAL, MUMBAI

NOTE: Wallach's Interpretation of Diagnostic Tests. 11th Ed, Editors: Rao LV. 2021

NOTE :-

The International Council for Standardization in Haematology (ICSH) recommends reporting of absolute counts of various WBC subsets for clinical decision making. This test has been performed on a fully automated 5 part differential cell counter which counts over 10,000 WBCs to derive differential counts. A complete blood count is a blood panel that gives information about the cells in a patient's blood, such as the cell count for each cell type and the concentrations of Hemoglobin and platelets. The cells that circulate in the bloodstream are generally divided into three types: white blood cells (leukocytes), red blood cells (erythrocytes), and platelets (thrombocytes). Abnormally high or low counts may be physiological or may indicate disease conditions, and hence need to be interpreted clinically.

ERYTHROCYTE

SEDIMENTATION RATE

(ESR)

ESR 18 mm/hr 0 - 20

Method: Westergren Method

INTERPRETATION :-

ESR is a non-specific phenomenon, its measurement is clinically useful in disorders associated with an increased production of acute-phase proteins. It provides an index of progress of the disease in rheumatoid arthritis or tuberculosis, and it is of considerable value in diagnosis of temporal arteritis and polymyalgia rheumatica. It is often used if multiple myeloma is suspected, but when the myeloma is non-secretory or light chain, a normal ESR does not exclude this diagnosis.

An elevated ESR occurs as an early feature in myocardial infarction. Although a normal ESR cannot be taken to exclude the presence of organic disease, the vast majority of acute or chronic infections and most neoplastic and degenerative diseases are associated with changes in the plasma proteins that increased ES values. An increased ESR in subjects who are HIV seropositive seems to be an early predictive marker of progression toward acquired immune deficiency syndrome (AIDS).

The ESR is influenced by age, stage of the menstrual cycle and medications taken (corticosteroids, contraceptive pills). It is especially low (0-1 mm) in polycythaemia, hypofibrinogenaemia and congestive cardiac failure and when there are abnormalities of the red cells such as poikilocytosis, spherocytosis, or sickle cells. In cases of performance enhancing drug intake by athletes the ESR values are generally lower than the usual value for the individual and as a result of the increase in haemoglobin (i.e. the effect of secondary polycythaemia).

End of Report



Dr. Ritesh Kharche

MD, PGD

HOD, Laboratory Medicine Dept.

RegNo: 2006/03/1680

LABORATORY INVESTIGATION REPORT

Patient Name : Mr. RAJ BEER SINGH

Age/Sex : 50 Year(s) / Male

UHID : SHHM.57122

Order Date : 24/01/2023 10:08

Episode : OP

Ref. Doctor : Self

Mobile No : 9991905005

DOB : 01/01/1973

Facility : SEVENHILLS HOSPITAL, MUMBAI

Stool Examination

Test Name

Result

Sample No : O0256830D

Collection Date : 24/01/23 11:16

Ack Date : 24/01/2023 11:19

Report Date : 24/01/23 13:35

Gross and Chemical

Examination

Consistency

Semi-Solid

COLOUR STOOL

Brown

Visible Blood

Absent

Mucus

Absent ▲

Occult Blood

NEGATIVE

Microscopic

Examination

Puscells

OCCASIONAL

RBC

ABSENT

Epithelial Cells

ABSENT

Parasites

Not Seen

Bacteria

Present

End of Report

Dr.Ritesh Kharche
MD, PGD

HOD, Laboratory Medicine Dept.

RegNo: 2006/03/1680

LABORATORY INVESTIGATION REPORT

Patient Name : Mr. RAJ BEER SINGH	Age/Sex : 50 Year(s) / Male
UHID : SHHM.57122	Order Date : 24/01/2023 10:08
Episode : OP	Mobile No : 9991905005
Ref. Doctor : Self	DOB : 01/01/1973
	Facility : SEVENHILLS HOSPITAL, MUMBAI

IMMUNOLOGY

Test Name	Result	Unit	Ref. Range
Sample No : O0256818C	Collection Date : 24/01/23 10:17	Ack Date : 24/01/2023 11:38	Report Date : 24/01/23 12:46

PSA -TOTAL-SERUM

PSA- Prostate Specific Antigen - SERUM	1.45	ng/ml	0.00 - 4.00
--	------	-------	-------------

*Biological Reference Interval :-
Conventional for all ages: <=4
60 - 69 yrs: 0 - 4.5*

Note : Change in method and Reference range

INTERPRETATION :

Prostate-specific antigen (PSA) is a glycoprotein that is produced by the prostate gland, the lining of the urethra, and the bulbourethral gland. PSA exists in serum mainly in two forms, complexed to alpha-1-anti-chymotrypsin (PSA-ACT complex) and unbound (free PSA). Increases in prostatic glandular size and tissue damage caused by benign prostatic hypertrophy, prostatitis, or prostate cancer may increase circulating PSA levels. Transient increase in PSA can also be seen following per rectal digital or sonological examinations.

NOTE:

Patients on Biotin supplement may have interference in some immunoassays. With individuals taking high dose Biotin (more than 5 mg per day) supplements, at least 8-hour wait time before blood draw is recommended.

Ref: Arch Pathol Lab Med—Vol 141, November 2017

T3 - SERUM	80.52	ng/dl	70.00 - 204.00
<i>Method - CLIA</i>			
T4 - SERUM	6.35	ug/dL	4.60 - 10.50
<i>Method - CLIA</i>			
TSH - SERUM	2.49	uIU/ml	0.40 - 4.50
<i>Method - CLIA</i>			

LABORATORY INVESTIGATION REPORT

Patient Name : Mr. RAJ BEER SINGH

Age/Sex : 50 Year(s) / Male

UHID : SHHM.57122

Order Date : 24/01/2023 10:08

Episode : OP

Ref. Doctor : Self

Mobile No : 9991905005

DOB : 01/01/1973

Facility : SEVENHILLS HOSPITAL, MUMBAI

Reference Ranges (T3) Pregnancy:

First Trimester 81 - 190

Second Trimester & Third Trimester 100 - 260

Reference Ranges (TSH) Pregnancy:

1st Trimester : 0.1 - 2.5

2nd Trimester : 0.2 - 3.0

3rd Trimester : 0.3 - 3.0

Reference:

1. Clinical Chemistry and Molecular Diagnostics, Tietz Fundamentals, 7th Edition & Endocrinology Guidelines

Interpretation :-

It is recommended that the following potential sources of variation should be considered while interpreting thyroid hormone results:

1. Thyroid hormones undergo rhythmic variation within the body this is called circadian variation in TSH secretion: Peak levels are seen between 2-4 am. Minimum levels seen between 6-10 am. This variation may be as much as 50% thus, influence of sampling time needs to be considered for clinical interpretation.

2. Circulating forms of T3 and T4 are mostly reversibly bound with Thyroxine binding globulins (TBG), and to a lesser extent with albumin and Thyroid binding PreAlbumin. Thus the conditions in which TBG and protein levels alter such as chronic liver disorders, pregnancy, excess of estrogens, androgens, anabolic steroids and glucocorticoids may cause misleading total T3, total T4 and TSH interpretations.

3. Total T3 and T4 levels are seen to have physiological rise during pregnancy and in patients on steroid treatment.

4. T4 may be normal the presence of hyperthyroidism under the following conditions : T3 thyrotoxicosis, Hypoproteinemia related reduced binding, during intake of certain drugs (eg Phenytoin, Salicylates etc)

5. Neonates and infants have higher levels of T4 due to increased concentration of TBG

6. TSH levels may be normal in central hypothyroidism, recent rapid correction of hypothyroidism or hyperthyroidism, pregnancy, phenytoin therapy etc.

7. TSH values of <0.03 uIU/mL must be clinically correlated to evaluate the presence of a rare TSH variant in certain individuals which is undetectable by conventional methods.

8. Presence of Autoimmune disorders may lead to spurious results of thyroid hormones

9. Various drugs can lead to interference in test results.

10. It is recommended that evaluation of unbound fractions, that is free T3 (fT3) and free T4 (fT4) for clinic-pathologic correlation, as these are the metabolically active forms.

End of Report

Dr. Ritesh Kharche

MD, PGD

HOD, Laboratory Medicine Dept.

RegNo: 2006/03/1680


ID : 2301240003
 Name : rajbeer singh
 Sex : Male
 Age : 50
 Divisions:
 Hospital No. :
 Hospital: seven hills hospital
 DataTime: 2023-01-24 10:44
 Height : cm
 Weight : kg
 BP : mmHg
 Bed No. :

HR 84 bpm
 P Dur/PR int 110/160ms
 QRS Dur 93 ms
 QT/QTc int 332/393 ms
 P/QRS/T axis 60/37/46°
 RV5/SV1 amp 0.766/0.885mV
 RV5+SV1 amp 1.651mV
 RV6/SV2 amp 0.516/0.981mV

Minnesota Code
 9-1-1(I, II, III)
 9-4-1(V3)
 Diagnosis Info
 800 Sinus Rhythmm
 131 Low Voltage(Limb Leads)

Handwritten signature





SEVEN HILLS HOSPITAL

MUMBAI
MUMBAI

Mr. RAJBEER SINGH
ID : 46947
DATE : 24-01-2023
AGE/SEX : 50 /M
HT/WT : 173 / 75
REF.BY : Self

TREADMILL TEST REPORT

PROTOCOL : Bruce
HISTORY : NIL
INDICATION : Routine
MEDICATION : NIL

PHASE	TOTAL TIME	STAGE TIME	SPEED Km/Hr	GRADE %	H.R. bpm	B.P. mmHg	RPP x100	ST LEVEL (MM)		METS
								II	V1	
SUPINE					107	126 / 80	134	3.1	0.7	3.7
STANDING					101	126 / 80	127	2	0.3	2.2
HYPERTENT		0:41			105	126 / 80	132	1.7	0.4	2.2
Stage 1	2:55	2:55	2.7	10	139	126 / 80	175	2.4	-0.4	4.67
Stage 2	5:55	2:55	4	12	152	130 / 90	197	2	-0.1	7.04
PK-EXERCISE	6:10	0:10	5.4	14	155	130 / 90	201	1.5	-0.1	7.26
RECOVERY	9:20	2:55			121	130 / 90	157	0.4	0.5	0.7
RECOVERY	10:19	3:54			110	110 / 70	121	0.8	0.3	1

RESULTS

EXERCISE DURATION : 6:10
MAX HEART RATE : 155 bpm 91 % of target heart rate 170 bpm
MAX BLOOD PRESSURE : 130 / 90 mm Hg
REASON OF TERMINATION : THR ACHIEVED
MAX WORK LOAD : 7.26 METS

BP RESPONSE :
ARRHYTHMIA :
H.R. RESPONSE :

IMPRESSIONS

GOOD EFFORT TOLERANCE.
NORMAL CHRONOTROPIC AND IONOTROPIC RESPONSES.
NO ANGINA / ARRHYTHMIA.
NO ST - T CHANGES.
STRESS TEST IS NEGATIVE FOR INDUCIBLE ISCHAEMIA.

Technician : VIKESH JADHAV

Dr. Jayshree Dash

LABORATORY INVESTIGATION REPORT

Patient Name : Mr. RAJ BEER SINGH	Age/Sex : 50 Year(s) / Male
UHID : SHHM.57122	Order Date : 24/01/2023 10:08
Episode : OP	Mobile No : 9991905005
Ref. Doctor : Self	DOB : 01/01/1973
	Facility : SEVENHILLS HOSPITAL, MUMBAI

Urinalysis

Test Name	Result	Unit	Ref. Range
Sample No : O0256820D	Collection Date : 24/01/23 10:21	Ack Date : 24/01/2023 10:54	Report Date : 24/01/23 13:40

Physical Examination

QUANTITY	50	ml	
Colour	Pale Yellow		
Appearance	Clear		
DEPOSIT	Absent		Absent
pH	Acidic		
Specific Gravity	1.020		

Chemical Examination

Protein	Absent		Absent
Sugar	POSITIVE (+)		Absent
ketones	Absent		Absent
Occult Blood	NEGATIVE		Absent
Bile Salt	Absent		Absent
Bile Pigments	Absent		Absent
Urobilinogen	NORMAL		Absent
NITRATE	Absent		
LEUKOCYTES	Absent		

Microscopic Examination

Puscells	OCCASIONAL	/HPF	
Epithelial Cells	OCCASIONAL	/HPF	
RBC	ABSENT	/HPF	Absent
Cast	Absent	/LPF	Absent
Crystal	Absent	/HPF	Absent
Amorphous Materials	Absent		Absent
Yeast	Absent		Absent
Bacteria	Absent		Absent

URINE SUGAR AND KETONE (FASTING)

Sugar	POSITIVE (+)
ketones	Absent

Sample No : O0256837E	Collection Date : 24/01/23 12:35	Ack Date : 24/01/2023 12:37	Report Date : 24/01/23 13:36
-----------------------	----------------------------------	-----------------------------	------------------------------

LABORATORY INVESTIGATION REPORT

Patient Name : Mr. RAJ BEER SINGH

Age/Sex : 50 Year(s) / Male

UHID : SHHM.57122

Order Date : 24/01/2023 10:08

Episode : OP

Ref. Doctor : Self

Mobile No : 9991905005

DOB : 01/01/1973

Facility : SEVENHILLS HOSPITAL, MUMBAI

URINE SUGAR AND KETONE (PP)

Sugar
ketones

POSITIVE (+++)

Absent

End of Report



Dr. Ritesh Kharche

MD, PGD

HOD, Laboratory Medicine Dept.

RegNo: 2006/03/1680

URINE SUGAR AND KETONE (FASTING)- Report has been amended at Jan 24 2023 1:36PM by Ritesh kharche.

DIAGNOSTICS REPORT

Patient Name	: Mr. RAJ BEER SINGH	Order Date	: 24/01/2023 10:08
Age/Sex	: 50 Year(s)/Male	Report Date	: 24/01/2023 13:11
UHID	: SHHM.57122	IP No	:
Ref. Doctor	: Self	Facility	: SEVENHILLS HOSPITAL, MUMBAI

USG ABDOMEN

FINDINGS:

Liver is normal in size (13.7 cm) and echotexture. No focal liver parenchymal lesion is seen.

Intrahepatic portal and biliary radicles are normal.

Gall-bladder is physiologically distended. No evidence of intraluminal calculus is seen. Wall thickness appears normal. No evidence of peri-cholecystic fluid is seen.

Portal vein and CBD are normal in course and calibre.

Pancreas is obscured due to excessive bowel gases.

Spleen is normal in size (8.6 cm) and echotexture. No focal lesion is seen in the spleen.

Right kidney measures 11 x 4.2 cm.

Left kidney measures 10.9 x 6.7 cm.

Both the kidneys are normal in size, shape and echotexture. Cortico-medullary differentiation is maintained. No evidence of calculus or hydronephrosis on either side.


Urinary bladder is well distended and appears normal. No evidence of intra-luminal calculus or mass lesion.

Prostate appears normal in size and echotexture. It measures 3.8 x 3.4 x 3.3 cm corresponding to 23.4 cc.

There is no free fluid in abdomen and pelvis.

IMPRESSION:

No significant abnormality is detected.



Dr. Shubham Asrani

Dr. Shubham Asrani, MBBS, MD

RegNo: 2020/01/0042

DIAGNOSTICS REPORT

Patient Name	: Mr. RAJ BEER SINGH	Order Date	: 24/01/2023 10:08
Age/Sex	: 50 Year(s)/Male	Report Date	: 24/01/2023 13:52
UHID	: SHHM.57122	IP No	:
Ref. Doctor	: Self	Facility	: SEVENHILLS HOSPITAL, MUMBAI

X-RAY CHEST PA VIEW

Both lungs are clear.

The frontal cardiac dimensions are normal.


The pleural spaces are clear.

Both hilar shadows are normal in position and density.

No diaphragmatic abnormality is seen.

The soft tissues and bony thorax are normal.

IMPRESSION: No pleuroparenchymal lesion is seen.



Dr. Rashmi Randive, MBBS, MD