PID No. : MED111551700 **Register On** : 25/03/2023 9:18 AM : 712309945 SID No. **Collection On** : 25/03/2023 10:09 AM

Age / Sex : 32 Year(s) / Female

Type : OP Ref. Dr : MediWheel

Report On : 27/03/2023 9:30 AM : 27/03/2023 12:38 PM Printed On



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HAEMATOLOGY			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood'Spectrophotometry)	13.8	g/dL	12.5 - 16.0
INTERPRETATION: Haemoglobin values vary in Mer blood loss, renal failure etc. Higher values are often due			
PCV (Packed Cell Volume) / Haematocrit (EDTA Blood/Derived)	40.2	%	37 - 47
RBC Count (EDTA Blood/Automated Blood cell Counter)	5.03	mill/cu.mm	4.2 - 5.4
MCV (Mean Corpuscular Volume) (EDTA Blood/Derived from Impedance)	80.0	fL	78 - 100
MCH (Mean Corpuscular Haemoglobin) (EDTA Blood/Derived)	27.4	pg	27 - 32
MCHC (Many Campus and a Harman lab.)	24.2	~/dI	22 26

RBC Count (EDTA Blood/Automated Blood cell Counter)	5.03	mill/cu.mm	4.2 - 5.4
MCV (Mean Corpuscular Volume) (EDTA Blood/Derived from Impedance)	80.0	fL	78 - 100
MCH (Mean Corpuscular Haemoglobin) (EDTA Blood/Derived)	27.4	pg	27 - 32
MCHC (Mean Corpuscular Haemoglobin concentration) (EDTA Blood/Derived)	34.3	g/dL	32 - 36
RDW-CV	13.8	%	11.5 - 16.0
(Derived)			
RDW-SD	38.64	fL	39 - 46
(Derived)			
Total WBC Count (TC) (EDTA Blood/Derived from Impedance)	11770	cells/cu.mm	4000 - 11000
Remark: Kindly correlate clinically.			
Neutrophils (Blood/Impedance Variation & Flow Cytometry)	65	%	40 - 75
Lymphocytes (Blood/Impedance Variation & Flow Cytometry)	27	%	20 - 45







 PID No.
 : MED111551700
 Register On
 : 25/03/2023 9:18 AM

 SID No.
 : 712309945
 Collection On
 : 25/03/2023 10:09 AM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Eosinophils (Blood/Impedance Variation & Flow Cytometry)	02	%	01 - 06
Monocytes (Blood/Impedance Variation & Flow Cytometry)	06	%	01 - 10
Basophils (Blood/Impedance Variation & Flow Cytometry)	00	%	00 - 02
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	7.65	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	3.18	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.24	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.71	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.00	10^3 / μl	< 0.2
Platelet Count (EDTA Blood/Derived from Impedance)	217	10^3 / μl	150 - 450
MPV (Blood/ <i>Derived</i>)	14.2	fL	8.0 - 13.3
PCT	0.31	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood/Automated ESR analyser)	10	mm/hr	< 20

: 27/03/2023 12:38 PM







 PID No.
 : MED111551700
 Register On
 : 25/03/2023 9:18 AM

 SID No.
 : 712309945
 Collection On
 : 25/03/2023 10:09 AM

 Age / Sex
 : 32 Year(s) / Female
 Report On
 : 27/03/2023 9:30 AM

Report On : 27/03/2023 9:30 AM

Type : OP **Printed On** : 27/03/2023 12:38 PM

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulfanilic Acid)	0.5	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.1	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.40	mg/dL	0.1 - 1.0
Total Protein (Serum/Biuret)	7.7	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.6	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	3.10	gm/dL	2.3 - 3.6
A : G Ratio (Serum/Derived)	1.48		1.1 - 2.2
INTERPRETATION: Remark : Electrophoresis is the p	preferred method		
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC / Kinetic)	15	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC / Kinetic)	22	U/L	5 - 41
Alkaline Phosphatase (SAP) (Serum/PNPP / Kinetic)	88	U/L	42 - 98
GGT(Gamma Glutamyl Transpeptidase)	34	U/L	< 38





(Serum/IFCC / Kinetic)





 PID No.
 : MED111551700
 Register On
 : 25/03/2023 9:18 AM

 SID No.
 : 712309945
 Collection On
 : 25/03/2023 10:09 AM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/Oxidase / Peroxidase method)	202	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol phosphate oxidase / peroxidase)	201	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

: 27/03/2023 12:38 PM

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the õusualö"circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	35	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/Calculated)	126.8	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	40.2	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	167.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219







Very High: >= 220

PID No. Register On : MED111551700 : 25/03/2023 9:18 AM : 712309945 SID No. Collection On : 25/03/2023 10:09 AM

Age / Sex : 32 Year(s) / Female Report On : 27/03/2023 9:30 AM

Printed On

Type : OP

(Serum/Calculated)

Ref. Dr : MediWheel



Investigation	<u>Observed</u> <u>Unit</u>	<u>Biological</u>
	Value	Reference Interval

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

: 27/03/2023 12:38 PM

Total Cholesterol/HDL Cholesterol Ratio (Serum/Calculated)	5.8	Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
Triglyceride/HDL Cholesterol Ratio (TG/HDL)	5.7	Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0

Optimal: 0.5 - 3.0 LDL/HDL Cholesterol Ratio 3.6 Borderline: 3.1 - 6.0 (Serum/Calculated)

High Risk: > 6.0

High Risk: > 5.0







 PID No.
 : MED111551700
 Register On
 : 25/03/2023 9:18 AM

 SID No.
 : 712309945
 Collection On
 : 25/03/2023 10:09 AM

Type : OP

Ref. Dr : MediWheel



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/ <i>HPLC</i>)	6.3	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4

: 27/03/2023 12:38 PM

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Printed On

Remark: Kindly correlate clinically.

Estimated Average Glucose 134.11 mg/dl

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.









Diabetic: >= 6.5

PID No. : MED111551700 : 712309945 SID No. Collection On : 25/03/2023 10:09 AM

Age / Sex : 32 Year(s) / Female Report On : 27/03/2023 9:30 AM

Printed On

Type : OP

Ref. Dr : MediWheel Register On : 25/03/2023 9:18 AM

: 27/03/2023 12:38 PM

Investigation	Observed	<u>Unit</u>	Biological
<u>investigation</u>	<u>Obscived</u>	OTTIL	
	<u>Value</u>		Reference Interval

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.19 0.7 - 2.04ng/ml

(Serum/Chemiluminescent Immunometric Assay

(CLIA))

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

Microg/dl 4.2 - 12.0T4 (Thyroxine) - Total 11.97

(Serum/Chemiluminescent Immunometric Assay

(CLIA))

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 2.259 μIU/mL 0.35 - 5.50

(Serum/Chemiluminescent Immunometric Assay

(CLIA))

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values&lt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.





APPROVED BY

 PID No.
 : MED111551700
 Register On
 : 25/03/2023 9:18 AM

 SID No.
 : 712309945
 Collection On
 : 25/03/2023 10:09 AM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	<u>Observed</u> <u>Ur</u>	<u>Biological</u>
	Value	Reference Interval

: 27/03/2023 12:38 PM

CLINICAL PATHOLOGY

PHYSICAL EXAMINATION

Colour Pale Yellow Yellow to Amber

(Urine/Physical examination)

Volume 30 ml

(Urine/Physical examination)

Appearance Clear

(Urine)

CHEMICAL EXAMINATION

pH 6.5 4.5 - 8.0

(Urine)

Specific Gravity 1.020 1.002 - 1.035

(Urine/Dip Stick o''Reagent strip method)

Protein Negative Negative

(Urine/Dip Stick oʻ'Reagent strip method)

Glucose Nil Nil

(Urine)

Ketone Nil Nil

(Urine/Dip Stick ó"Reagent strip method)

Leukocytes Negative leuco/uL Negative

(Urine)

Nitrite Nil Nil

(Urine/Dip Stick ó"Reagent strip method)

Bilirubin Negative mg/dL Negative

(Urine)

Blood Nil Nil

(Urine)

Urobilinogen Normal Within normal limits

(Urine/Dip Stick o''Reagent strip method)





PID No. : MED111551700

SID No. : 712309945

Age / Sex : 32 Year(s) / Female

Type : OP

Ref. Dr : MediWheel

Register On : 25/03/2023 9:18 AM

Collection On : 25/03/2023 10:09 AM

Report On : 27/03/2023 9:30 AM

Printed On : 27/03/2023 12:38 PM



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Urine Microscopy Pictures</u>			
RBCs (Urine/Microscopy)	Nil	/hpf	NIL
Pus Cells (Urine/Microscopy)	4-5	/hpf	< 5
Epithelial Cells (Urine/Microscopy)	4-5	/hpf	No ranges
Others (Urine)	Nil		Nil





APPROVED BY

PID No. : MED111551700 **Register On** : 25/03/2023 9:18 AM

: 712309945 SID No. Collection On : 25/03/2023 10:09 AM

Age / Sex : 32 Year(s) / Female Report On : 27/03/2023 9:30 AM

Printed On

Type : OP

Ref. Dr : MediWheel

Investigation **Observed** <u>Unit</u> **Biological** Reference Interval **Value**

: 27/03/2023 12:38 PM

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING 'O' 'Positive'

(EDTA Blood/Agglutination)

Remark: Test to be confirmed by Gel method.







PID No. Register On : MED111551700 : 712309945 SID No. Collection On : 25/03/2023 10:09 AM

Printed On

Age / Sex : 32 Year(s) / Female Report On

Ref. Dr : MediWheel

: OP

Type

: 25/03/2023 9:18 AM

: 27/03/2023 12:38 PM

27/03/2023 9:30 AM



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	11.5		
Glucose Fasting (FBS) (Plasma - F/GOD- POD)	93	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Urine sugar, Fasting (Urine - F)	Nil		Nil
Glucose Postprandial (PPBS) (Plasma - PP/GOD - POD)	96	mg/dL	70 - 140

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Sugar (PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	8.1	mg/dL	7.0 - 21
Creatinine (Serum/Jaffe Kinetic)	0.7	mg/dL	0.6 - 1.1

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetylcyteine, chemotherapeutic agent such as flucytosine

Uric Acid 5.3 mg/dL 2.6 - 6.0

(Serum/Uricase/Peroxidase)







APPROVED BY

-- End of Report --



Name	POOJA O N	ID	MED111551700
Age & Gender	32Y/F	Visit Date	Mar 25 2023 9:18AM
Ref Doctor	MediWheel		

X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

Impression: No significant abnormality detected.

DR. MOHAN. B

(DMRD, DNB, EDIR, FELLOW IN CARDIAC

MRI)

CONSULTANT RADIOLOGIST