PID No.
 : MED111017412
 Register On
 : 12/03/2022 10:14 AM

 SID No.
 : 422019562
 Collection On
 : 12/03/2022 11:56 AM

 Age / Sex
 : 49 Year(s) / Female
 Report On
 : 14/03/2022 10:32 AM

Printed On

: 19/03/2022 4:57 PM

Type : OP

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HAEMATOLOGY			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood'Spectrophotometry)	10.5	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood/Derived from Impedance)	34.6	%	37 - 47
RBC Count (EDTA Blood/Impedance Variation)	4.91	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (EDTA Blood/Derived from Impedance)	71.0	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood/Derived from Impedance)	21.4	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood/Derived from Impedance)	30.4	g/dL	32 - 36
RDW-CV (EDTA Blood/Derived from Impedance)	17.2	%	11.5 - 16.0
RDW-SD (EDTA Blood/Derived from Impedance)	42.74	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood/Impedance Variation)	13700	cells/cu.mm	4000 - 11000
Neutrophils (EDTA Blood/Impedance Variation & Flow Cytometry)	60.5	%	40 - 75
Lymphocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	33.4	%	20 - 45



VERIFIED BY



 Name
 : Ms. RAJINI K

 PID No.
 : MED111017412
 Register On
 : 12/03/2022 10:14 AM

 SID No.
 : 422019562
 Collection On
 : 12/03/2022 11:56 AM

 Age / Sex
 : 49 Year(s) / Female
 Report On
 : 14/03/2022 10:32 AM

 Type
 : OP
 Printed On
 : 19/03/2022 4:57 PM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Eosinophils (EDTA Blood/Impedance Variation & Flow Cytometry)	1.1	%	01 - 06
Monocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	4.1	%	01 - 10
Basophils (Blood/Impedance Variation & Flow Cytometry)	0.9	%	00 - 02
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	8.29	10^3 / μ1	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	4.58	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.15	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.56	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.12	10^3 / μl	< 0.2
Platelet Count (EDTA Blood/Impedance Variation)	443	$10^3 / \mu l$	150 - 450
MPV (EDTA Blood/Derived from Impedance)	9.1	fL	8.0 - 13.3
PCT (EDTA Blood/Automated Blood cell Counter)	0.40	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood/Modified Westergren)	13	mm/hr	< 20



VERIFIED BY



 PID No.
 : MED111017412
 Register On
 : 12/03/2022 10:14 AM

 SID No.
 : 422019562
 Collection On
 : 12/03/2022 11:56 AM

 Age / Sex
 : 49 Year(s) / Female
 Report On
 : 14/03/2022 10:32 AM

 Type
 : OP
 Printed On
 : 19/03/2022 4:57 PM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulfanilic Acid)	0.3	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.2	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.1	mg/dL	0.1 - 1.0
Total Protein (Serum/Biuret)	7.0	gm/dL	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.2	gm/dL	3.5 - 5.2
Globulin (Serum/Derived)	2.8	gm/dL	2.3 - 3.6
A : G Ratio (Serum/Derived)	1.5		1.1 - 2.2
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC Kinetic)	15	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC / Kinetic)	20	U/L	5 - 41
Alkaline Phosphatase (SAP) (Serum/IFCC Kinetic)	144	U/L	42 - 98
GGT(Gamma Glutamyl Transpeptidase) (Serum/SZASZ standarised IFCC)	61	U/L	< 38



VERIFIED BY



 PID No.
 : MED111017412
 Register On
 : 12/03/2022 10:14 AM

 SID No.
 : 422019562
 Collection On
 : 12/03/2022 11:56 AM

 Age / Sex
 : 49 Year(s) / Female
 Report On
 : 14/03/2022 10:32 AM

Printed On

Type : OP

Ref. Dr : MediWheel

: 19/03/2022 4:57 PM

Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/Cholesterol oxidase/Peroxidase)	164	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol phosphate oxidase / peroxidase)	142	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the õusualö circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	41	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/Calculated)	94.6	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	28.4	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	123.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220



VERIFIED BY



 PID No.
 : MED111017412
 Register On
 : 12/03/2022 10:14 AM

 SID No.
 : 422019562
 Collection On
 : 12/03/2022 11:56 AM

 Age / Sex
 : 49 Year(s) / Female
 Report On
 : 14/03/2022 10:32 AM

Printed On

Type : OP

Ref. Dr : MediWheel

<u>Investigation</u> <u>Observed</u> <u>Unit</u> <u>Biological</u> <u>Value</u> <u>Reference Interval</u>

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2. It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

: 19/03/2022 4:57 PM

Total Cholesterol/HDL Cholesterol Ratio 4 Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 4.5 - 11.0 High Risk: 4.5 - 11.0

Triglyceride/HDL Cholesterol Ratio 3.5 Optimal: < 2.5 (TG/HDL) Mild to moderate risk: 2.5 - 5.0

(Serum/Calculated) High Risk: > 5.0

LDL/HDL Cholesterol Ratio

(Serum/Calculated)

2.3

Optimal: 0.5 - 3.0

Borderline: 3.1 - 6.0

High Risk: > 6.0



VERIFIED BY



PID No. : MED111017412 Register On : 12/03/2022 10:14 AM : 422019562 SID No. Collection On : 12/03/2022 11:56 AM Age / Sex : 49 Year(s) / Female Report On : 14/03/2022 10:32 AM

Type : OP

Printed On : 19/03/2022 4:57 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/ <i>HPLC</i>)	8.0	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 182.9 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.



VERIFIED BY



PID No. : MED111017412 Register On : 12/03/2022 10:14 AM : 422019562 SID No. Collection On : 12/03/2022 11:56 AM

Age / Sex : 49 Year(s) / Female Report On : 14/03/2022 10:32 AM : OP **Type**

Printed On

Ref. Dr : MediWheel

Investigation <u>Observed</u> <u>Unit</u> **Biological** Value Reference Interval

: 19/03/2022 4:57 PM

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.14 ng/mL 0.7 - 2.04

(Serum/CMIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is

Metabolically active.

4.2 - 12.0T4 (Thyroxine) - Total 12.42 μg/dL

(Serum/CMIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

µIU/mL 0.35 - 5.50TSH (Thyroid Stimulating Hormone) 4.58

(Serum/Chemiluminescent Microparticle

Immunoassay(CMIA))

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester: 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.

2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.

3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.





APPROVED BY

Page 7 of 11

 PID No.
 : MED111017412
 Register On
 : 12/03/2022 10:14 AM

 SID No.
 : 422019562
 Collection On
 : 12/03/2022 11:56 AM

 Age / Sex
 : 49 Year(s) / Female
 Report On
 : 14/03/2022 10:32 AM

 Type
 : OP
 Printed On
 : 19/03/2022 4:57 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	Value		Reference Interval

Pale Yellow

CLINICAL PATHOLOGY

PHYSICAL EXAMINATION

(Urine)
Appearance Clear Cle

Appearance Clear Clear (Urine)

Volume 15 mL

(Urine)

CHEMICAL EXAMINATION(Automated-

<u>Urineanalyser)</u>

pH 6.0 4.5 - 8.0

(Urine/AUTOMATED URINANALYSER)

Specific Gravity 1.020 1.002 - 1.035

(Urine)

Colour

Ketones Negative Negative

(Urine)

Urobilinogen 0.2 0.2 - 1.0

(Urine/AUTOMATED URINANALYSER)

Blood Negative Negative

(Urine/AUTOMATED URINANALYSER)

Nitrite Negative Negative Negative

(Urine/AUTOMATED URINANALYSER)

Bilirubin Negative Negative

(Urine/AUTOMATED URINANALYSER)

Protein Negative Negative

(Urine)





 PID No.
 : MED111017412
 Register On
 : 12/03/2022 10:14 AM

 SID No.
 : 422019562
 Collection On
 : 12/03/2022 11:56 AM

 Age / Sex
 : 49 Year(s) / Female
 Report On
 : 14/03/2022 10:32 AM

 Type
 : OP
 Printed On
 : 19/03/2022 4:57 PM

Ref. Dr : MediWheel

Investigation Glucose (Urine)	Observed Value Negative	<u>Unit</u>	Biological Reference Interval Negative
Leukocytes (Urine)	Negative	leuco/uL	Negative
MICROSCOPY(URINE DEPOSITS)			
Pus Cells (Urine/Flow cytometry)	0-2	/hpf	3-5
Epithelial Cells (Urine)	0-2	/hpf	1-2
RBCs (Urine/Flow cytometry)	Nil	/hpf	2-3
Others (Urine)	Nil		Nil
Casts (Urine/Flow cytometry)	Nil	/hpf	0 - 1
Crystals (Urine)	Nil		NIL





PID No. Register On : 12/03/2022 10:14 AM : MED111017412 : 422019562 SID No. Collection On : 12/03/2022 11:56 AM

Age / Sex : 49 Year(s) / Female Report On : 14/03/2022 10:32 AM : OP Type

Printed On

Ref. Dr : MediWheel

<u>Unit</u> **Investigation** <u>Observed</u> **Biological** <u>Value</u> Reference Interval

: 19/03/2022 4:57 PM

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING 'A' 'Positive'

(EDTA Blood/Agglutination)



VERIFIED BY



 PID No.
 : MED111017412
 Register On
 : 12/03/2022 10:14 AM

 SID No.
 : 422019562
 Collection On
 : 12/03/2022 11:56 AM

 Age / Sex
 : 49 Year(s) / Female
 Report On
 : 14/03/2022 10:32 AM

 Type
 : OP
 Printed On
 : 19/03/2022 4:57 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	17		6 - 22
Glucose Fasting (FBS) (Plasma - F/GOD - POD)	80	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose Fasting - Urine	Negative		Negative
(Urine - F)			
Glucose Postprandial (PPBS)	54	mg/dL	70 - 140

(Plasma - PP/GOD - POD)

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Glucose Postprandial - Urine (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease-GLDH)	14	mg/dL	7.0 - 21
Creatinine (Serum/Jaffe Kinetic)	0.8	mg/dL	0.6 - 1.1

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine etc.

Uric Acid 5.5 mg/dL 2.6 - 6.0

(Serum/Uricase/Peroxidase)



VERIFIED BY



APPROVED BY

-- End of Report --