

Name	MRS.MANORAMA TIWARI	ID	MED120841310
Age & Gender	53/FEMALE	Visit Date	13/02/2023
Ref Doctor Name	MediWheel		

X-ray mammogram (mediolateral oblique & craniocaudal views) followed by Sonomammography.

## **BILATERAL MAMMOGRAPHY**

Breast composition Type B (These are scattered areas of fibroglandular density).

No evidence of focal soft tissue lesion.

No evidence of cluster microcalcification.

Subcutaneous fat deposition is within normal limits.

#### **BILATERAL SONOMAMMOGRAPHY**

Both the breasts show normal echopattern.

No evidence of focal solid / cystic areas.

Few prominent ducts are seen in the retroareolar region, largest measuring 1.5 mm in maximum diameter. No solid components / internal echoes.

Bilateral axillary lymphnodes are noted with preserved fatty hilum.

#### **IMPRESSION:**

- Prominent ducts in bilateral retroareolar region. No other breast lesions.
- Bilateral benign axillary lymph nodes.

## **ASSESSMENT: BI-RADS CATEGORY -2**

## **BI-RADS CLASSIFICATION**

## **CATEGORY RESULT**

## 2 Benign finding. Routine mammogram in 1 year recommended.

#### REPORT DISCLAIMER

- 1. This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- $10. \\ Reports are subject to interpretation in their entirety, partial or selective interpretation may lead to false opinion.$
- $11. Disputes, if any \ , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only. \\$



Name	MRS.MANORAMA TIWARI	ID	MED120841310
Age & Gender	53/FEMALE	Visit Date	13/02/2023
Ref Doctor Name	MediWheel		

# DR. HEMANANDINI V.N CONSULTANT RADIOLOGIST Hn/mj

#### REPORT DISCLAIMER

- 1.This is only a radiologincal imperssion.Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11. Disputes, if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.

 PID No.
 : MED120841310
 Register On
 : 13/02/2023 8:10 AM

 SID No.
 : 522302256
 Collection On
 : 13/02/2023 8:53 AM

 Age / Sex
 : 53 Year(s) / Female
 Report On
 : 13/02/2023 3:09 PM

**Printed On** 

: 21/02/2023 11:51 AM



Type : OP

Ref. Dr : MediWheel

Investigation  HAEMATOLOGY	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> Reference Interval
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood/Spectrophotometry)	11.6	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	36.0	%	37 - 47
RBC Count (EDTA Blood)	4.11	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (EDTA Blood)	87.4	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood)	28.2	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood)	32.2	g/dL	32 - 36
RDW-CV	14.4	%	11.5 - 16.0
RDW-SD	44.05	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood)	5300	cells/cu.mm	4000 - 11000
Neutrophils (Blood)	50.5	%	40 - 75
Lymphocytes (Blood)	34.5	%	20 - 45
Eosinophils (Blood)	5.9	%	01 - 06



 PID No.
 : MED120841310
 Register On
 : 13/02/2023 8:10 AM

 SID No.
 : 522302256
 Collection On
 : 13/02/2023 8:53 AM

 Age / Sex
 : 53 Year(s) / Female
 Report On
 : 13/02/2023 3:09 PM

**Printed On** 



Ref. Dr : MediWheel

: OP

Type

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Monocytes (Blood)	8.2	%	01 - 10
Basophils (Blood)	0.9	%	00 - 02
INTERPRETATION: Tests done on Automated Five P	art cell counter. All	abnormal results are	reviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood)	2.68	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood)	1.83	10^3 / μ1	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood)	0.31	10^3 / μ1	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood)	0.43	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood)	0.05	10^3 / μl	< 0.2
Platelet Count (EDTA Blood)	258	10^3 / μl	150 - 450
MPV (Blood)	9.4	fL	8.0 - 13.3
PCT (Automated Blood cell Counter)	0.24	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood)	7	mm/hr	< 30

: 21/02/2023 11:51 AM



: MediWheel

 PID No.
 : MED120841310
 Register On
 : 13/02/2023 8:10 AM

 SID No.
 : 522302256
 Collection On
 : 13/02/2023 8:53 AM

 Age / Sex
 : 53 Year(s) / Female
 Report On
 : 13/02/2023 3:09 PM

**Printed On** 



Type : OP

Ref. Dr

Investigation BIOCHEMISTRY	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	0.27	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.15	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.12	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	23.70	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	44.80	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	48.30	U/L	< 38
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	85.6	U/L	53 - 141
Total Protein (Serum/Biuret)	7.14	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.65	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	2.49	gm/dL	2.3 - 3.6
A : G RATIO	1.87		1.1 - 2.2

: 21/02/2023 11:51 AM

Dr Anusha.K.S Sr.Consultant Pathologist Reg No : 100674 APPROVED BY

(Serum/Derived)

PID No. : MED120841310 Register On : 13/02/2023 8:10 AM : 522302256 SID No.

Age / Sex : 53 Year(s) / Female Report On : 13/02/2023 3:09 PM

**Printed On** 

**Type** : OP

Ref. Dr : MediWheel Collection On : 13/02/2023 8:53 AM

: 21/02/2023 11:51 AM



Investigation **Piological** Limit

<u>investigation</u>	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	153.72	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	93.86	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual\_ circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	50.13	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/Calculated)	84.8	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	18.8	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	103.6	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220



Type : OP

PID No.

SID No.

Printed On : 21/02/2023 11:51 AM

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	Value		Reference Interval

**INTERPRETATION:** 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2. It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio

(Serum/Calculated)

3.1

Optimal: < 3.3

Low Risk: 3.4 - 4.4

Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio 1.9 Optimal: < 2.5

(TG/HDL)

(Serum/Calculated)

LDL/HDL Cholesterol Ratio 1.7

(Serum/Calculated)

Mild to moderate risk: 2.5 - 5.0

High Risk: > 5.0

Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0

High Risk: > 6.0



Type : OP

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)  HbA1C (Whole Blood/HPLC)	6.2	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4
(			Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

**Printed On** 

Estimated Average Glucose 131.24 mg/dL

(Whole Blood)

#### **INTERPRETATION: Comments**

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.

Dr Anusha.K.S Sr.Consultant Pathologist Reg No : 100674

: 21/02/2023 11:51 AM

> : MED120841310 Register On : 13/02/2023 8:10 AM

> > **Printed On**

: 522302256 SID No. Collection On : 13/02/2023 8:53 AM

Age / Sex : 53 Year(s) / Female Report On : 13/02/2023 3:09 PM

Type : OP

Ref. Dr : MediWheel

PID No.



<u>Investigation</u>	<u>Observed</u> <u>Un</u>	<u>it</u> <u>Biological</u>
	Value	Reference Interval

: 21/02/2023 11:51 AM

# **IMMUNOAS**SAY

#### THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.12 ng/ml 0.4 - 1.81

(Serum/ECLIA)

#### INTERPRETATION:

#### **Comment:**

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is

Metabolically active.

4.2 - 12.0T4 (Tyroxine) - Total 9.50 μg/dl

(Serum/ECLIA)

#### INTERPRETATION:

#### **Comment:**

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

9.44 μIU/mL 0.35 - 5.50TSH (Thyroid Stimulating Hormone)

(Serum/ECLIA)

#### INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester: 0.3-3.0

(Indian Thyroid Society Guidelines)

#### **Comment:**

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



 PID No.
 : MED120841310
 Register On
 : 13/02/2023 8:10 AM

 SID No.
 : 522302256
 Collection On
 : 13/02/2023 8:53 AM

 Age / Sex
 : 53 Year(s) / Female
 Report On
 : 13/02/2023 3:09 PM



Type : OP

OP **Printed On** : 21/02/2023 11:51 AM

Ref. Dr : MediWheel

<u>Investigation</u>	<u>Observed</u> <u>l</u>	<u>Jnit</u> <u>Biological</u>
	Value	Reference Interval

# **CLINICAL PATHOLOGY**

# PHYSICAL EXAMINATION (URINE COMPLETE)

Colour Yellow Yellow Yellow to Amber

(Urine)

Appearance Clear Clear

(Urine)

Volume(CLU) 25

(Urine)

# CHEMICAL EXAMINATION (URINE

**COMPLETE**)

pH 5.0 4.5 - 8.0

(Urine)

Specific Gravity 1.013 1.002 - 1.035

(Urine)

Ketone Negative Negative

(Urine)

Urobilinogen Normal Normal

(Urine)

Blood Negative Negative

Nitrite Negative

(Urine)

(Urine)

Bilirubin Negative Negative

(Urine)

(Urine)

Protein Negative Negative

DR SHAMIM JAVED
MD PATHOLOGY
KMC 88902

**APPROVED BY** 

Negative

PID No. : MED120841310 Register On : 13/02/2023 8:10 AM : 522302256 SID No. Collection On : 13/02/2023 8:53 AM Age / Sex : 53 Year(s) / Female Report On : 13/02/2023 3:09 PM



Type : OP

(Urine)

**Printed On** : 21/02/2023 11:51 AM

Ref. Dr : MediWheel

Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glucose (Urine/GOD - POD)	Negative		Negative
Leukocytes(CP) (Urine)  MICROSCOPIC EXAMINATION (URINE COMPLETE)	Negative		Negative
Pus Cells (Urine)	0-2	/hpf	NIL
Epithelial Cells (Urine)	0-1	/hpf	NIL
RBCs (Urine)	NIL	/HPF	NIL
Others	NIL		

INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.



**Type** : OP **Printed On** : 21/02/2023 11:51 AM

Ref. Dr : MediWheel

PID No.



InvestigationObservedUnitBiologicalValueReference Interval

# **IMMUNOHAEMATOLOGY**

BLOOD GROUPING AND Rh TYPING 'B' 'Positive'

(EDTA Blood/Agglutination)

INTERPRETATION: Note: Slide method is screening method. Kindly confirm with Tube method for transfusion.

Dr Anusha.K.S Sr.Consultant Pathologist Reg No : 100674

PID No. : MED120841310

: 522302256

Age / Sex : 53 Year(s) / Female

**Type** : OP

SID No.

Ref. Dr : MediWheel Register On : 13/02/2023 8:10 AM

Collection On : 13/02/2023 8:53 AM

Report On : 13/02/2023 3:09 PM

**Printed On** : 21/02/2023 11:51 AM



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<b>BIOCHEMISTRY</b>			
BUN / Creatinine Ratio	17.5		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	103.31	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine)	Negative		Negative
(Urine - F/GOD - POD)			
Glucose Postprandial (PPBS)	116.28	mg/dL	70 - 140
(Plasma - PP/GOD-PAP)			

**INTERPRETATION:** 

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Blood Urea Nitrogen (BUN)	12.6	mg/dL	7.0 - 21
(Serum/ <i>Urease UV / derived</i> )			
Creatinine	0.72	mg/dL	0.6 - 1.1
(Serum/Modified Jaffe)			

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetylcyteine, chemotherapeutic agent such as flucytosine

Uric Acid 2.85 2.6 - 6.0mg/dL (Serum/Enzymatic)



**APPROVED BY** 

-- End of Report --