

.

1

V1	V2	V3	. V4
TH TARUN KUMAR le 'Dr. MMHSRATH.S. MBBS Years kg Regd. No: 41964 DDRC SRL, KANNI Utrun normal Und	S,MD,DNB		
2	V6	Standard	
: 111 ms : 151 ms IS : 81 ms C/QTe : 352/396 ms QRS/T : 48/64/29 ° V5/SV1 : 2.002/1.257 mV			




ECHOCARDIOGRAPHY REPORT


Name:	KETHA TARUN KUMAR	Age: 30	Date:	11/03/2023
Ref By:	BANK OF BARODA	Sex: M	SRD No:	WC000958
M MODE	2D ECHO		MV/A	Area : Normal
AORTA : 29 mm	MV : Norma	al	PV: N	lormal
LA : 32 mm	AV: Normal	1	TV: N	ormal
LVIDD: 47 mm	RWMA: Nil		RA: N	lormal
LVIDS : 33 mm	LA: Normal		RV: N	lormal
IVSD: 10 mm	LV: Normal		IVS: N	lormal
IVSS : 12 mm	IAS: Intact		A-V R	elationship : Normal
LVPWD:11 mm	Situs: Solitu	IS		Vessel Relationship: Norma
LVPWS: 13 mm	V-A Relation	nship: Normal		onary V Drainage: Normal
LVEF : 60 %	Syst. V. Dra	inage: Normal		
FS: 30 %	Pericardial	Effusion: Nil		
DOPPLER				
Pul Velocity : 1.0	m/s			
MV Velocity :	E : 0.81 m/s	A:0.71 m/s	F/A ·	1.14 MV Area (PHT):
AV Velocity :1.16		1.0.71 11/5	цл.	1.14 WY Area (FIII).
AO Area :				
TV Velocity : 0.6	m/s	RVSP : mmHg		
revelocity . 0.0	11/3	NV5F . Milling		
COLOUR				
MR: Trivial	ASD:			
AR: Nil	VSD:			
TR: Nil	PDA			
PR:Nil	CoA:			
	ormalities : Nil			

Wall motion abnormalities : Nil	
Pericardium: No pericardial effusion	
Vegetation/ Thrombus: Nil	

IMPRESSION:

- Trivial MR
- No RWMA
- Normal LV Systolic function











CLIENT CODE : CA00010147 - MEDIWHEEL CLIENT'S NAME AND ADDRESS ! THOADE I MATTER

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED F701A, LADO SARAI, NEW DELHI, SOUTH DELHI, DELHI, SOUTH DELHI 110030 DELHI INDIA 8800465156

DDRC SRL DIAGNOSTICS


KANNUR KERALA, INDIA Tel: 93334 93334 Email : customercare.ddrc@srl.in

PATIENT NAME : KETHA TARUN	KUMAR	PATIENT ID : KETHM1008924053
ACCESSION NO : 4053WC000958	AGE : 30 Years SEX : Male	ABHA NO :
DRAWN :	RECEIVED : 11/03/2023 09:26	REPORTED : 11/03/2023 17:56
REFERRING DOCTOR : DR. MEDIWH	IEEL	CLIENT PATIENT ID :
Test Report Status <u>Final</u>	Results	Biological Reference Interval Units
MEDIWHEEL HEALTH CHECKUP	BELOW 40(M)2DECHO	
OPTHAL		
OPTHAL	COMPLETED	
PHYSICAL EXAMINATION		

COMPLETED PHYSICAL EXAMINATION

















CLIENT CODE : CA00010147 - MEDIWHEEL CLIENT'S NAME AND ADDRESS ! THOADE I MATTER

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED F701A, LADO SARAI, NEW DELHI, SOUTH DELHI, DELHI, SOUTH DELHI 110030 DELHI INDIA 8800465156


DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334 Email : customercare.ddrc@srl.in

PATIENT NAME : KETHA TARUN KUMAR		PATIENT ID : KETHM1008924053
ACCESSION NO : 4053WC000958 AGE : 30 Ye	ears SEX : Male	ABHA NO :
DRAWN : RECEIVED :	11/03/2023 09:26	REPORTED : 11/03/2023 17:56
REFERRING DOCTOR : DR. MEDIWHEEL		CLIENT PATIENT ID :
Test Report Status <u>Final</u>	Results	Units
DIRECT LDL CHOLESTEROL	110	Optimum : < 100 mg/dL Above Optimum : 100-139 Borderline High : 130-159 High : 160-189 Very High : >or= 190
NON HDL CHOLESTEROL	119	Desirable-Less than 130 mg/dL Above Desirable-130-159 Borderline High-160-189 High-190-219 Very High- >or =220
VERY LOW DENSITY LIPOPROTEIN	11.3	= 30.0 mg/dL</td
CHOL/HDL RATIO	3.1 Lov	 3.3 - 4.4 Low Risk 4.5 - 7.0 Average Risk 7.1 - 11.0 Moderate Risk > 11.0 High Risk
LDL/HDL RATIO	1.9	0.5-3 Desirable/Low risk 3.1-6 Borderline/Moderate risk >6.0 High Risk







Test Report Status <u>Final</u> Results Units

Interpretation(s)

1) Cholesterol levels help assess the patient risk status and to follow the progress of patient under treatment to lower serum cholesterol concentrations.

2) Serum Triglyceride (TG) are a type of fat and a major source of energy for the body. Both quantity and composition of the diet impact on plasma triglyceride concentrations. Elevations in TG levels are the result of overproduction and impaired clearance. High TG are associated with increased risk for CAD (Coronary artery disease) in patients with other risk factors, such as low HDL-C, some patient groups with elevated apolipoprotein B concentrations, and patients with forms of LDL that may be particularly atherogenic.

3)HDL-C plays a crucial role in the initial step of reverse cholesterol transport, this considered to be the primary atheroprotective function of HDL

4) LDL -C plays a key role in causing and influencing the progression of atherosclerosis and, in particular, coronary sclerosis. The majority of cholesterol stored in atherosclerotic plaques originates from LDL, thus LDL-C value is the most powerful clinical predictor.

5)Non HDL cholesterol: Non-HDL-C measures the cholesterol content of all atherogenic lipoproteins, including LDL hence it is a better marker of risk in both primary and secondary prevention studies. Non-HDL-C also covers, to some extent, the excess ASCVD risk imparted by the sdLDL, which is significantly more atherogenic than the normal large buoyant particles, an elevated non-HDL-C indirectly suggests greater proportion of the small, dense variety of LDL particles

Serum lipid profile is measured for cardiovascular risk prediction.Lipid Association of India recommends LDL-C as primary target and Non HDL-C as co-primary treatment target.

Risk Category				
Extreme risk group	A.CAD with > 1 feature of high risk group			
	B. CAD with > 1 feature of Very high risk group or recurrent ACS (within 1 year) despite LDL-C			
	< or $=$ 50 mg/dl or polyvascular disease			
Very High Risk		major risk factors or evidence of end organ damage 3.		
	Familial Homozygous Hypercholesterolemi			
High Risk		abetes with 1 major risk factor or no evidence of end		
		DL >190 mg/dl 5. Extreme of a single risk factor. 6.		
		7. Lipoprotein a >/= 50mg/dl 8. Non stenotic carotid		
	plaque	seelt and see		
Moderate Risk	2 major ASCVD risk factors			
Low Risk	0-1 major ASCVD risk factors			
Major ASCVD (Ath	erosclerotic cardiovascular disease) Risk Fa	ictors		
1. Age $>$ or $=$ 45 year	s in males and $>$ or $=$ 55 years in females	3. Current Cigarette smoking or tobacco use		
2. Family history of premature ASCVD		4. High blood pressure		
5. Low HDL				

Newer treatment goals and statin initiation thresholds based on the risk categories proposed by LAI in 2020.

Risk Group	Treatment Goals		Consider Drug Therapy	
	LDL-C (mg/dl)	Non-HDL (mg/dl)	LDL-C (mg/dl)	Non-HDL (mg/dl)









SEX : Male



CLIENT CODE: CA00010147 - MEDIWHEEL CLIENT'S NAME AND ADDRESS !! THOADE I MITTED

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED F701A, LADO SARAI, NEW DELHI, SOUTH DELHI, DELHI, SOUTH DELHI 110030 DELHI INDIA 8800465156

PATIENT NAME : KETHA TARUN KUMAR

ACCESSION NO : 4053WC000958 AGE : 30 Years

DRAWN :

REFERRING DOCTOR : DR. MEDIWHEEL

Test Report Status **Final**

Extreme Risk Group	<50 (Optional goal	< 80 (Optional goal	>OR = 50	>OR = 80
Category A	< OR = 30)	< OR = 60)		
Extreme Risk Group	<or 30<="" =="" td=""><td><or 60<="" =="" td=""><td>> 30</td><td>>60</td></or></td></or>	<or 60<="" =="" td=""><td>> 30</td><td>>60</td></or>	> 30	>60
Category B				
Very High Risk	<50	<80	>OR= 50	>OR= 80
High Risk	<70	<100	>OR= 70	>OR=100
Moderate Risk	<100	<130	>OR=100	>OR=130
Low Risk	<100	<130	>OR=130*	>OR=160

RECEIVED : 11/03/2023 09:26

Results

*After an adequate non-pharmacological intervention for at least 3 months.

References: Management of Dyslipidaemia for the Prevention of Stroke: Clinical Practice Recommendations from the Lipid Association of India. Current Vascular Pharmacology, 2022, 20, 134-155.

LIVER FUNCTION TEST WITH GGT

BILIRUBIN, TOTAL	0.50	General Range : < 1.1	mg/dL
BILIRUBIN, DIRECT	0.21	General Range : < 0.3	mg/dL
BILIRUBIN, INDIRECT	0.29	0.00 - 0.60	mg/dL
TOTAL PROTEIN	7.8	Ambulatory : 6.4 - 8.3 Recumbant : 6 - 7.8	g/dL
ALBUMIN	5.0	20-60yrs : 3.5 - 5.2	g/dL
GLOBULIN	2.8	2.0 - 4.0	g/dL
ALBUMIN/GLOBULIN RATIO	1.8	1.0 - 2.0	RATIO
ASPARTATE AMINOTRANSFERASE (AST/SGOT)	22	Adults : < 40	U/L
ALANINE AMINOTRANSFERASE (ALT/SGPT)	29	Adults : < 45	U/L
ALKALINE PHOSPHATASE	112	Adult(<60yrs): 40 - 130	U/L
GAMMA GLUTAMYL TRANSFERASE (GGT)	27	Adult(male) : < 60	U/L
TOTAL PROTEIN, SERUM			
TOTAL PROTEIN	7.8	Ambulatory : 6.4 - 8.3 Recumbant : 6 - 7.8	g/dL
URIC ACID, SERUM			
URIC ACID	5.3	Adults : 3.4-7	mg/dL
ABO GROUP & RH TYPE, EDTA WHOLE BLOOD			
ABO GROUP	TYPE A		
RH TYPE	POSITIVE		
BLOOD COUNTS, EDTA WHOLE BLOOD			
HEMOGLOBIN	16.2	13.0 - 17.0	g/dL





DDRC SRL DIAGNOSTICS

PATIENT ID :

CLIENT PATIENT ID :

ABHA NO:

REPORTED :

KANNUR KERALA, INDIA Tel: 93334 93334 Email : customercare.ddrc@srl.in

11/03/2023 17:56

KETHM1008924053

Units







KETHM1008924053

CLIENT CODE : CA00010147 - MEDIWHEEL CLIENT'S NAME AND ADDRESS? THOADE HIMITED

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED F701A, LADO SARAI, NEW DELHI, SOUTH DELHI, DELHI, SOUTH DELHI 110030 DELHI INDIA 8800465156

PATIENT NAME : KETHA TARUN KUMAR

DDRC SRL DIAGNOSTICS

PATIENT ID :

CLIENT PATIENT ID :

Email : customercare.ddrc@srl.in

KANNUR

KERALA, INDIA

Tel: 93334 93334

ACCESSION NO: 4053WC000958 AGE: 30 Years SEX : Male ABHA NO: RECEIVED : 11/03/2023 09:26 REPORTED : 11/03/2023 17:56 DRAWN :

REFERRING DOCTOR : DR. MEDIWHEEL

Test Report Status <u>Final</u>	Results			Units
	F 40			
RED BLOOD CELL COUNT	5.49		4.5 - 5.5	mil/µL
WHITE BLOOD CELL COUNT	7.54		4.0 - 10.0	thou/µL
PLATELET COUNT	272		150 - 410	thou/µL
RBC AND PLATELET INDICES			40 50	0/
HEMATOCRIT	48.7		40 - 50	%
MEAN CORPUSCULAR VOL	88.6		83 - 101	fL
MEAN CORPUSCULAR HGB.	29.4		27.0 - 32.0	pg
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION	33.2		31.5 - 34.5	g/dL
RED CELL DISTRIBUTION WIDTH	13.1		11.6 - 14.0	%
MENTZER INDEX	16.1			
MEAN PLATELET VOLUME	9.2		6.8 - 10.9	fL
WBC DIFFERENTIAL COUNT				
SEGMENTED NEUTROPHILS	63		40 - 80	%
LYMPHOCYTES	30		20 - 40	%
MONOCYTES	2		2 - 10	%
EOSINOPHILS	4		1 - 6	%
BASOPHILS	1		0 - 2	%
ABSOLUTE NEUTROPHIL COUNT	4.75		2.0 - 7.0	thou/µL
ABSOLUTE LYMPHOCYTE COUNT	2.26		1 - 3	thou/µL
ABSOLUTE MONOCYTE COUNT	0.15	Low	0.20 - 1.00	thou/µL
ABSOLUTE EOSINOPHIL COUNT	0.30		0.02 - 0.50	thou/µL
NEUTROPHIL LYMPHOCYTE RATIO (NLR)	2.1			
ERYTHROCYTE SEDIMENTATION RATE (ESR),W BLOOD	HOLE			
SEDIMENTATION RATE (ESR) SUGAR URINE - POST PRANDIAL	21	High	0 - 14	mm at 1 hr
SUGAR URINE - POST PRANDIAL THYROID PANEL, SERUM	NOT DETECTED		NOT DETECTED	
Т3	147.80		80.00 - 200.00	ng/dL
T4	10.82		5.10 - 14.10	µg/dl
TSH 3RD GENERATION	3.010		21-50 yrs : 0.4 - 4.2	µIU/mL







Test Report Status	<u>Final</u>	Results	Units

Interpretation(s)

Triiodothyronine T3, Thyroxine T4, and Thyroid Stimulating Hormone TSH are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hyperthyroidism, TSH levels are low. Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3.Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism.Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Sr. No.	TSH	Total T4	FT4	Total T3	Possible Conditions
1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3)
		(a a.e.			Post Thyroidectomy (4) Post Radio-Iodine treatment
2	High	Normal	Normal	Normal	(1)Subclinical Hypothyroidism (2) Patient with insufficient thyroid
					hormone replacement therapy (3) In cases of Autoimmune/Hashimoto
					thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical
					inflammation, drugs like amphetamines, Iodine containing drug and
					dopamine antagonist e.g. domperidone and other physiological reasons.
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre
					(3)Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid
					hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4
					replacement therapy (7) First trimester of Pregnancy
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism
6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent
					treatment for Hyperthyroidism
8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies

REF: 1. TIETZ Fundamentals of Clinical chemistry 2.Guidlines of the American Thyroid association during pregnancy and Postpartum, 2011. **NOTE: It is advisable to detect Free T3,FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.**TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

PHYSICAL EXAMINATION, URINE

COLOR	
APPEARANCE	
CHEMICAL EXAMINATION, URINE	

PALE YELLOW CLEAR

6

4.7 - 7.5



PH









/HPF

CLIENT CODE : CA00010147 - MEDIWHEEL CLIENT'S NAME AND ADDRESS !! THOADE I MATTER

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED F701A, LADO SARAI, NEW DELHI, SOUTH DELHI, DELHI, SOUTH DELHI 110030 DELHI INDIA 8800465156

DDRC SRL DIAGNOSTICS

NOT DETECTED

NOT DETECTED

KANNUR KERALA, INDIA Tel: 93334 93334 Email : customercare.ddrc@srl.in

PATIENT NAME : KETHA TARUN	KUMAR	PATIENT ID : K	ETHM1008924053
ACCESSION NO : 4053WC000958	AGE : 30 Years SEX : Male	ABHA NO :	
DRAWN :	RECEIVED : 11/03/2023 09:26	REPORTED : 11/03/2023	17:56
REFERRING DOCTOR : DR. MEDIWH	EEL	CLIENT PATIENT ID :	
Test Report Status <u>Final</u>	Results		Units
SPECIFIC GRAVITY	1.015	1.003 - 1.035	
PROTEIN	NOT DETECTED	NOT DETECTED	
GLUCOSE	NOT DETECTED	NOT DETECTED	
KETONES	NOT DETECTED	NOT DETECTED	
BILIRUBIN	NOT DETECTED	NOT DETECTED	
UROBILINOGEN	NORMAL	NORMAL	
NITRITE	NOT DETECTED	NOT DETECTED	
MICROSCOPIC EXAMINATION, U	JRINE		
RED BLOOD CELLS	NOT DETECTED	NOT DETECTED	/HPF
WBC	1-2	0-5	/HPF

DETECTED

ABSENT ABSENT

(OCCASIONAL)

NOT DETECTED

CASTS CRYSTALS

EPITHELIAL CELLS

BACTERIA

Scan to View Details









CLIENT CODE: CA00010147 - MEDIWHEEL CLIENT'S NAME AND ADDRESS' THOADE HIMITED

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED F701A, LADO SARAI, NEW DELHI, SOUTH DELHI, DELHI, SOUTH DELHI 110030 DELHI INDIA 8800465156

DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334 Email : customercare.ddrc@srl.in

PATIENT NAME : KETHA TARU	JN KUMAR	PATIENT ID : KETHM1008924053
ACCESSION NO : 4053WC0009	58 AGE : 30 Years SEX : Male	ABHA NO :
DRAWN :	RECEIVED : 11/03/2023 09:26	REPORTED : 11/03/2023 17:56
REFERRING DOCTOR : DR. MEDIV	VHEEL	CLIENT PATIENT ID :
Test Report Status <u>Final</u>	Results	Units

Interpretation(s)

The following table describes the probable conditions, in which the analytes are present in urine

Presence of	Conditions
Proteins	Inflammation or immune illnesses
Pus (White Blood Cells)	Urinary tract infection, urinary tract or kidney stone, tumors or any kind of kidney impairment
Glucose	Diabetes or kidney disease
Ketones	Diabetic ketoacidosis (DKA), starvation or thirst
Urobilinogen	Liver disease such as hepatitis or cirrhosis
Blood	Renal or genital disorders/trauma
Bilirubin	Liver disease
Erythrocytes	Urological diseases (e.g. kidney and bladder cancer, urolithiasis), urinary tract infection and glomerular diseases
Leukocytes	Urinary tract infection, glomerulonephritis, interstitial nephritis either acute or chronic, polycystic kidney disease, urolithiasis, contamination by genital secretions
Epithelial cells	Urolithiasis, bladder carcinoma or hydronephrosis, ureteric stents or bladder catheters for prolonged periods of time
Granular Casts	Low intratubular pH, high urine osmolality and sodium concentration, interaction with Bence-Jones protein
Hyaline casts	Physical stress, fever, dehydration, acute congestive heart failure, renal diseases
Calcium oxalate	Metabolic stone disease, primary or secondary hyperoxaluria, intravenous infusion of large doses of vitamin C, the use of vasodilator naftidrofuryl oxalate or the gastrointestinal lipase inhibitor orlistat, ingestion of ethylene glycol or of star fruit (Averrhoa carambola) or its juice
Uric acid	arthritis
Bacteria	Urinary infectionwhen present in significant numbers & with pus cells.
Trichomonas vaginalis	Vaginitis, cervicitis or salpingitis

SUGAR URINE - FASTING

SUGAR URINE - FASTING PHYSICAL EXAMINATION, STOOL

COLOUR CONSISTENCY MUCUS VISIBLE BLOOD NOT DETECTED

NOT DETECTED

YELLOW SEMI LIQUID NOT DETECTED ABSENT

NOT DETECTED ABSENT











CLIENT CODE : CA00010147 - MEDIWHEEL CLIENT'S NAME AND ADDRESS ! THOADE I MATTER

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED F701A, LADO SARAI, NEW DELHI, SOUTH DELHI, DELHI, SOUTH DELHI 110030 DELHI INDIA 8800465156

VEGETABLE CELLS

DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334 Email : customercare.ddrc@srl.in

PATIENT NAME : KETHA TARUN	KUMAR	PATIENT ID : K	ETHM1008924053
ACCESSION NO : 4053WC000958	AGE : 30 Years SEX : Male	ABHA NO:	
DRAWN :	RECEIVED : 11/03/2023 09:26	REPORTED : 11/03/2023	17:56
REFERRING DOCTOR : DR. MEDIWH	EEL	CLIENT PATIENT ID :	
Test Report Status <u>Final</u>	Results		Units
MICROSCOPIC EXAMINATION,S	TOOL		
PUS CELLS	0-1		/hpf
RED BLOOD CELLS	NOT DETECTED	NOT DETECTED	/HPF
CYSTS	NOT DETECTED	NOT DETECTED	
OVA	NOT DETECTED		

ABSENT











CLIENT CODE: CA00010147 - MEDIWHEEL CLIENT'S NAME AND ADDRESS ! THOADE HIMITED

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED F701A, LADO SARAI, NEW DELHI, SOUTH DELHI, DELHI, SOUTH DELHI 110030 DELHI INDIA 8800465156

DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334 Email : customercare.ddrc@srl.in

PATIENT NAME : KETH	IA TARUN KUMAR	PATIENT ID : KETHM1008924053
ACCESSION NO : 4053V	VC000958 AGE : 30 Years SEX : M	ale ABHA NO :
DRAWN :	RECEIVED : 11/03/2023 09	0:26 REPORTED : 11/03/2023 17:56
REFERRING DOCTOR : DF	R. MEDIWHEEL	CLIENT PATIENT ID :
Test Report Status	Final Results	Units

Interpretation(s)

Stool routine analysis is only a screening test for disorders of gastrointentestinal tract like infection, malabsorption, etc. The following table describes the probable conditions, in which the analytes are present in stool.

PRESENCE OF	CONDITION
Pus cells	Pus in the stool is an indication of infection
Red Blood cells	Parasitic or bacterial infection or an inflammatory bowel condition such as ulcerative colitis
Parasites	Infection of the digestive system. Stool examination for ova and parasite detects presence of parasitic infestation of gastrointestinal tract. Various forms of parasite that can be detected include cyst, trophozoite and larvae. One negative result does not rule out the possibility of parasitic infestation. Intermittent shedding of parasites warrants examinations of multiple specimens tested on consecutive days. Stool specimens for parasitic examination should be collected before initiation of antidiarrheal therapy or antiparasitic therapy. This test does not detect presence of opportunistic parasites like Cyclospora, Cryptosporidia and Isospora species. Examination of Ova and Parasite has been carried out by direct and concentration techniques.
Mucus	Mucus is a protective layer that lubricates, protects& reduces damage due to bacteria or viruses.
Charcot-Leyden crystal	Parasitic diseases.
Ova & cyst	Ova & cyst indicate parasitic infestation of intestine.
Frank blood	Bleeding in the rectum or colon.
Occult blood	Occult blood indicates upper GI bleeding.
Macrophages	Macrophages in stool are an indication of infection as they are protective cells.
Epithelial cells	Epithelial cells that normally line the body surface and internal organs show up in stool when there is inflammation or infection.
Fat	Increased fat in stool maybe seen in conditions like diarrhoea or malabsorption.
рН	Normal stool pH is slightly acidic to neutral. Breast-fed babies generally have an acidic stool.

ADDITIONAL STOOL TESTS :

- Stool Culture:- This test is done to find cause of GI infection, make decision about best treatment for GI infection & to find out if 1. treatment for GI infection worked.
- 2. Fecal Calprotectin: It is a marker of intestinal inflammation. This test is done to differentiate Inflammatory Bowel Disease (IBD) from Irritable Bowel Syndrome (IBS).
- 3. Fecal Occult Blood Test(FOBT): This test is done to screen for colon cancer & to evaluate possible cause of unexplained anaemia.
- 4. Clostridium Difficile Toxin Assay: This test is strongly recommended in healthcare associated bloody or waterydiarrhoea, due to overuse of broad spectrum antibiotics which alter the normal GI flora.
- 5. Biofire (Film Array) GI PANEL: In patients of Diarrhoea, Dysentry, Rice watery Stool, FDA approved, Biofire Film Array Test, (Real Time Multiplex PCR) is strongly recommended as it identifies organisms, bacteria, fungi, virus, parasite and other opportunistic pathogens, Vibrio cholera infections only in 3 hours. Sensitivity 96% & Specificity 99%.
- 6. Rota Virus Immunoassay: This test is recommended in severe gastroenteritis in infants & children associated with watery











CLIENT CODE: CA00010147 - MEDIWHEEL CLIENT'S NAME AND ADDRESS !! TUCADO

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED F701A, LADO SARAI, NEW DELHI, SOUTH DELHI, DELHI, SOUTH DELHI 110030 DELHI INDIA 8800465156

DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334 Email : customercare.ddrc@srl.in

PATIENT NAME : KETHA TARUN	KUMAR	PATIENT ID : KETHM1008924053
ACCESSION NO : 4053WC000958	AGE : 30 Years SEX : Male	ABHA NO :
DRAWN :	RECEIVED : 11/03/2023 09:26	REPORTED : 11/03/2023 17:56
REFERRING DOCTOR : DR. MEDIWH	EEL	CLIENT PATIENT ID :
Test Report Status Final	Results	Units

diarrhoea, vomitting& abdominal cramps. Adults are also affected. It is highly contagious in nature.

Interpretation(s)

CREATININE, SERUM-Higher than normal level may be due to:

 Blockage in the urinary tract Kidney problems, such as kidney damage or failure, infection, or reduced blood flow

Loss of body fluid (dehydration)
Muscle problems, such as breakdown of muscle fibers

• Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:

Myasthenia Gravis

Muscular dystrophy

GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.Additional test HbA1c GLUCOSE FASTING, FLUORIDE PLASMA- TEST DESCRIPTION

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in the urine.

Increased in

Diabetes mellitus, Cushing' s syndrome (10 - 15%), chronic pancreatitis (30%). Drugs:corticosteroids, phenytoin, estrogen, thiazides.

Decreased in

Pancreatic islet cell disease with increased insulin, insulinoma, adrenocortical insufficiency, hypopituitarism, diffuse liver disease, malignancy (adrenocortical,

stomach,fibrosarcoma), infant of a diabetic mother, enzyme deficiency diseases(e.g., galactosemia),Drugs- insulin,

ethanol, propranolol; sulfonylureas,tolbutamide, and other oral hypoglycemic agents. NOTE: While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values), there is wide fluctuation within individuals. Thus, glycosylated hemoglobin (HbA1c) levels are favored to monitor glycemic control.

High fasting glucose level in comparison to post practial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-Used For:

Evaluating the long-term control of blood glucose concentrations in diabetic patients.

2.Diagnosing diabetes. 3.Identifying patients at increased risk for diabetes (prediabetes).

The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for well-controlled type 2 diabetic patients) to determine whether a patients metabolic control has remained continuously within the target range.

1.eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.

eAG gives an evaluation of blood glucose levels for the last couple of months.
 eAG is calculated as eAG (mg/dl) = 28.7 * HbA1c - 46.7

HbA1c Estimation can get affected due to :

I.Shortened Erythrocyte survival : Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss, hemolytic anemia) will falsely lower HbA1c test results. Fructosamine is recommended in these patients which indicates diabetes control over 15 days II. Vitamin C & E are reported to falsely lower test results. (possibly by inhibiting glycation of hemoglobin.

III. Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates addiction are reported to interfere with some assay methods, falsely increasing results.

VI.Interference of hemoglobinopathies in HbA1c estimation is seen in a.Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c. b.Heterozygous state detected (D10 is corrected for HbS & HbC trait.)

c.HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c.Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy

TOTAL PROTEIN, SERUM-Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum .. Protein in the plasma is made up of albumin and globulin









CLIENT CODE: CA00010147 - MEDIWHEEL CLIENT'S NAME AND ADDRESS ! THOADE HIMITED

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED F701A, LADO SARAI, NEW DELHI, SOUTH DELHI, DELHI, SOUTH DELHI 110030 DELHI INDIA 8800465156

DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334 Email : customercare.ddrc@srl.in

PATIENT NAME : K	ETHA TARUN KU	JMAR		PA	TIENT ID :	KETHM1008924053
ACCESSION NO : 40	53WC000958	AGE: 30 Years	SEX : Male	ABHA NO :		
DRAWN :		RECEIVED : 11/03	/2023 09:26	REPORTED :	11/03/202	3 17:56
REFERRING DOCTOR :	DR. MEDIWHEEL	-		CLIEN	PATIENT ID	:
		_				

-	Test Report Status	<u>Final</u>	Results	Units

syndrome, Protein-losing enteropathy etc.

URIC ACID, SERUM-Causes of Increased levels:-Dietary(High Protein Intake, Prolonged Fasting, Rapid weight loss), Gout, Lesch nyhan syndrome, Type 2 DM, Metabolic syndrome

Causes of decreased levels-Low Zinc intake,OCP,Multiple Sclerosis ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-

Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same.

The test is performed by both forward as well as reverse grouping methods.

BLOOD COUNTS, EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology. RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13)

from Beta thalassaemia trait

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients ; A.-P. Yang, et al.; International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope.

ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD-**TEST DESCRIPTION** :-Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic; it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change.

TEST INTERPRETATION

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy, Estrogen medication, Aging.

Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias, Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis). In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum.

Decreased in: Polycythermia vera, Sickle cell anemia

LIMITATIONS

False elevated ESR : Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia

False Decreased : Poikilocytosis, (SickleCells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs (Quinine, salicvlates)

REFERENCE

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition; 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin; 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th edition. SUGAR URINE - POST PRANDIAL-METHOD: DIPSTICK/BENEDICT'S TEST SUGAR URINE - FASTING-METHOD: DIPSTICK/BENEDICT'S TEST







REPORT COMPLETED CHEST X-RAY WITH REPORT

REPORT COMPLETED

> **End Of Report** Please visit www.srlworld.com for related Test Information for this accession

JINSHA KRISHNAN LAB TECHNOLOGIST



DR.INDUSARATH S CONSULTANT PATHOLOGIST

iticate

NIMISHA K LAB TECHNOLOGIST

Trees

SREENA A LAB TECHNOLOGIST







OPTHALMOLOGY REPORT

TO WHOM-SO-EVER IT MAY CONCERN

This is to certify that I have examined Mr. KETHA TARUN KUMAR, 30 years Male on 11.03.2023 and his visual standards are as follows:

	OD	OS
UNCORRECTED DISTANCE VISUAL ACUITY	5/60	6/60
UNCORRECTED NEAR VISUAL ACUITY	N6	N6
BEST CORRECTED VISUAL ACUITY	6/6,N6	6/6,N6
COLOUR VISION	NORMAL	NORMAL

NOTE : HISTORY SPECS SINCE 10 YEARS ; LAST CHANGED 1 YEARS BACK. NO RELEVANT MEDICAL HISTORY.

SHEHZIYA V P OPTOMETRIST



DATE: 11.03.2023



Name	Mr. KETHA TARUN KUMAR	Age/Sex	30Yrs/Male
Ref: By:	MEDI WHEEL	Date	11.03.2023

Thanks for referral

CHEST X-RAY - PA VIEW

Trachea is central. Carina and principal bronchi are normal.

Cardio-thoracic ratio is within normal limits.

Both lungs show normal Broncho-vascular markings. No definite focal opacities noted.

No volume loss in either hemithorax.

No definite mediastinal widening or other abnormalities noted.

CP angles, diaphragm, bony cage and soft tissue shadows - not remarkable.

IMPRESSION:

Normal X-ray chest

DR. P. NIYAZI NASIR, MBBS, DMRD

(Because of technical and technological limitation complete diagnosis cannot be assured on imaging sonography. Clinical correlation, consultation if required repeat imaging required in the event of controversies. This document is not for legal purposes).



Dr. P. NIYAZI NASIR. MBBS, DMRD REG. No. 41419 CONSULTANT RADIOLOGIST DDRC SRL DIAGNOSTIC (P) LTD. KANNUR





Name	Mr. KETHA TARUN KUMAR	Age/Sex	30Yrs/Male
Ref: By:	MEDI WHEEL	Date	11.03.2023

Thanks for referral

CHEST X-RAY - PA VIEW

Trachea is central. Carina and principal bronchi are normal.

Cardio-thoracic ratio is within normal limits.

Both lungs show normal Broncho-vascular markings. No definite focal opacities noted.

No volume loss in either hemithorax.

No definite mediastinal widening or other abnormalities noted.

CP angles, diaphragm, bony cage and soft tissue shadows - not remarkable.

IMPRESSION:

Normal X-ray chest

DR. P. NIYAZI NASIR, MBBS, DMRD

(Because of technical and technological limitation complete diagnosis cannot be assured on imaging sonography. Clinical correlation, consultation if required repeat imaging required in the event of controversies. This document is not for legal purposes).



Dr. P. NIYAZI NASIR. MBBS, DMRD REG. No. 41419 CONSULTANT RADIOLOGIST DDRC SRL DIAGNOSTIC (P) LTD. KANNUR

