

CLIENT'S NAME AND ADDRESS:
ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )
F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL Ltd S.K. Tower,Hari Niwas, LBS Marg THANE, 400602

MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: HARESH G GULRAJANI

PATIENT ID: HAREM180571181

ACCESSION NO: 0181VI000319 AGE: 51 Years SEX: Male

DRAWN: RECEIVED: 10/09/2022 09:57

REPORTED: 13/09/2022 15:18

REFERRING DOCTOR: SELF

CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units **Final** 

# MEDI WHEEL FULL BODY HEALTH CHECK UP ABOVE 40 MALE

PHYSICAL EXAMINATION, URINE			
COLOR	PALE YELLOW		
METHOD: VISUAL INSPECTION			
APPEARANCE	CLEAR		
METHOD: VISUAL INSPECTION			
SPECIFIC GRAVITY	1.005	1.003 - 1.035	
METHOD: IONIC CONCENTRATION METHOD			
BLOOD COUNTS,EDTA WHOLE BLOOD			
HEMOGLOBIN	16.1	13.0 - 17.0	g/dL
METHOD: SLS-HEMOGLOBIN DETECTION METHOD			
RED BLOOD CELL COUNT	5.49	4.5 - 5.5	mil/µL
METHOD: HYDRODYNAMIC FOCUSING BY DC DETECTION			
WHILE BLOOD CELL COUNT	5.75	4.0 - 10.0	thou/µL
METHOD: FLUORESCENCE FLOW CYTOMETRY			
PLATELET COUNT	232	150 - 410	thou/µL
METHOD: HYDRODYNAMIC FOCUSING BY DC DETECTION			
RBC AND PLATELET INDICES			
HEMATOCRIT	48.3	40.0 - 50.0	%
METHOD : CUMULATIVE PULSE HEIGHT DETECTION METHOD			
MEAN CORPUSCULAR VOL	88.0	83.0 - 101.0	†L
METHOD: CALCULATED FROM RBC & HCT			
MEAN CORPUSCULAR HGB.	29.3	27.0 - 32.0	pg
METHOD: CALCULATED FROM THE RBC & HGB			
MEAN CORPUSCULAR HEMOGLOBIN	33.3	31.5 - 34.5	g/dL
CONCENTRATION  METHOD: CALCULATED FROM THE HGB & HCT			
MENTZER INDEX	16.0		
RED CELL DISTRIBUTION WIDTH	11.7	11.6 - 14.0	%
METHOD : CALCULATED FROM RBC SIZE DISTRIBUTION CURVE			
MEAN PLATELET VOLUME	9.4	6.8 - 10.9	fL
METHOD : CALCULATED FROM PLATELET COUNT & PLATELET HEMA	- ' '		
CHEMICAL EXAMINATION, URINE			
P⊢	7.0	4.7 - 7.5	
METHOD: DOUBLE INDICATOR PRINCIPLE		. <del>-</del>	
PROTEIN	NOT DETECTED	NOT DETECTED	



METHOD: TETRA BROMOPHENOL BLUE/SULFOSALICYLIC ACID

Page 1 Ot 12 Scan to View Report



CLIENT'S NAME AND ADDRESS:
ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI

NEW DELHI 110030 DELHI INDIA 8800465156

SRL Ltd

S.K. Tower,Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: HARESH G GULRAJANI

PATIENT ID: HAREM180571181

ACCESSION NO: 0181VI000319 AGE: 51 Years SEX: Male

DRAWN: RECEIVED: 10/09/2022 09:57 REPORTED: 13/09/2022 15:18

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results		Biological Reference Interva	al Units
GLUCOSE	NOT DETECTED		NOT DETECTED	
METHOD : GLUCOSE OXIDASE PEROXIDASE	NOT DETECTED		NOT DETECTED	
KETONES	NOT DETECTED		NOT DETECTED	
METHOD: NITROPRUSSIDE REACTION	NOT DETECTED		NOT DETECTED	
BLOOD	NOT DETECTED		NOT DETECTED	
METHOD : PEROXIDASE	NOT BETEGIES		NOT BETEGTED	
UROBILINOGEN	NORMAL		NORMAL	
METHOD: MODIFIED EHRLICH REACTION	T TOTAL IL			
NITRITE	NOT DETECTED		NOT DETECTED	
METHOD: 1,2,3,4-TETRAHYDROBENZO(H)QUINOLIN-3-OL				
LEUKOCYTE ESTERASE	NOT DETECTED		NOT DETECTED	
WBC DIFFERENTIAL COUNT - NLR				
SEGMENTED NEUTROPHILS	45		40 - 80	%
METHOD : FLOW CYTOMETRY WITH LIGHT SCATTERING				
ABSOLUTE NEUTROPHIL COUNT	2.59		2.0 - 7.0	thou/µL
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING				
LYMPHOCYTES	47	High	20 - 40	%
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING				
ABSOLUTE LYMPHOCYTE COUNT	2.69		1.0 - 3.0	thou/µL
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING				
NEUTROPHIL LYMPHOCYTE RATIC (NLR)	1.0			
EOSINOPHILS	3		1 - 6	%
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING				
ABSOLUTE EOSINOPHIL COUNT	0.19		0.02 - 0.50	thou/µL
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING				
MONOCYTES	5		2 - 10	%
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING				
ABSOLUTE MONOCYTE COUNT	0.31		0.2 - 1.0	thou/µL
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING				
DIFFERENTIAL COUNT PERFORMED ON:	EDTA SMEAR			
MICROSCOPIC EXAMINATION, URINE				
PUS CELL (WBC'S)	1-2		0-5	/HPF
METHOD: MICROSCOPIC EXAMINATION				
EPITHELIAL CELLS	1-2		0-5	/HPF
METHOD: MICROSCOPIC EXAMINATION				
ERYTHROCYTES (RBC'S)	NOT DETECTED		NOT DETECTED	/HPF
METHOD: MICROSCOPIC EXAMINATION				
CASTS	NOT DETECTED			



Page 2 Ot 12 Scan to View Report



CLIENT'S NAME AND ADDRESS:
ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI

NEW DELHI 110030 DELHI INDIA 8800465156

SRL Ltd

S.K. Tower,Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: HARESH G GULRAJANI

PATIENT ID: HAREM180571181

ACCESSION NO: 0181VI000319 AGE: 51 Years SEX: Male

DRAWN: RECEIVED: 10/09/2022 09:57 REPORTED: 13/09/2022 15:18

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Reference Interv	al Units
METHOD I MICROSCOPIC EVAMINATION			
METHOD: MICROSCOPIC EXAMINATION  CRYSTALS	NOT DETECTED		
METHOD: MICROSCOPIC EXAMINATION	NOT DETECTED		
BACTERIA	NOT DETECTED	NOT DETECTED	
METHOD: MICROSCOPIC EXAMINATION	NOI DETECTED	NOT BETECTED	
YEAST	NOT DETECTED	NOT DETECTED	
MORPHOLOGY			
RBC	NORMOCYTIC NORMOCH	ROMIC	
WBC	NORMAL MORPHOLOGY	(3)/10	
METHOD: MICROSCOPIC EXAMINATION	NORMAL MONTHOLOGY		
PLATELETS	ADEQUATE		
ERYTHRO SEDIMENTATION RATE, BLOOD			
SEDIMENTATION RAIL (ESR)	05	0 - 14	mm at 1 hr
METHOD: WESTERGREN METHOD	00	0 14	mm de i m
GLYCOSYLATED HEMOGLOBIN, EDTA WHOLE B	BLOOD		
GLYCOSYLATED HEMOGLOBIN (HBA1C)	5.3	Non-diabetic: < 5.7 Pre-diabetics: 5.7 - 6.4 Diabetics: > or = 6.5 ADA Target: 7.0 Action suggested: > 8.0	%
METHOD: HPLC	105.4	. 446.0	
MEAN PLASMA GLUCOSE  METHOD: CALCULATED PARAMETER	105.4	< 116.0	mg/dL
GLUCOSE, FASTING, PLASMA			
	99	Normal 75 - 99	ma/dl
GLUCOSE, FASTING, PLASMA	99	Pre-diabetics: 100 - 125 Diabetic: > or = 126	mg/dL
METHOD: ENZYMATIC REFERENCE METHOD WITH HEXOKINASE			
GLUCOSE, POST-PRANDIAL, PLASMA			
GLUCOSE, POST-PRANDIAL, PLASMA	92	70 - 139	mg/dL
METHOD: ENZYMATIC REFERENCE METHOD WITH HEXOKINASE			
CORONARY RISK PROFILE, SERUM			
CHOLESTEROL	183	Desirable cholesterol level < 200 Borderline high cholesterol 200 - 239 High cholesterol > / = 240	mg/dL
METHOD - ENTRAFTED OOL ODINETDIO 4004V			

METHOD: ENZYMATIC COLORIMETRIC ASSAY



Page 3 Ot 12 Scan to View Report



CLIENT'S NAME AND ADDRESS:
ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI

NEW DELHI 110030 DELHI INDIA 8800465156

SRL Ltd

S.K. Tower,Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: HARESH G GULRAJANI

PATIENT ID: HAREM180571181

ACCESSION NO: 0181VI000319 AGE: 51 Years SEX: Male

DRAWN: RECEIVED: 10/09/2022 09:57 REPORTED: 13/09/2022 15:18

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results		Biological Reference Interva	al Units
TRIGLYCERIDES  METHOD: ENZYMATIC COLORIMETRIC ASSAY	88		Normal: < 150 Borderline high: 150 - 199 High: 200 - 499 Very High: >/= 500	mg/dL
HDL CHOLESTEROL	41.1		Low HDL Cholesterol <40	mg/dL
TIBLE OF IGLEST ENGE	71.1		Low Fibe difforesteror 140	mg/ac
METHOD : ENZYMATIC, COLORIMETRIC			High HDL Cholesterol >/= 60	
CHOLESTEROL LDL	124	High	Adult levels: Optimal < 100 Near optimal/above optimal: 11 129 Borderline high: 130-159 High: 160-189 Very high: = 190	mg/dL 00-
METHOD : ENZYMATIC COLORIMETRIC ASSAY			very riigir . — 190	
NON HDL CHOLESTEROL	142	High	Desirable : < 130 Above Desirable : 130 -159 Borderline High : 160 - 189 High : 190 - 219 Very high : > / = 220	mg/dL
CHOL/HDL RATIO  LDL/HDL RATIO	<b>4.5</b> 3.0	High	Low Risk: 3.3 - 4.4 Average Risk: 4.5 - 7.0 Moderate Risk: 7.1 - 11.0 High Risk: > 11.0 0.5 - 3.0 Desirable/Low Risk	
BUTTELLATIO	3.0		3.1 - 6.0 Borderline/Moderate Risk >6.0 High Risk	
VERY LOW DENSITY LIPOPROTEIN	17.6		< OR = 30.0	mg/dL
LIVER FUNCTION PROFILE, SERUM				
BILIRUBIN, TOTAL METHOD: COLORIMETRIC DIAZO	0.60		Upto 1.2	mg/dL
BILIRUBIN, DIRECT	0.26		< 0.30	mg/dL
BILIRUBIN, INDIRECT	0.34		0.1 - 1.0	mg/dL
TOTAL PROTEIN	7.4		6.0 - 8.0	g/dL
METHOD: COLORIMETRIC				
ALBUMIN	5.0	High	3.97 - 4.94	g/dL
METHOD: COLORIMETRIC				
GLOBULIN	2.4		2.0 - 3.5	g/dL
ALBUMIN/GLOBULIN RATIO	2.1		1.0 - 2.1	RATIO
ASPARTATE AMINOTRANSFERASE (AST/SGOT)  METHOD: UV ABSORBANCE	28		< OR = 50	U/L







CLIENT'S NAME AND ADDRESS:
ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI

NEW DELHI 110030 DELHI INDIA 8800465156

SRL Ltd

S.K. Tower,Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: HARESH G GULRAJANI

PATIENT ID: HAREM180571181

ACCESSION NO: 0181VI000319 AGE: 51 Years SEX: Male

DRAWN: RECEIVED: 10/09/2022 09:57

REPORTED: 13/09/2022 15:18

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Reference In	terval Units
ALANINIE AMINIOTRANICEERACE (ALTICORT)	26	< OD _ F0	1.171
ALANINE AMINOTRANSFERASE (ALT/SGPT) METHOD: UV ABSORBANCE	26	< OR = 50	U/L
ALKALINE PHOSPHATASE	64	40 - 129	U/L
METHOD : COLORIMETRIC			J, L
GAMMA GLUTAMYL TRANSFERASE (GGT)	8	0 - 60	U/L
METHOD : ENZYMATIC, COLORIMETRIC			
LACTATE DEHYDROGENASE	180	125 - 220	U/L
METHOD: UV ABSORBANCE			
SERUM BLOOD UREA NITROGEN			
BLOOD UREA NITROGEN	13	6 - 20	mg/dL
METHOD: ENZYMATIC ASSAY			
CREATININE, SERUM			
CREATININE	1.01	0.7 - 1.2	mg/dL
METHOD: COLORIMETRIC			
BUN/CREAT RATIO			
BUN/CREAT RATIO	12.87	8.0 - 15.0	
URIC ACID, SERUM			
URIC ACID	6.4	3.4 - 7.0	mg/dL
METHOD: ENZYMATIC COLORIMETRIC ASSAY			
TOTAL PROTEIN, SERUM			
TOTAL PROTEIN	7.4	6.0 - 8.0	g/dL
METHOD: COLORIMETRIC			
ALBUMIN, SERUM			
ALBUMIN	5.0	<b>High</b> 3,97 - 4,94	g/dL
METHOD: COLORIMETRIC			
GLOBULIN			
GLOBULIN	2.4	2.0 - 3.5	g/dL
ELECTROLYTES (NA/K/CL), SERUM			
SODIUM	143	136 - 145	mmol/L
POTASSIUM	4.93	3.5 - 5.1	mmol/L
CHLORIDE	104	98 - 107	mmol/L
THYROID PANEL, SERUM			
T3	95.5	80 - 200	ng/dL
METHOD: ELECTROCHEMILUMINESCENCE			3, ==
T4	6.79	5.1 - 14.1	µg/dL
METHOD: ELECTROCHEMILUMINESCENCE			
TSH 3RD GENERATION	1.620	0.27 - 4.2	μIU/mL



Page 5 Ot 12 Scan to View Report



ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT ID:

HAREM180571181

Email: customercare.thane@srl.in

PATIENT NAME: HARESH G GULRAJANI

ACCESSION NO: 0181VI000319 AGE: 51 Years SEX: Male

DRAWN: RECEIVED: 10/09/2022 09:57 REPORTED: 13/09/2022 15:18

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units <u>Final</u>

METHOD: ELECTROCHEMILUMINESCENCE

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

ABO GROUP TYPE O

METHOD: GEL COLUMN AGGLUTINATION METHOD.

RH TYPE **POSITIVE** 

METHOD: GEL COLUMN AGGLUTINATION METHOD.

XRAY-CHEST

**IMPRESSION** NO ABNORMALITY DETECTED

TMT OR ECHO

TMT OR ECHO 2D ECHO :-

Structurally normal valves. Mild Concentric LVH.

No regional wall motion abnormality.

Good Left Ventricular systolic function. LVEF 60 %

Normal LV Diastolic function. No e/o pulmonary hypertension.

**ECG** 

ECG WITHIN NORMAL LIMITS

MEDICAL HISTORY

RELEVANT PRESENT HISTORY NOT SIGNIFICANT

RELEVANT PAST HISTORY COVID 2 YEARS BACK, ISOLATED IN QUARANTINE.

RELEVANT PERSONAL HISTORY MARRIED / 2 CHILD / MIXED DIET / NO ALLERGIES / NO SMOKING /

OCC ALCOHOL.

RELEVANT FAMILY HISTORY HIGH BLOOD PRESSURE & DIABETS: MOTHER.

ANTHROPOMETRIC DATA & BMI

HEIGHT IN METERS 1.67 mts WEIGHT IN KGS. 74 Kgs BMI BMI & Weight Status as follows: kg/sqmts 27

Below 18.5: Underweight 18.5 - 24.9: Normal 25.0 - 29.9: Overweight

30.0 and Above: Obese

GENERAL EXAMINATION

MENTAL / EMOTIONAL STATE NORMAL PHYSICAL ATTITUDE NORMAL GENERAL APPEARANCE / NUTRITIONAL STATUS **OVERWEIGHT** BUILT / SKELETAL FRAMEWORK **AVERAGE** FACIAL APPEARANCE NORMAL SKIN NORMAL UPPER LIMB NORMAL



Page 6 Ot 12 回路深級區 Scan to View Report



ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI DELHI INDIA 8800465156

NEW DELHI 110030

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

78/MIN.REGULAR, ALL PERIPHERAL PULSES WELL FELT, NO CAROTID

Email: customercare.thane@srl.in

S.K. Tower, Hari Niwas, LBS Marg

SRL Ltd

THANE, 400602 MAHARASHTRA, INDIA

PATIENT NAME: HARESH G GULRAJANI

PATIENT ID: HAREM180571181

ACCESSION NO: 0181VI000319 AGE: 51 Years SEX: Male

DRAWN: RECEIVED: 10/09/2022 09:57 REPORTED: 13/09/2022 15:18

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units **Final** LOWER LIMB NORMAL NECK NORMAL NECK LYMPHATICS / SALIVARY GLANDS NOT ENLARGED OR TENDER THYROID GLAND NOT ENLARGED CAROTID PULSATION NORMAL **TEMPERATURE** NORMAL

**BRUIT** RESPIRATORY RATE NORMAL

CARDIOVASCULAR SYSTEM

**PULSE** 

130/70 MM HG mm/Hg

(SUPINE)

PERICARDIUM NORMAL NORMAL APEX BEAT HEART SOUNDS NORMAL **MURMURS ABSENT** 

RESPIRATORY SYSTEM

SIZE AND SHAPE OF CHEST NORMAL MOVEMENTS OF CHEST SYMMETRICAL BREATH SOUNDS INTENSITY NORMAL

BREATH SOUNDS QUALITY VESICULAR (NORMAL)

ABSENT ADDED SOUNDS

PER ABDOMEN

**APPEARANCE** NORMAL VENOUS PROMINENCE ABSENT NOT PALPABLE **LIVER SPLEEN** NOT PALPABLE

**HERNIA ABSENT** 

CENTRAL NERVOUS SYSTEM

HIGHER FUNCTIONS NORMAL CRANIAL NERVES NORMAL CEREBELLAR FUNCTIONS NORMAL SENSORY SYSTEM NORMAL MOTOR SYSTEM NORMAL **REFLEXES** NORMAL







ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: HARESH G GULRAJANI

PATIENT ID: HAREM180571181

ACCESSION NO: 0181VI000319 AGE: 51 Years SEX: Male

DRAWN: RECEIVED: 10/09/2022 09:57 REPORTED: 13/09/2022 15:18

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Results Biological Reference Interval Test Report Status Units <u>Final</u>

MUSCULOSKELETAL SYSTEM

SPINE NORMAL JOINTS NORMAL

BASIC EYE EXAMINATION

CONJUNCTIVA NORMAL **EYELIDS** NORMAL EYE MOVEMENTS NORMAL CORNEA NORMAL

DISTANT VISION RIGHT EYE WITHOUT GLASSES REDUCED VISUAL ACUITY 6/18

DISTANT VISION LEFT EYE WITHOUT GLASSES WITHIN NORMAL LIMIT

NEAR VISION RIGHT EYE WITHOUT GLASSES REDUCED VISUAL ACUITY N/36 NEAR VISION LEFT EYE WITHOUT GLASSES REDUCED VISUAL ACUITY N/18

NEAR VISION RIGHT EYE WITH GLASSES WITHIN NORMAL LIMIT NEAR VISION LEFT EYE WITH GLASSES WITHIN NORMAL LIMIT

COLOUR VISION NORMAL

SUMMARY

NOT SIGNIFICANT RELEVANT HISTORY RELEVANT GP EXAMINATION FINDINGS OVERWEIGHT: BMI 27

1) LOW FAT, LOW CALORIE, LOW CARBOHYDRATE, HIGH FIBRE DIET, REMARKS / RECOMMENDATIONS

2) REGULAR EXERCISE.REGULAR WALK FOR 30-40 MIN DAILY.

3) REPEAT LIPID PROFILE AFTER 3 MONTHS OF DIET AND EXERCISE.

Interpretation(s)

BLOOD COUNTS, EDTA WHOLE BLOOD-

The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

WBC DIFFERENTIAL COUNT - NLRThe optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years

old and NLR - 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients; A.-P. Yang, et al.; International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope.
MICROSCOPIC EXAMINATION, URINERoutine urine analysis assists in screening and diagnosis of various metabolic, urological, kidney and liver disorders

Protein: Elevated proteins can be an early sign of kidney disease. Urinary protein excretion can also be temporarily elevated by strenuous exercise, orthostatic proteinuria, dehydration, urinary tract infections and acute illness with fever

Glucose: Uncontrolled diabetes mellitus can lead to presence of glucose in urine. Other causes include pregnancy, hormonal disturbances, liver disease and certain medications.

Ketones: Uncontrolled diabetes mellitus car lead to presence of ketones in urine. Ketones can also be seen in starvation, frequent vomiting, pregnancy and strenuous exercise.

Blood: Occult blood can occur in urine as intact erythrocytes or haemoglobin, which can occur in various urological, nephrological and bleeding disorders. Leukocytes: An increase in leukocytes is an indication of inflammation in urinary tract or kidneys. Most common cause is bacterial urinary tract infection.







ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI **NEW DELHI 110030** DELHI INDIA 8800465156

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: HARESH G GULRAJANI

PATIENT ID:

HAREM180571181

ACCESSION NO: 0181VI000319 AGE: 51 Years

SEX: Male

REPORTED:

13/09/2022 15:18

DRAWN:

REFERRING DOCTOR: SELF

RECEIVED: 10/09/2022 09:57

CLIENT PATIENT ID:

Test Report Status

<u>Final</u>

Results

Biological Reference Interval Units

Nitrite: Many bacteria give positive results when their number is high. Nitrite concentration during infection increases with length of time the urine specimen is retained in bladder prior to collection.

pH: The kidneys play an important role in maintaining acid base balance of the body. Conditions of the body producing acidosis/ alkalosis or ingestion of certain type of food

can affect the pH of urine.

Specific gravity: Specific gravity gives an indication of how concentrated the urine is. Increased specific gravity is seen in conditions like dehydration, glycosuria and proteinuria while decreased specific gravity is seen in excessive fluid intake, renal failure and diabetes insipidus.

Bilirubin: In certain liver diseases such as biliary obstruction or hepatitis, bilirubin gets excreted in urine.

Urobilinogen: Positive results are seen in liver diseases like hepatitis and cirrhosis and in cases of hemolytic anemia

ERYTHRO SEDIMENTATION RATE, BLOODErythrocyte sedimentation rate (ESR) is a non - specific phenomena and is clinically useful in the diagnosis and monitoring of disorders associated with an increased production of acute phase reactants. The ESR is increased in pregnancy from about the 3rd month and returns to normal by the 4th week post partum. ESR is influenced by age, sex, menstrual cycle and drugs (eg. corticosteroids, contraceptives). It is especially low (0-1mm) in polycythaemia, hypofibrinogenemia or congestive cardiac failure and when there are abnormalities of the red cells such as polikilocytosis, spherocytosis or sickle cells.

- 1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin
- 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th Edition"

GLYCOSYLATED HEMOGLOBIN, EDTA WHOLE BLOOD-Glycosylatec hemoglobin (GHb) has been firmly established as an index of long-term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. Formation of GHb is essentially irreversible, and the concentration in the blood depends on both the life span of the red blood cell (average 120 days) and the blood glucose concentration. Because the rate of formation of GHb is directly proportional to the concentration of glucose in the blood,

the GHb concentration represents the integrated values for glucose over the preceding 6-8 weeks.

Any condition that alters the life span of the red blood cells has the potential to alter the GHb level. Samples from patients with hemolytic anemias will exhibit decreased glycated hemoglobin values due to the shortened life span of the red cells. This effect will depend upon the severity of the anemia. Samples from patients with polycythemia

or post-splenectomy may exhibit increased glycated hemoglobin values due to a somewhat longer life span of the recicells.

Glycosylated hemoglobins results from patients with HbSS, HbCC, and HbSC and HbD must be interpreted with caution, given the pathological processes, including anemia, increased recicel turnover, transfusion requirements, that adversely impact HbA1c as a marker of long-term glycemic control. In these conditions, alternative forms of testing such as glycated serum protein (fructosamine) should be considered.
"Targets should be individualized; More or less stringent glycemic goals may be appropriate for individual patients. Goals should be individualized based on duration of

diabetes, age/life expectancy, comorbid conditions, known CVD or advanced microvascular complications, hypoglycemia unawareness, and individual patien considerations.

# References

- 1. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics, edited by Carl A Burtis, Edward R.Ashwood, David E Bruns, 4th Edition, Elsevier publication, 2006, 879-884.
- 2016-064.
  2. Forsham PH. Diabetes Mellitus: A rational plan for management. Postgrad Med 1982, 71,139-154.
  3. Mayer TK, Freedman ZR: Protein glycosylation in Diabetes Mellitus: A review of laboratory measurements and their clinical utility. Clin Chim Acta 1983, 127, 147-184.
  GLUCOSE, FASTING, PLASMAADA 2021 guidelines for adults, after 8 hrs fasting is as follows:

Pre-diabetics: 100 - 125 mg/dL Diabetic: > or = 126 mg/dL

GLUCOSE, POST-PRANDIAL, PLASMA-ADA Guidelines for 2hr post prandial glucose levels is only after ingestion of 75 grams of glucose in 300 ml water, over a period of 5

minutes. LIVER FUNCTION PROFILE, SERUM-

LIVER FUNCTION PROFILE

Bilirubin is a yellowish pigment found in bile and is a breakdowr product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. Elevatec levels results from increased bilirubir production (eg, hemolysis and ineffective erythropolesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg., hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors & Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

actaches sugar molecules to bilirubin.
AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood.ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health. AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis.

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Paget's disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Mainutrition, Protein deficiency, Wilson's disease. GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc. Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease. Lower-than-normal levels may be due to:



Page 9 Ot 12 Scan to View Report



ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI **NEW DELHI 110030** DELHI INDIA

8800465156

MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT ID:

HAREM180571181

Email: customercare.thane@srl.in

S.K. Tower, Hari Niwas, LBS Marg

SRL Ltd

THANE, 400602

PATIENT NAME: HARESH G GULRAJANI

ACCESSION NO: 0181VI000319 AGE: 51 Years SEX: Male

DRAWN: RECEIVED: 10/09/2022 09:57 REPORTED: 13/09/2022 15:18

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Results Biological Reference Interval Test Report Status Units <u>Final</u>

Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc. Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrobic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc SERUM BLOOD UREA NITROGEN-

Causes of Increased levels

Pre renal

- High protein diet, Increased protein databolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal
- Renal Failure Post Renal

· Malignancy, Nephrolithiasis, Prostatism

Causes of decreased levels

- Liver disease
- SIADH.

CREATININE, SERUM-

Higher than normal level may be due to:

- Blockage in the urinary tract
- Kidney problems, such as kidney damage or failure, infection, or reduced blood flow
   Loss of body fluid (dehydration)

- Muscle problems, such as breakdown of muscle fibers
   Problems during pregnancy, such as seizures (ediampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:
• Myasthenia Gravis

- Muscular dystrophy URIC ACID, SERUM-

Causes of Increasec levels

- Dietary
   High Protein Intake.
- Prolonged Fasting,
  Rapid weight loss.
- Gout

Lesch nyhan syndrome.

Type 2 DM.

Metabolic syndrome.

Causes of decreased levels

- Low Zinc Intake
- OCP's
- Multiple Sclerosis

Nutritional tips to manage increased Uric acid levels

- Drink plenty of fluids
   Limit animal proteins
   High Fibre foods
   Vit C Intake

Antioxidant rich foods
TOTAL PROTEIN, SERUM-

Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

ABUMIN, SERUMHuman serum albumir is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc. ELECTROLYTES (NA/K/CL), SERUM-

Sodium levels are Increased in dehydration, cushing's syndrome, aldosteronism & decreased in Addison's disease, hypopituitarism, liver disease. Hypokalemia (low K) is common in vomiting, diarrhea, alcoholism, folic acid deficiency and primary aldosteronism. Hyperkalemia may be seen in end-stage renal failure, hemolysis, trauma, Addison's disease, metabolic acidosis, acute starvation, dehydration, and with rapid K infusion. Chloride is increased in dehydration, renal tubular acidosis (hyperchloremia metabolic acidosis), acute renal failure, metabolic acidosis associated with prolonged diarrhea and loss of sodium bicarbonate, diabetes insipidus, adrenocortical hypertuction, salicylate intoxication and with excessive infusion of isotonic saline or extremely high dietary intake of salt. Chloride is decreased in overhydration, chronic respiratory acidosis, salt-losing nephritis, metabolic alkalosis, congestive heart failure, Addisonian crisis, certain types of metabolic acidosis, persistent gastric secretion and prolonaec vomitina.



Page 10 Ot 12 Scan to View Report



ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI **NEW DELHI 110030** DELHI INDIA 8800465156

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: HARESH G GULRAJANI

PATIENT ID: HAREM180571181

ACCESSION NO: 0181VI000319 AGE: 51 Years SEX: Male

DRAWN: RECEIVED: 10/09/2022 09:57 REPORTED: 13/09/2022 15:18

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Results Biological Reference Interval Test Report Status Units <u>Final</u>

### THYROID PANEL, SERUM-

Triiodothyronine T3 , is a thyroid hormone. It affects almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate. Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Thyroxine T4, Thyroxine's principal function is to stimulate the metabolism of all cells and tissues in the body. Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low.

Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3 Levels in TOTAL T4 TSH3G TOTAL T3

Pregnancy (µg/dL) (µIU/mL) (ng/dL) 0.1 - 2.5 0.2 - 3.0 0.3 - 3.0 81 - 190 First Trimester 6.6 - 12.4 2nd Trimester 6.6 - 15.5 100 - 260 100 - 260 3rc Trimester 6.6 - 15.5

Below mentioned are the guidelines for age related reference ranges for T3 and T4.

Т3 (ng/dL) New Born: 75 - 260 (µg/dL) 1-3 day: 8.2 **-** 19.9 1 Week: 6.0 - 15.9

NOTE: TSH concentrations in apparently normal euthyroid subjects are known to be highly skewed, with a strong tailed distribution towards higher TSH values. This is well documented in the pediatric population including the infant age group.

Kindly note: Method specific reference ranges are appearing or the report under biological reference range.

#### Reterence:

- 1. Burts C.A., Ashwood E. R. Bruns D.E. Teitz textbook of Clinical Chemistry and Molecular Diagnostics, 4th Edition.
  2. Gowenlock A.H. Varley's Practical Clinical Biochemistry, 6th Edition.

ABO GROUP & RH TYPE, EDTA WHOLE BLOODBlood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Flease note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same.

The test is performed by both forward as well as reverse grouping methods.

MEDICAL HISTORY-\*\*\*\* THIS REPORT CARRIES THE SIGNATURE OF OUR LABORATORY DIRECTOR, THIS IS AN INVIOLABLE FEATURE OF OUR LAB MANAGEMENT SOFTWARE, HOWEVER, ALL EXAMINATIONS AND INVESTIGATIONS HAVE BEEN CONDUCTED BY OUR PANEL OF DOCTORS.







CLIENT'S NAME AND ADDRESS:
ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )
F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL Ltd

S.K. Tower,Hari Niwas, LBS Marg THANE, 400602

MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

REPORTED:

PATIENT NAME: HARESH G GULRAJANI

PATIENT ID:

HAREM180571181

Test Report Status

ACCESSION NO: 0181VI000319 AGE: 51 Years

**Final** 

SEX: Male

13/09/2022 15:18

DRAWN:

RECEIVED: 10/09/2022 09:57

CLIENT PATIENT ID:

Biological Reference Interval

REFERRING DOCTOR: SELF

Results

Units

# MEDI WHEEL FULL BODY HEALTH CHECK UP ABOVE 40 MALE

ULTRASOUND ABDOMEN ULTRASOUND ABDOMEN GRADE | FATTY LIVER

GALL BLADDER POLYP WITH ECHOGENIC SLUDGE.

\*\*End Of Report\*\*

Please visit www.srlworld.com for related Test Information for this accession

Dr.Priyal Chinchkhede Consultant Pathologist

Bhinchkhede

Dr. Ushma Wartikar Consultant Pathologist Dr.(Mrs)Neelu K Bhojani

Lab Head



