

Name : Mrs. SHIKHA KUMARI

PID No. : MED111165270

SID No. : 222010939

Age / Sex : 36 Year(s) / Female

Type : OP

Ref. Dr : MediWheel

Register On : 16/06/2022 10:40 AM

Collection On : 16/06/2022 10:56 AM

Report On : 16/06/2022 6:31 PM

Printed On : 29/06/2022 12:56 PM



<u>Investigation</u>	<u>Observed Value</u>	<u>Unit</u>	<u>Biological Reference Interval</u>
----------------------	-----------------------	-------------	--------------------------------------

BLOOD GROUPING AND Rh TYPING	'AB' 'Positive'		
------------------------------	-----------------	--	--

(EDTA Blood/Agglutination)

**INTERPRETATION:** Reconfirm the Blood group and Typing before blood transfusion

**Complete Blood Count With - ESR**

Haemoglobin (EDTA Blood/Spectrophotometry)	<b>11.7</b>	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood/Derived from Impedance)	<b>35.0</b>	%	37 - 47
RBC Count (EDTA Blood/Impedance Variation)	<b>3.84</b>	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (EDTA Blood/Derived from Impedance)	91.1	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood/Derived from Impedance)	30.5	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood/Derived from Impedance)	33.4	g/dL	32 - 36
RDW-CV (EDTA Blood/Derived from Impedance)	13.0	%	11.5 - 16.0
RDW-SD (EDTA Blood/Derived from Impedance)	41.6	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood/Impedance Variation)	7400	cells/cu.mm	4000 - 11000
Neutrophils (EDTA Blood/Impedance Variation & Flow Cytometry)	65.8	%	40 - 75
Lymphocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	25.5	%	20 - 45
Eosinophils (EDTA Blood/Impedance Variation & Flow Cytometry)	1.6	%	01 - 06



VERIFIED BY



APPROVED BY

The results pertain to sample tested.

Page 1 of 7

Lab Address: MEDALL HEALTHCARE PRIVATE LIMITED,#17,RACE VIEW COLONY,2ND STREET, RACE COURSE ROAD, GUINDY, CHENNAI, TAMIL NADU, INDIA,.

Name : Mrs. SHIKHA KUMARI

PID No. : MED111165270

SID No. : 222010939

Age / Sex : 36 Year(s) / Female

Type : OP

Ref. Dr : MediWheel

Register On : 16/06/2022 10:40 AM

Collection On : 16/06/2022 10:56 AM

Report On : 16/06/2022 6:31 PM

Printed On : 29/06/2022 12:56 PM



<u>Investigation</u>	<u>Observed Value</u>	<u>Unit</u>	<u>Biological Reference Interval</u>
Monocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	6.8	%	01 - 10
Basophils (EDTA Blood/Impedance Variation & Flow Cytometry)	0.3	%	00 - 02
<b>INTERPRETATION:</b> Tests done on Automated Five Part cell counter. All abnormal results are reviewed and confirmed microscopically.			
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	4.9	10 <sup>3</sup> / $\mu$ l	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	1.9	10 <sup>3</sup> / $\mu$ l	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.1	10 <sup>3</sup> / $\mu$ l	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.5	10 <sup>3</sup> / $\mu$ l	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.0	10 <sup>3</sup> / $\mu$ l	< 0.2
Platelet Count (EDTA Blood/Impedance Variation)	227	10 <sup>3</sup> / $\mu$ l	150 - 450
MPV (EDTA Blood/Derived from Impedance)	10.7	fL	8.0 - 13.3
PCT (EDTA Blood/Automated Blood cell Counter)	0.242	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Blood/Automated - Westergren method)	2	mm/hr	< 20
BUN / Creatinine Ratio	13.1		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	79.3	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

  
**DR. GURUPRIYA J**  
PATHOLOGIST  
Reg No : 13-48036

VERIFIED BY

  
Dr. E. Saravanan M.D(Path)  
Consultant Pathologist  
Reg No : 73347

APPROVED BY

The results pertain to sample tested.

Page 2 of 7

Name : Mrs. SHIKHA KUMARI

PID No. : MED111165270

SID No. : 222010939

Age / Sex : 36 Year(s) / Female

Type : OP

Ref. Dr : MediWheel

Register On : 16/06/2022 10:40 AM

Collection On : 16/06/2022 10:56 AM

Report On : 16/06/2022 6:31 PM

Printed On : 29/06/2022 12:56 PM



<u>Investigation</u>	<u>Observed Value</u>	<u>Unit</u>	<u>Biological Reference Interval</u>
----------------------	-----------------------	-------------	--------------------------------------

**INTERPRETATION:** Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine) (Urine - F/GOD - POD)	Negative		Negative
---------------------------------------------------	----------	--	----------

Glucose Postprandial (PPBS) (Plasma - PP/GOD-PAP)	75.3	mg/dL	70 - 140
------------------------------------------------------	------	-------	----------

**INTERPRETATION:**

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

**Remark:** Please correlate clinically.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
-------------------------------------------	----------	--	----------

Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	8.0	mg/dL	7.0 - 21
----------------------------------------------------------	-----	-------	----------

Creatinine (Serum/Modified Jaffe)	0.61	mg/dL	0.6 - 1.1
--------------------------------------	------	-------	-----------

**INTERPRETATION:** Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetylcysteine, chemotherapeutic agent such as flucytosine etc.

Uric Acid (Serum/Enzymatic)	3.2	mg/dL	2.6 - 6.0
--------------------------------	-----	-------	-----------

**Liver Function Test**

Bilirubin(Total) (Serum/DCA with ATCS)	0.68	mg/dL	0.1 - 1.2
-------------------------------------------	------	-------	-----------

Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.26	mg/dL	0.0 - 0.3
---------------------------------------------------------	------	-------	-----------

Bilirubin(Indirect) (Serum/Derived)	0.42	mg/dL	0.1 - 1.0
----------------------------------------	------	-------	-----------

SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	24.4	U/L	5 - 40
----------------------------------------------------------------	------	-----	--------

**DR GURUPRIYA J**  
PATHOLOGIST  
Reg No : 13-48036

VERIFIED BY

**Dr. E. Saravanan M.D (Path)**  
Consultant Pathologist  
Reg No : 73347

APPROVED BY

Name : Mrs. SHIKHA KUMARI

PID No. : MED111165270

SID No. : 222010939

Age / Sex : 36 Year(s) / Female

Type : OP

Ref. Dr : MediWheel

Register On : 16/06/2022 10:40 AM

Collection On : 16/06/2022 10:56 AM

Report On : 16/06/2022 6:31 PM

Printed On : 29/06/2022 12:56 PM



<u>Investigation</u>	<u>Observed Value</u>	<u>Unit</u>	<u>Biological Reference Interval</u>
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	25.5	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	11.8	U/L	< 38
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	65.2	U/L	42 - 98
Total Protein (Serum/Biuret)	7.12	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.17	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	2.95	gm/dL	2.3 - 3.6
A : G RATIO (Serum/Derived)	1.41		1.1 - 2.2
<b><u>Lipid Profile</u></b>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	179.1	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	75.4	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

**INTERPRETATION:** The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the usual circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	<b>63.0</b>	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
---------------------------------------------	-------------	-------	--------------------------------------------------------------------------------



VERIFIED BY



APPROVED BY

Name : Mrs. SHIKHA KUMARI

PID No. : MED111165270

SID No. : 222010939

Age / Sex : 36 Year(s) / Female

Type : OP

Ref. Dr : MediWheel

Register On : 16/06/2022 10:40 AM

Collection On : 16/06/2022 10:56 AM

Report On : 16/06/2022 6:31 PM

Printed On : 29/06/2022 12:56 PM



<u>Investigation</u>	<u>Observed Value</u>	<u>Unit</u>	<u>Biological Reference Interval</u>
LDL Cholesterol (Serum/Calculated)	101	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	15.1	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	116.1	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

**INTERPRETATION:** 1.Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol.  
2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio (Serum/Calculated)	2.8	Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
---------------------------------------------------------------	-----	--------------------------------------------------------------------------------------------------------------------

Triglyceride/HDL Cholesterol Ratio (TG/HDL) (Serum/Calculated)	1.2	Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
----------------------------------------------------------------------	-----	------------------------------------------------------------------------

LDL/HDL Cholesterol Ratio (Serum/Calculated)	1.6	Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0
-------------------------------------------------	-----	-----------------------------------------------------------------

**Glycosylated Haemoglobin (HbA1c)**

HbA1C (Whole Blood/HPLC)	4.6 %	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5
-----------------------------	-------	-----------------------------------------------------------------

**INTERPRETATION:** If Diabetes - Good control : 6.1 - 7.0 % , Fair control : 7.1 - 8.0 % , Poor control >= 8.1 %



VERIFIED BY



APPROVED BY

The results pertain to sample tested.

Page 5 of 7

Name : Mrs. SHIKHA KUMARI

PID No. : MED111165270

SID No. : 222010939

Age / Sex : 36 Year(s) / Female

Type : OP

Ref. Dr : MediWheel

Register On : 16/06/2022 10:40 AM

Collection On : 16/06/2022 10:56 AM

Report On : 16/06/2022 6:31 PM

Printed On : 29/06/2022 12:56 PM



<u>Investigation</u>	<u>Observed Value</u>	<u>Unit</u>	<u>Biological Reference Interval</u>
Estimated Average Glucose (Whole Blood)	85.32	mg/dL	

**INTERPRETATION: Comments**

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glyceimic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.

**THYROID PROFILE / TFT**

T3 (Triiodothyronine) - Total (Serum/Chemiluminescent Immunometric Assay (CLIA))	1.03	ng/ml	0.7 - 2.04
----------------------------------------------------------------------------------------	------	-------	------------

**INTERPRETATION:**

**Comment :**

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Tyroxine) - Total (Serum/Chemiluminescent Immunometric Assay (CLIA))	6.87	µg/dl	4.2 - 12.0
--------------------------------------------------------------------------------	------	-------	------------

**INTERPRETATION:**

**Comment :**

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) (Serum/Chemiluminescent Immunometric Assay (CLIA))	2.28	µIU/mL	0.35 - 5.50
--------------------------------------------------------------------------------------------	------	--------	-------------

**INTERPRETATION:**

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5

2 nd trimester 0.2-3.0

3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

**Comment :**

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.

2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM.The variation can be of the order of 50%,hence time of the day has influence on the measured serum TSH concentrations.

3.Values&amplt;0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



VERIFIED BY



APPROVED BY

Name : Mrs. SHIKHA KUMARI

PID No. : MED111165270

SID No. : 222010939

Age / Sex : 36 Year(s) / Female

Type : OP

Ref. Dr : MediWheel

Register On : 16/06/2022 10:40 AM

Collection On : 16/06/2022 10:56 AM

Report On : 16/06/2022 6:31 PM

Printed On : 29/06/2022 12:56 PM



<u>Investigation</u>	<u>Observed Value</u>	<u>Unit</u>	<u>Biological Reference Interval</u>
<b><u>Urine Analysis - Routine</u></b>			
COLOUR (Urine)	Pale yellow		Yellow to Amber
APPEARANCE (Urine)	Clear		Clear
Protein (Urine/Protein error of indicator)	Negative		Negative
Glucose (Urine/GOD - POD)	Negative		Negative
Pus Cells (Urine/Automated $\delta$ Flow cytometry )	1 - 2	/hpf	NIL
Epithelial Cells (Urine/Automated $\delta$ Flow cytometry )	1 - 2	/hpf	NIL
RBCs (Urine/Automated $\delta$ Flow cytometry )	NIL	/hpf	NIL
Casts (Urine/Automated $\delta$ Flow cytometry )	NIL	/hpf	NIL
Crystals (Urine/Automated $\delta$ Flow cytometry )	NIL	/hpf	NIL
Others (Urine)	NIL		

**INTERPRETATION:**Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.

**DR GURUPRIYA J**  
PATHOLOGIST  
Reg No : 13-48036

VERIFIED BY

**Dr. E. Saravanan M.D (Path)**  
Consultant Pathologist  
Reg No : 73347

APPROVED BY

-- End of Report --