Age / Gender : 31 years / Female

Patient ID: 8363

Source : MEDI WHEEL

## Referral : SELF

Collection Time : Aug 19, 2022, 02:04 p.m.

Reporting Time : Aug 19, 2022, 05:37 p.m. Sample ID :



	004323122F		
Test Description	Value(s)	Reference Range	Unit
CBC; Complete Blood Count			
Hemoglobin (Hb)*	13.6	12.0 - 15.0	gm/dL
Method : Cynmeth Photometric Measurement			
Erythrocyte (RBC) Count*	4.7	3.8 - 4.8	mil/cu.mm
Method : Electrical Impedence			
Packed Cell Volume (PCV)*	42	36 - 46	%
Method : Calculated			
Mean Cell Volume (MCV)*	90	83 - 101	f∟
Method : Electrical Impedence			
Mean Cell Haemoglobin (MCH)*	29	27 - 32	pg
Method : Calculated			
Mean Corpuscular Hb Concn. (MCHC)*	31.9	31.5 - 34.5	gm/dL
Method : Calculated			
Red Cell Distribution Width (RDW)*	13.4	11.6 - 14.0	%
Method : Electrical Impedence			
Total Leucocytes (WBC) Count*	8400	4000-10000	cell/cu.mm
Method : Electrical Impedence			
Neutrophils*	49	40 - 80	%
Method : VCSn Technology			
Lymphocytes*	44	20 - 40	%
Method : VCSn Technology			
Monocytes*	6	2 - 10	%
Method : VCSn Technology			
Eosinophils*	1	1 - 6	%
Method : VCSn Technology			
Basophils	0	0 - 1	
Platelet Count*	2.58	1.5 - 4.5	10^3/ul
Method : Electrical Impedence			
Mean Platelet Volume (MPV)*	8.1	7.2 - 11.7	fL
Method : Electrical Impedence			

Auptin

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Approved by



Age / Gender : 31 years / Female

#### Patient ID: 8363

Source : MEDI WHEEL

# Referral : SELF

Collection Time : Aug 19, 2022, 02:04 p.m.

Reporting Time : Aug 19, 2022, 05:37 p.m. Sample ID :



Test Description	Value(s)	Reference Range	Unit
PCT*	0.2	0.2 - 0.5	%
Method : Calculated			
PDW*	17.7	9.0 - 17.0	%
Method : Calculated			

Tests done on Automated Three Part Cell Counter. (WBC, RBC, Platelet count by impedance method, colorimetric method for Hemoglobin, WBC differential by flow cytometry using laser technology other parameters are calculated). All Abnormal Haemograms are reviewed confirmed microscopically.

Esr, Erythrocyte Sedimentation Rate			
Esr, Erythrocyte Sedimentation Rate	31	0-30	mm/hr
(Westergren)			
Interpretation			

#### Interpretation:

- It indicates presence and intensity of an inflammatory process. It does not diagnose a specific disease. Changes in the ESR are
  more significant than the abnormal results of a single test.
- It is a prognostic test and used to monitor the course or response to treatment of diseases like tuberculosis, bacterial
  endocarditis, acute rheumatic fever, rheumatoid arthritis, SLE, Hodgkins disease, temporal arteritis and polymyalgia rheumatica.

"B" + (POSITIVE)

• It is also increased in pregnancy, multiple myeloma, menstruation, and hypothyroidism.

# Blood Group & Rh Type

#### **Blood Grouping & Rh Typing**

Method : Forward and Reverse By Tube Method

Methodology

This is done by forward and reverse grouping by tube Agglutination method.

#### Interpretation

Newborn baby does not produce ABO antibodies until 3 to 6 months of age. So the blood group of the Newborn baby is done by ABO antigen grouping (forward grouping) only, antibody grouping (reverse grouping) is not required.Confirmation of the New-born's blood group is indicatedwhen the A and B antigen expression and the isoagglutinins are fully developed (2–4 years).

appen

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Approved by



Age / Gender : 31 years / Female

Patient ID: 8363

Source : MEDI WHEEL

## Referral : SELF

Collection Time : Aug 19, 2022, 02:04 p.m.

Reporting Time : Aug 19, 2022, 05:37 p.m. Sample ID :



		004323122P	
Test Description	Value(s)	Reference Range	Unit
Fasting - Glucose			
Glucose Fasting* Method : Plasma, Hexokinase	73	Normal: 70-110 Impaired Fasting Glucose (IFG): 110-125 Diabetes Mellitus: >= 126 (On more than one occasion) (American Diabetes Association guidelines 2017)	mg/dL
Fasting Urine Sugar			
Fasting Urine Sugar	NEGATIVE	NEGATIVE -	
Lipid Profile			
Cholesterol-Total Method : Serum, Cholesterol oxidase esterase, peroxidase	125	Desirable: <= 200 Borderline High: 201-239 High: > 239 Ref: The National Cholesterol Education Program (NCEP) Adult Treatment Panel III Report.	mg/dL
Triglycerides Method : Serum, Enzymatic, endpoint	89	Normal: < 150 Borderline High: 150-199 High: 200-499 Very High: >= 500	mg/dL
Cholesterol-HDL Direct Method : Serum, Direct measure-PEG	45	Normal: > 40 Major Heart Risk: < 40	mg/dL

adapter

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Approved by



Age / Gender : 31 years / Female

#### Patient ID: 8363

Source : MEDI WHEEL

## Referral : SELF

Collection Time : Aug 19, 2022, 02:04 p.m.

Reporting Time : Aug 19, 2022, 05:37 p.m. Sample ID :



Test Description	Value(s)	Reference Range	Unit
LDL Cholesterol	62.2	Optimal: < 100	mg/dL
Method : Serum		Near optimal/above optimal: 100-12	9
		Borderline high: 130-159	
		High: 160-189	
		Very High: >= 190	
Non - HDL Cholesterol, Serum	80	Desirable: < 130 mg/dL	mg/dL
Method : calculated		Borderline High: 130-159mg/dL	
		High: 160-189 mg/dL	
		Very High: > or = 190 mg/dL	
VLDL Cholesterol	17.8	6 - 38	mg/dL
Method : calculated			
CHOL/HDL RATIO	2.78	3.5 - 5.0	ratio
Method : calculated			
LDL/HDL RATIO	1.38	Desirable / low risk - 0.5 -3.0	ratio
Method : calculated		Low/ Moderate risk - 3.0- 6.0	
		Elevated / High risk - > 6.0	
HDL/LDL RATIO	0.7	Desirable / low risk - 0.5 -3.0	ratio
Method : calculated		Low/ Moderate risk - 3.0- 6.0	
		Elevated / High risk - > 6.0	

#### **Liver Function Test** Bilirubin - Total 0.6 Adults and Children: < 1.2 mg/dL Method : Serum, Diazotization Adults and Children: < 0.5 0.2 mg/dL **Bilirubin - Direct** Method : Serum, Diazotization **Bilirubin - Indirect** 0.4 0.1 - 1.0 mg/dL Method : Serum, Calculated

abuptin

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Approved by



Age / Gender : 31 years / Female

Patient ID: 8363

Source : MEDI WHEEL

## Referral : SELF

Collection Time : Aug 19, 2022, 02:04 p.m.

Reporting Time : Aug 19, 2022, 05:37 p.m. Sample ID :



Test Description	Value(s)	Reference Range	Unit
SGOT	28	< 50	U/L
Method : Serum, UV with P5P, IFCC 37 degree			
SGPT	22	< 50	U/L
Method : Serum, UV with P5P, IFCC 37 degree			
Alkaline Phosphatase-ALPI	58	30-120	U/L
Method : Serum, PNPP, AMP Buffer, IFCC 37 degree			
Total Protein	7.7	6.6 - 8.3	g/dL
Method : Serum, Biuret, reagent blank end point			
Globulin	3.4	1.8 - 3.6	g/dL
Method : Calculated			
Albumin	4.3	Adults: 3.5 - 5.2	g/dL
Method : Serum, Bromcresol purple			
A/G Ratio	1.26	1.2 - 2.2	ratio
Method : Calculated			
GGT-Gamma Glutamyl Transpeptidae	18	< 55	U/L
Method : Serum, G-glutamyl-carboxy-nitoanilide			
KIDNEY FUNCTION TEST			
Urea *	20	15- 50	mg/dL
Method : Serum			
Blood Urea Nitrogen-BUN*	9.3	7 - 24	mg/dL
Method : Serum, Urease			
Uric Acid*	4.9	3.5 - 7.2	mg/dL
Method : Serum, Uricase/POD			
Creatinine*	0.7	0.6 - 1.1	mg/dL
Method : Serum, Jaffe IDMS			

Colour\* Transparency (Appearance)\*

Pale Yellow Clear Pale Yellow Clear

Auptin

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Approved by



Age / Gender : 31 years / Female

#### Patient ID: 8363

Source : MEDI WHEEL

# Referral : SELF

Collection Time : Aug 19, 2022, 02:04 p.m.

Reporting Time : Aug 19, 2022, 05:37 p.m. Sample ID :



Test Description	Value(s)	Reference Range	Unit
Deposit*	Absent	Absent	
Reaction (pH)*	5.0	4.5 - 8	
Specific Gravity*	1.025	1.010 - 1.030	
Chemical Examination (Automated D	ipstick Method) Urine		
Urine Glucose (sugar)*	Absent	Absent	
Urine Protein (Albumin)*	Absent	Absent	
Urine Ketones (Acetone)*	Absent	Absent	
Blood*	Absent	Absent	
Bile pigments*	Absent	Absent	
Nitrite*	Absent	Absent	
Urobilinogen*	Normal	Normal	
Microscopic Examination Urine			
Pus Cells (WBCs)*	2-3	0 - 5	/hpf
Epithelial Cells*	0-1	0 - 4	/hpf
Red blood Cells*	Absent	Absent	/hpf
Crystals*	Absent	Absent	
Cast*	Absent	Absent	
Trichomonas Vaginalis*	Absent	Absent	
Yeast Cells*	Absent	Absent	
Amorphous deposits*	Absent	Absent	
Bacteria*	Absent	Absent	

5.2

Glyco Hb (HbA1C)	
Method : EDTA Whole blood, HPLC	

Non-Diabetic: <=5.9 Pre Diabetic:6.0-6.4 Diabetic: >=6.5 %

abuptin

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Approved by



Age / Gender : 31 years / Female Patient ID: 8363 Sample ID : Source : MEDI WHEEL **Test Description** Value(s) **Reference Range** Estimated Average Glucose : 102 Interpretations 1. HbA1C has been endorsed by clinical groups and American Diabetes Association guidelines 2017 for diagnosing diabetes using a cut off point of 6.5% 2. Low glycated haemoglobin in a non diabetic individual are often associated with systemic inflammatory diseases, chronic anaemia (especially severe iron deficiency and haemolytic), chronic renal failure and liver diseases. Clinical correlation suggested. 3. In known diabetic patients, following values can be considered as a tool for monitoring the glycemic control. Poor Control – More than 10 % **Thyroid Function Test (TFT)** 

Referral : SELF

Collection Time : Aug 19, 2022, 02:04 p.m.

Reporting Time : Aug 19, 2022, 05:37 p.m.

Unit

mg/dL



Excellent control-6-7 %	
Fair to Good control – 7-8 %	
Unsatisfactory control – 8 to 10 %	

Patient Name : MRS. M. THARMILA RANI

THYROID STIMULATING HORMONE (TSH)	7.0	0.46 – 8.10 : 1 Yrs – 5 Yrs	uIU/mL
Method : CLIA		0.36 – 5.80 : 6 Yrs – 18 Yrs	
		0.35 – 5.50 : 18 Yrs – 55 Yrs	
		0.50 – 8.90 : >55 Yrs	
		Pregnancy Ranges::::	
		lst Tri :0.1 - 2.5	
		llnd Tri :0.2 - 3.0	
		IIIrd Tri:0.3 - 3.0	
TOTAL TRIIODOTHYRONINE (T3)	132	126 – 258 : 1 Yr – 5 Yr	ng/dl
Method : CLIA		96 – 227 : 6 Yr – 15 Yr	
		91 – 164 : 16 Yr – 18 Yr	
		60 – 181 : > 18 Years	
		Pregnancy :	
		1st Trimester : 81 - 190	
		2nd & 3rd Trimester:100 - 260	

Auptin

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Approved by



Age / Gender : 31 years / Female

Patient ID: 8363

Source : MEDI WHEEL

Referral : SELF

Collection Time : Aug 19, 2022, 02:04 p.m.

Reporting Time : Aug 19, 2022, 05:37 p.m. Sample ID :



Test Description	Value(s)	Reference Range	Unit			
TOTAL THYROXINE (T4)	5.5		μg/dL			
Method : CLIA		4.6 - 10.9				
		Pregnancy:				
		4.6 – 16.5 : 1st Trimester				
		4.6 – 18.5 : 2nd & 3rd Tri				
Comments:	IF NOT ON I	DRUGS SUGGESTED FT3 & FT4 I	ESTIMATION			
	Please corre	elate with clinical conditions.				
	Note : Serum	Note : Serum T3, T4 and TSH form the three components of thyroid screening				
	panel, useful in diagnosing various disorders of the thyroid gland. Primary					
	Hypothyroidism is accompanied by depressed serum T3 and T4 values and					
	elevated serum TSH levels. Although elevated TSH levels are nearly always					
	indicative of Primary Hypothyroidism, rarely they can from TSH secreting					
	pituitary tumors (Secondary hyperthyroidism)To confirm diagnosis - evaluate					
	FT3 and FT4					
Post Prandial Urine Sugar						
Post Prandial Urine Sugar	NEGATIVE					
Post Prandial Blood Sugar						
Blood Glucose-Post Prandial*	54	70-140	mg/dL			
Method : Plasma - P, Hexokinase						

\*\*END OF REPORT\*\*

Auptin

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Approved by

