Buy Now to Create PDF with wal Watermark!!

MC-5333

PATIENT NAME: RAM JI LAL MEENA REF. DOCTOR: SELF

CODE NAME & ADDRESS: C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO: **0251WA001949** PATIENT ID: RAMJM280170251

CLIENT PATIENT ID: 012301280017 ABHA NO : AGE/SEX :53 Years Male
DRAWN :28/01/2023 09:38:00
RECEIVED :28/01/2023 10:13:50
REPORTED :28/01/2023 16:06:26

Test Report Status Final Results Biological Reference Interval Units

	HAEMATOLOGY - CBC		
MEDI WHEEL FULL BODY HEALTH CHECK UP A	ABOVE 40 MALE		
BLOOD COUNTS,EDTA WHOLE BLOOD			
HEMOGLOBIN (HB) METHOD: CYANIDE FREE DETERMINATION	15.8	13.0 - 17.0	g/dL
RED BLOOD CELL (RBC) COUNT METHOD: ELECTRICAL IMPEDANCE	4.33 Low	4.5 - 5.5	mi l /μL
WHITE BLOOD CELL (WBC) COUNT METHOD: ELECTRICAL IMPEDANCE	6 . 90	4.0 - 10.0	thou/μL
PLATELET COUNT METHOD: ELECTRONIC IMPEDANCE	170	150 - 410	thou/μL
RBC AND PLATELET INDICES			
HEMATOCRIT (PCV) METHOD: CALCULATED PARAMETER	47.8	40 - 50	%
MEAN CORPUSCULAR VOLUME (MCV) METHOD: CALCULATED PARAMETER	109,0 High	83 - 101	fL
MEAN CORPUSCULAR HEMOGLOBIN (MCH) METHOD: CALCULATED PARAMETER	36.4 High	27.0 - 32.0	pg
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION (MCHC) METHOD: CALCULATED PARAMETER	33.5	31.5 - 34.5	g/dL
RED CELL DISTRIBUTION WIDTH (RDW) METHOD: CALCULATED PARAMETER	12.2	11.6 - 14.0	%
MENTZER INDEX	25.2	$\langle \langle \langle \rangle \rangle$	
MEAN PLATELET VOLUME (MPV) METHOD: CALCULATED PARAMETER	11.3 High	6.8 - 10.9	fL
WBC DIFFERENTIAL COUNT			
NEUTROPHILS METHOD: IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY	53	40 - 80	%
LYMPHOCYTES METHOD: IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY	36	20 - 40	%
MONOCYTES METHOD: IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY	06	2 - 10	%
EOSINOPHILS METHOD: IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY	05	1 - 6	%

A CONTRACTOR OF THE PARTY OF TH


Dr. Akansha Jain Consultant Pathologist



Page 1 Of 14

View Details

View Report





MC-5333

PATIENT NAME: RAM JI LAL MEENA

CODE NAME & ADDRESS : C000049066 SRL JATPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

REF. DOCTOR: SELF

ACCESSION NO: 0251WA001949 PATIENT ID : RAMJM280170251

CLIENT PATIENT ID: 012301280017

ABHA NO

AGE/SEX :53 Years DRAWN :28/01/2023 09:38:00 RECEIVED: 28/01/2023 10:13:50 REPORTED :28/01/2023 16:06:26

Test Report Status <u>Final</u>	Results	Biological Reference	Interval Units
BASOPHILS METHOD: IMPEDANCE WITH HYDRO-FOCUS AND MICROSCO	00	0 - 2	%
ABSOLUTE NEUTROPHIL COUNT METHOD: CALCULATED PARAMETER	3.66	2.0 - 7.0	thou/μL
ABSOLUTE LYMPHOCYTE COUNT METHOD: CALCULATED PARAMETER	2.48	1.0 - 3.0	thou/µL
ABSOLUTE MONOCYTE COUNT METHOD: CALCULATED PARAMETER	0.41	0.2 - 1.0	thou/µL
ABSOLUTE EOSINOPHIL COUNT METHOD: CALCULATED PARAMETER	0.34	0.02 - 0.50	thou/µL
ABSOLUTE BASOPHIL COUNT NEUTROPHIL LYMPHOCYTE RATIO (NLR)	0 Low 1.5	0.02 - 0.10	thou/µL

Interpretation(s)
BLOOD COUNTS,EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs, However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13)

from Beta thalassaemia trait

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical contellation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

was understand that the second of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR = 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients; A-Parang, et al.; International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope.



Dr. Akansha Jain Consultant Pathologist Page 2 Of 14







Buy Now to Create PDF without Trial Watermar

PATIENT NAME: RAM JI LAL MEENA

CODE NAME & ADDRESS : C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN

AAKRITI LABS PVT LTD. A-430, AGRASEN MARG JAIPUR 302017

9314660100

REF. DOCTOR: SELF

ACCESSION NO: 0251WA001949

: RAMJM280170251

CLIENT PATIENT ID: 012301280017

ABHA NO

AGE/SEX :53 Years Male DRAWN :28/01/2023 09:38:00

RECEIVED: 28/01/2023 10:13:50 REPORTED :28/01/2023 16:06:26

Test Report Status

Final

Results

Biological Reference Interval

Units

HAEMATOLOGY

MEDI WHEEL FULL BODY HEALTH CHECK UP ABOVE 40 MALE

ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE

BLOOD

E.S.R 0 - 14METHOD: AUTOMATED (PHOTOMETRICAL CAPILLARY STOPPED FLOW KINETIC ANALYSIS)"

mm at 1 hr

Interpretation(s)

ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD-TEST DESCRIPTION :-

Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowaday's fully automated instruments are available to measure ESR.

ESR is not diagnostic; it is a non-specific test that may be elevated in a ruprber of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change. **TEST INTERPRETATION**

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Areignia Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy, Estrogen medication, Aging,

Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined syriptons directs the physician to search for a systemic disease (Paraproteinemias,

Disseminated malignancies, connective tissue disease, severe infections such as vacterial endocarditis).

In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum. Decreased in: Polycythermia vera, Sickle cell anemia

False elevated ESR: Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholester olemia
False Decreased: Poikilocytosis,(SickleCells,spherocytes),Microcytosis, Low fibrinogen, Very High WBC counts, Drugs(Quinine, salicylates)

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition; 2. Paediatric reference intervals. 🕰 Press, 7th edition. Edited by S. Soldin; 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th edition.

Dr. Akansha Jain Consultant Pathologist Page 3 Of 14





PERFORMED AT:

C/o Aakriti Labs P C td. 3. Mahatma Gardhi Marg,Gandhi Rajasthan, INDIA



Buy Now to Create PDF without Trial Watermar

PATIENT NAME: RAM JI LAL MEENA

CODE NAME & ADDRESS : C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

REF. DOCTOR: SELF

ACCESSION NO: 0251WA001949

: RAMJM280170251 CLIENT PATIENT ID: 012301280017

ABHA NO

AGE/SEX :53 Years Male DRAWN :28/01/2023 09:38:00 RECEIVED: 28/01/2023 10:13:50

REPORTED :28/01/2023 16:06:26

Test Report Status

Final

Results

Biological Reference Interval Units

IMMUNOHAEMATOLOGY

MEDI WHEEL FULL BODY HEALTH CHECK UP ABOVE 40 MALE

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

ABO GROUP

RH TYPF

TYPE B

METHOD: TUBE AGGLUTINATION

METHOD: TUBE AGGLUTINATION

POSITIVE

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-

agnant women. Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Croup) for pregnant women are not available, please check with the patient records for availability of the same."

The test is performed by both forward as well as reverse grouping methods.

Page 4 Of 14





PERFORMED AT:

Dr. Akansha Jain Consultant Pathologist

C/o Aakriti Labs Prestd. 3. Mahatma Gardhi Marg,Gandhi Rajasthan, INDIA



MC-5333

PATIENT NAME: RAM JI LAL MEENA

CODE NAME & ADDRESS : C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

REF. DOCTOR: SELF

ACCESSION NO: 0251WA001949 AGE/SEX

: RAMJM280170251

CLIENT PATIENT ID: 012301280017

ABHA NO

PATIENT ID

:53 Years Male DRAWN :28/01/2023 09:38:00 RECEIVED: 28/01/2023 10:13:50

REPORTED :28/01/2023 16:06:26

Test Report Status

Final

Results

Biological Reference Interval Units

BIOCHEMISTRY

MEDI WHEEL FULL BODY HEALTH CHECK UP ABOVE 40 MALE

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE

BLOOD HBA1C

6.6 High

Non-diabetic: < 5.7

Pre-diabetics: 5.7 - 6.4 Diabetics: > or = 6.5Therapeutic goals: < 7.0 Action suggested: > 8.0

(ADA Guideline 2021)

METHOD: HIGH PERFORMANCE LIQUID CHROMATOGRAPHY (HPLC)

ESTIMATED AVERAGE GLUCOSE(EAG)

METHOD: CALCULATED PARAMETER

142.7 High

< 116.0

mg/dL

%

GLUCOSE FASTING, FLUORIDE PLASMA

FBS (FASTING BLOOD SUGAR)

METHOD: GLUCOSE OXIDASE

173 High

74 - 99

mg/dL

GLUCOSE, POST-PRANDIAL, PLASMA

PPBS(POST PRANDIAL BLOOD SUGAR)

240 High

70 - 140

mg/dL

METHOD: GLUCOSE OXIDASE LIPID PROFILE, SERUM

CHOLESTEROL, TOTAL

159

< 200 Desirable

mg/dL

200 - 239 Borderline High

>/= 240 High

METHOD: CHOLESTEROL OXIDASE

TRIGLYCERIDES

HDL CHOLESTEROL

126

★ 150 Normal

mg/dL

150 - 199 Borderline High

200 - 499 High

≥/≠500 Very High

43

< 4040W >/=60 High

mg/dL

METHOD: DIRECT CLEARANCE METHOD

METHOD: LIPASE/GPO-PAP NO CORRECTION

Dr. Akansha Jain Consultant Pathologist Page 5 Of 14







MC-5333

PATIENT NAME: RAM JI LAL MEENA **REF. DOCTOR:** SELF

CODE NAME & ADDRESS : C000049066 SRL JATPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

ACCESSION NO: 0251WA001949 PATIENT ID : RAMJM280170251 CLIENT PATIENT ID: 012301280017

ABHA NO

AGE/SEX :53 Years Male :28/01/2023 09:38:00 DRAWN RECEIVED: 28/01/2023 10:13:50 REPORTED :28/01/2023 16:06:26


		<u> </u>	
Test Report Status <u>Final</u>	Results	Biological Reference In	terval Units
CHOLESTEROL LDL	91	< 100 Optimal 100 - 129	mg/dL
		Near optimal/ above op	timal
	\supset	130 - 159	
	3	Borderline High	
)55	160 - 189 High >/= 190 Very High	
NON HDL CHOLESTEROL	116	Desirable: Less than 13	30 mg/dL
		Above Desirable: 130 -	
		Borderline High: 160 -	189
	v v	High: 190 - 219 Very high: > or = 220	
METHOD: CALCULATED PARAMETER		very mgm. > 01 = 220	
VERY LOW DENSITY LIPOPROTE	Z5.2	= 30.0</td <td>mg/dL</td>	mg/dL
CHOL/HDL RATIO	3,7	3.3 - 4.4	
		Low Risk 4.5 - 7.0	
	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	Average Risk	
		7.1 - 11.0	
		Moderate Risk > 11.0	
		High Risk	
LDL/HDL RATIO	2.1	0.5 - 3.0 Desirable/Low	Risk
		3.1 - 6.0 Borderline/Mo	derate
		Risk >6.0 High Risk	
Interpretation(s)	V	A COLO HIGH KISK	
		(\bigcirc) \bigcirc	
LIVER FUNCTION PROFILE, SER	UM		
BILIRUBIN, TOTAL	0.94	0-1	mg/dL
METHOD: DIAZO WITH SULPHANILIC ACID	0.00 11111	2.25	
BILIRUBIN, DIRECT METHOD: DIAZO WITH SULPHANILIC ACID	0.32 High	0.00 - 0.25	mg/dL
BILIRUBIN, INDIRECT	0.62	0.1 - 1.0	mg/dL
METHOD : CALCULATED PARAMETER	3.32		5, -
TOTAL PROTEIN	8.2	6.4 - 8.2	g/dL
METHOD: BIURET REACTION, END POINT			
ALBUMIN	4.9 High	3.8 - 4.4	g/dL
			9)

Dr. Akansha Jain **Consultant Pathologist**





Page 6 Of 14



Buy Now to Create PDF with watermark!!

MC-5333

PATIENT NAME: RAM JI LAL MEENA REF. DOCTOR: SELF

CODE NAME & ADDRESS : C000049066

SRL JATPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO: **0251WA001949**PATIENT ID: RAMJM280170251
CLIENT PATIENT ID: 012301280017

ABHA NO :

AGE/SEX :53 Years Male
DRAWN :28/01/2023 09:38:00
RECEIVED :28/01/2023 10:13:50
REPORTED :28/01/2023 16:06:26

	<u> </u>	<u> </u>	
Test Report Status <u>Final</u>	Results	Biological Reference Interv	al Units
METHOD : BROMOCRESOL GREEN			
GLOBULIN	3.3	2.0 - 4.1	g/dL
METHOD : CALCULATED PARAMETER			
ALBUMIN/GLOBULIN RATIO	1.5	1.0 - 2.1	RATIO
METHOD: CALCULATED PARAMETER			
ASPARTATE AMINOTRANSFERASE (AST/SGOT) METHOD: TRIS BUFFER NO P5P IFCC / SFBC 37	31	0 - 37	U/L
ALANINE AMINOTRANSFERASE (ALTISOPT) METHOD: TRIS BUFFER NO P5P IFCC / SFBC 37° C	38	0 - 40	U/L
ALKALINE PHOSPHATASE METHOD: AMP OPTIMISED TO IFCC 37° C	103	39 - 117	U/L
GAMMA GLUTAMYL TRANSFERASE (GGT) METHOD: GAMMA GLUTAMYL-3 CARBOXY-4 NITROANILIDE (IFCC)	65 High	11 - 50	U/L
LACTATE DEHYDROGENASE	346	230 - 460	U/L
BLOOD UREA NITROGEN (BUN), SERUM			
BLOOD UREA NITROGEN	9	5.0 - 18.0	mg/dL
METHOD : UREASE KINETIC	(\bigcirc)		
CREATININE, SERUM			
CREATININE	0.99	0.8 - 1.3	mg/dL
METHOD: ALKALINE PICRATE NO DEPROTEINIZATION			
BUN/CREAT RATIO	Sym		
BUN/CREAT RATIO	9.09		
METHOD : CALCULATED PARAMETER	\wedge	(\mathcal{O})	
URIC ACID, SERUM	3.0	7.0	ma/dl
URIC ACID METHOD: URICASE PEROXIDASE WITH ASCORBATE OXIDASE	3.9	3,4 - 7.0	mg/dL
TOTAL PROTEIN, SERUM			
TOTAL PROTEIN	8.2	6.4 - 8.3	g/dL
METHOD : BIURET REACTION, END POINT	0.2	0.4 0.9	9, 42
ALBUMIN, SERUM			
ALBUMIN	4.9 High	3.8 - 4.4	g/dL
METHOD: BROMOCRESOL GREEN	-		5.
GLOBULIN			
GLOBULIN	3.3	2.0 - 4.1	g/dL
			>



Dr. Akansha Jain Consultant Pathologist

Page 7 Of 14

View Details

View Report



MC-5333

PATIENT NAME: RAM JI LAL MEENA

CODE NAME & ADDRESS : C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

REF. DOCTOR: SELF ACCESSION NO: 0251WA001949

PATIENT ID : RAMJM280170251

CLIENT PATIENT ID: 012301280017

ABHA NO

AGE/SEX :53 Years DRAWN :28/01/2023 09:38:00 RECEIVED: 28/01/2023 10:13:50 REPORTED: 28/01/2023 16:06:26

Test Report Status <u>Final</u>	Results	Biological Reference	e Interval Units
ELECTROLYTES (NA/K/CL), SERUM			
SODIUM, SERUM METHOD: ION-SELECTIVE ELECTROPE	138.7	137 - 145	mmo l /L
POTASSIUM, SERUM METHOD: ION-SELECTIVE ELECTRODE	3.94	3.6 - 5.0	mmol/L
CHLORIDE, SERUM METHOD: ION-SELECTIVE ELECTRODE	98.5	98 - 107	mmo l /L
Interpretation(s)			

Interpretation(s)

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-Used For

- 1.Evaluating the long-term control of blood glucose concentrations in diabetic patients.
- 3.Identifying patients at increased risk for diabetes (prediabetes).

The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1) and poorly controlled type 2 diabetic patients, and 2 times per year for well-controlled type 2 diabetic patients) to determine whether a patients metabolic control has remained continuously within the target range.

1.eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.

- 2. eAG gives an evaluation of blood glucose levels for the last couple of months.

 3. eAG is calculated as eAG (mg/dl) = 28.7 * HbA1c 46.7

HbA1c Estimation can get affected due to: I.Shortened Erythrocyte survival: Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss, hemolytic anemia) will falsely lower HbA1c test results. Fructosamine is recommended in these patients which indicates diabetes control over 15 days.

II.Vitamin C & E are reported to falsely lower test results (possibly by inhibiting glycation of hemoglobics.
III.Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilifubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates addiction are reported to interfere with some assay methods, falsely increasing results. IV.Interference of hemoglobinopathies in HbA1c estimation is seen in

a.Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c.

b. Heterozygous state detected (D10 is corrected for HbS & HbC trait.)

c.HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c. Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy GLUCOSE FASTING,FLUORIDE PLASMA-TEST DESCRIPTION

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readify available to tissues and sothat no glucose is excreted in the urine

Increased in

Diabetes mellitus, Cushing's syndrome (10 – 15%), chronic pancreatitis (30%). Drugs:corticosteroids,phenytoin, estrogen, thiazides.

Decreased in

Pancreatic islet cell disease with increased insulin,insulinoma,adrenocortical insufficiency, hypopituitarism,diffuse liver disease, malignancy (adrenocortical, stomach,fibrosarcoma), infant of a diabetic mother, enzyme deficiency diseases(e.g., galactosemia),Drugs- insulin, ethanol, propranolol; sulfonylureas,tolbutamide, and other oral hypoglycemic agents.

NOTE:

While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values), there المرافعة والمرافعة والمرافعة المرافعة المرافعة والمرافعة المرافعة المرافع glycosylated hemoglobin(HbA1c) levels are favored to monitor glycemic control.

glycosylated nemoglobin(HBALC) levels are ravored to monitor glycemic control. High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.

GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc. Additional test HbA1c LIVER FUNCTION PROFILE, SERUM-LIVER FUNCTION PROFILE



Consultant Pathologist





Page 8 Of 14

View Report

Buy Now to Create PDF with 🚳 🔞 al

MC-5333

PATIENT NAME: RAM JI LAL MEENA

CODE NAME & ADDRESS : C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

REF. DOCTOR: SELF

ACCESSION NO: 0251WA001949 AGE/SEX :53 Years Male

DRAWN PATIENT ID : RAMJM280170251

CLIENT PATIENT ID: 012301280017 ABHA NO

:28/01/2023 09:38:00 RECEIVED: 28/01/2023 10:13:50

REPORTED: 28/01/2023 16:06:26

Test Report Status

Final

Biological Reference Interval Units

Bilirubin is a yellowish pigment ինսից in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abjorma bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, (Dryo/reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the tody. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured

Results

clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney fallure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis

hepatitis, obstruction of bile ducts, cirrhosis.

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Paget'''s disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilson'''s disease. GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intesting spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are postructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc. Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammatic bleeding (hemorrhage). Burns, Glomerulonephritis. Liver disease. Malabsoration. Malnutrition. Nephrotic. disease Lower-than-normal levels may be due to: Agammaglobulinemia, legding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc. Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein Low blood albumin levels (hypoalbuminemia) can be caused by:Liver disease like cirrhosis of the liver, nephrotic syndrome,protein-losing

enteropathy, Burns, hemodilution, increased vascular permeability or degreased lymphatic clearance, malnutrition and wasting etc

BLOOD UREA NITROGEN (BUN), SERUM-Causes of Increased levels include Prevenal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)

Causes of decreased level include Liver disease, SIADH.

CREATININE, SERUM-Higher than normal level may be due to:

- Blockage in the urinary tract
 Kidney problems, such as kidney damage or failure, infection, or reduced blood flow
- Loss of body fluid (dehydration)
 Muscle problems, such as breakdown of muscle fibers
- Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:

- Myasthenia GravisMuscular dystrophy

URIC ACID, SERUM-Causes of Increased levels:-Dietary(High Protein Intake,Prolonged Fasting, Kapid weight loss),Gout,Lesch nyhan syndrome,Type 2 DM,Metabolic syndrome

Causes of decreased levels-Low Zinc intake,OCP,Multiple Sclerosis

TOTAL PROTEIN, SERUM-Serum total protein,also known as total protein, is a biochemical test for measyring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin

Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage),Burns,Glomerulonephitis/Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

ALBUMIN, SERUM-Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum

protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.

Dr. Akansha Jain

Consultant Pathologist

Page 9 Of 14





View Report

MC-5333

PATIENT NAME: RAM JI LAL MEENA

CODE NAME & ADDRESS: C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100 REF. DOCTOR: SELF

ACCESSION NO: **0251WA001949** AGE/SEX :53 Years
PATIENT ID : RAMJM280170251 DRAWN :28/01/20

CLIENT PATIENT ID: 012301280017

ABHA NO :

DRAWN :28/01/2023 09:38:00 RECEIVED :28/01/2023 10:13:50 REPORTED :28/01/2023 16:06:26

Male

Test Report Status <u>Final</u> Results Biological Reference Interval Units

CLINICAL PATH - URINALYSIS

MEDI WHEEL FULL BODY HEALTH CHECK UP ABOVE 40 MALE

PHYSICAL EXAMINATION, URINE

COLOR PALE YELLOW

METHOD : GROSS EXAMINATION

APPEARANCE SLIGHTLY HAZY

METHOD : GROSS EXAMINATION

CHEMICAL EXAMINATION, URINE

PH 6.0 4.7 - 7.5

METHOD: DOUBLE INDICATOR PRINCIPLE

SPECIFIC GRAVITY = 1.005 1.003 - 1.035

METHOD: IONIC CONCENTRATION METHOD

PROTEIN NOT DETECTED NOT DETECTED

METHOD: PROTEIN ERROR OF INDICATORS WITH REFLECTANCE

GLUCOSE

MOT DETECTED

NOT DETECTED

METHOD: GLUCOSE OXIDASE PEROXIDASE / BENEDICTS

KETONES NOT DETECTED NOT DETECTED

METHOD: SODIUM NITROPRUSSIDE REACTION

BLOOD

NOT DETECTED

NOT DETECTED

METHOD : PEROCIDASE ANTI PEROXIDASE

BILIRUBIN NOT DETECTED NOT DETECTED

UROBILINOGEN NORMAL NORMAL

METHOD: EHRLICH REACTION REFLECTANCE

NITRITE NOT DETECTED NOT DETECTED

METHOD: NITRATE TO NITRITE CONVERSION METHOD

LEUKOCYTE ESTERASE DETECTED NOT DETECTED

MICROSCOPIC EXAMINATION, URINE

RED BLOOD CELLS NOT DETECTED NOT DETECTED /HPF

METHOD: MICROSCOPIC EXAMINATION

PUS CELL (WRC'S)

5-7

0-5

PUS CELL (WBC'S) 5-7 0-5 /HPF

METHOD: DIPSTICK, MICROSCOPY

EPITHELIAL CELLS

1-2

0-5

PITHELIAL CELLS 1-2 0-5 /HPF
METHOD: MICROSCOPIC EXAMINATION

CASTS NOT DETECTED

1

METHOD: DIPSTICK

Dr. Akansha Jain Consultant Pathologist



Page 10 Of 14

View Details

View Report

PERFORMED AT:

JAIPUR, 302015 Created by eDocPrinter PDF Rajasthan, INDIA



MC-5333

PATIENT NAME: RAM JI LAL MEENA

CODE NAME & ADDRESS : C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

REF. DOCTOR: SELF

ACCESSION NO: 0251WA001949

PATIENT ID : RAMJM280170251 CLIENT PATIENT ID: 012301280017

ABHA NO

AGE/SEX :53 Years Male DRAWN :28/01/2023 09:38:00 RECEIVED: 28/01/2023 10:13:50 REPORTED :28/01/2023 16:06:26

Units

Test Report Status **Biological Reference Interval** Results <u>Final</u>

METHOD : MICROSCOPIC EXAMINATION

CRYSTALS

METHOD: MICROSCOPIC EXAMINATION

BACTERIA METHOD: MICROSCOPIC EXAMINATION

YEAST

Interpretation(s)

NOT DETECTED

N NOT.

Dr. Akansha Jain Consultant Pathologist

Page 11 Of 14

MC-5333

PATIENT NAME: RAM JI LAL MEENA

CODE NAME & ADDRESS : C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

REF. DOCTOR: SELF

ACCESSION NO: 0251WA001949

: RAMJM280170251

CLIENT PATIENT ID: 012301280017

ABHA NO

PATIENT ID

AGE/SEX :53 Years Male DRAWN :28/01/2023 09:38:00 RECEIVED: 28/01/2023 10:13:50

REPORTED :28/01/2023 16:06:26

Test Report Status

Final

Results

Biological Reference Interval Units

CLINICAL PATH - STOOL ANALYSIS

MEDI WHEEL FULL BODY HEALTH CHECK UP ABOVE 40 MALE

PHYSICAL EXAMINATION, STOOL

COLOUR METHOD: GROSS EXAMINATION

Dr. Abhishek Sharma Consultant Microbiologist



Page 12 Of 14

PERFORMED AT:

C/o Aakriti Labs Pyc'td, 3. Mahatma Gar dhi Marg,Gandhi Rajasthan, INDIA



MC-5333

PATIENT NAME: RAM JI LAL MEENA **REF. DOCTOR:** SELF

CODE NAME & ADDRESS : C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

ACCESSION NO: 0251WA001949 PATIENT ID : RAMJM280170251 CLIENT PATIENT ID: 012301280017

ABHA NO

AGE/SEX :53 Years Male DRAWN :28/01/2023 09:38:00 RECEIVED: 28/01/2023 10:13:50 REPORTED: 28/01/2023 16:06:26

Test Report Status Results **Biological Reference Interval** Units **Final**

SPECIALISED CHEMISTRY - HORMONE

MEDI WHEEL FULL BODY HEALTH CHECK UP ABOVE 40 MALE

THYROID PANEL, SERUM

60.0 - 181.0 ng/dL T3 96.96

METHOD: CHEMILUMINESCENCE 11.30 High 4.5 - 10.9 µg/dL T4

METHOD: CHEMILUMINESCENCE

0.550 - 4.780 μIU/mL 1.772 TSH (ULTRASENSITIVE) METHOD: CHEMILUMINESCENCE

Interpretation(s)

Triiodothyronine T3, Thyroxine T4, and Thyroid Stipulating Hormone TSH are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabotism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, white in secondary and tertiary hyporthyroidism, TSH levels are low. owidetlparowidetlparBelow mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3. Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Sr. No.	TSH	Total T4	FT4	Total T3	Possible Conditions
1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3)
	1000				Post Thyroidectomy (4) Post Radio-Iodine treatment
2	High	Normal	Normal	Normal	(1)Subclinical Hypothyroidism (2) Patient with insufficient thyroid
					hormone replacement therapy (3) In cases of Autoimmune/Hashimoto
					thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical
					inflammation, drugs like amphetamines, Iodine containing drug and
					dopamine antagonist e.g. domperidone and other physiological reasons.
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre
					(3) Toxic Nodular Goitre (4) Thyroddits (5) Over treatment of thyroid
					hormone (6) Drug effect e.g. Glusocorticoids, dopamine, T4
					replacement therapy (7) First trimester of Pregnancy
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism
6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent
					treatment for Hyperthyroidism



Dr. Akansha Jain Consultant Pathologist Page 13 Of 14







MC-5333

PATIENT NAME: RAM JI LAL MEENA **REF. DOCTOR:** SELF

CODE NAME & ADDRESS : C000049066 SRL JATPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

ACCESSION NO: 0251WA001949 PATIENT ID : RAMJM280170251 CLIENT PATIENT ID: 012301280017

ABHA NO

AGE/SEX :53 Years Male DRAWN :28/01/2023 09:38:00 RECEIVED: 28/01/2023 10:13:50

REPORTED :28/01/2023 16:06:26

Test Report Status

Final

Results

Biological Reference Interval Units

8	Normal/Low	B	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness
9	Low	7	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies

REF: 1. TIETZ Fundamentals of Clinical chemistry 2. Guidlines of the American Thyroid association during pregnancy and Postpartum, 2011. NOTE: It is advisable to detect Free 13, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.TSH is not affected by variation in thyroid - hinding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

or related Tex Please visit www.srlworld.com for related Test Information for this accession

Dr. Akansha Jain

Consultant Pathologist

Page 14 Of 14









Aakriti Labs

3 Mahatma Gandhi Marg, Gandhi Nagar Mod Tonk Road, Jaipur (Raj.) Ph.: 0141-2710661

www.aakritilabs.com

CIN NO.: U85195RJ2004PTC019563

NAME	MR.	RAM JI L	AL MI	EENA	A	GE	53 YRS		SEX	MALE	
REF BY	MEDI WHEEL					DATE 28/01/20		023	REG NO		
			E	CHC	CARDIOGR	AM RE	PORT				
WINDO	W- POC	R/ADEQ	UATE	/GO	DDVALVE						
MITRAL			NOR			TRICUSPID			NORM	AL	
AORTIC			NOR	MAL		PULN	IONARY		NORM	AL	
2D/M-N	OD						dan .				
IVSD mr	n	10.8			IVSS mm	14.	2	AORT	A mm	23.0	
LVID mn	n	49.7			LVIS mm	31.	1	LA m	m	26.7	
LVPWD	mm	10.1			LVPWS mm	13.	5	EF%		60%	
CHAMB	ERS										
LA	LA			NORMAL		RA	RA			NORMAL	
LV				NORMAL		RV	RV		NC	NORMAL	
PERICARDIUM			NORMAL								
DOPPLE	R STUD	Y MITRA	AL								
PEAK VE	LOCITY	m/s E/A	1	0.92/0.85		1000	AK GRADIAN				
MEAN V	ELOCIT	Y m/s				ME	AN GRADIAI	NT Mm	Hg		
MVA cm	12 (PLA	NITMETE	RY)	_0000000000000000000000000000000000000		MV	MVA cm2 (PHT)				
MR					- Approximation and the second	STATE STATE	Ma View		A		
AORTIC							7970	16			
PEAK VELOCITY m/s			1.61			PEAK GRADIANT MmHg					
MEAN VELOCITY m/s				48	ME	AN GRADIAI	NT Mm	Hg			
AR							OMITS NO.				
TRICUSI	PID					and the same	many Established	WWW.			
PEAK VE	LOCITY	m/s		0.7	9	PE	AK GRADIAN	T Mml	g		

MEAN GRADIANT MmHg

PEAK GRADIANT MmHg

MEAN GRADIANT MmHg

PASP mmHg

RVEDP mmHg

IMPRESSION

PULMONARY

MEAN VELOCITY m/s

PEAK VELOCITY m/s

MEAN VELOCITY m/s

TR

PR

NORMAL LV SYSTOLIC & DIASTOLIC FUNCTION

TRACE

1.42

MILD

- NO RWMA LVEF 60%
- NORMAL RV FUNCTION
- TRACE TR
- MILD PR
- NORMAL CHAMBER DIMENSIONS
- NORMAL VALVULAR ECHO
- INTACT IAS / IVS
- NO THROMBUS, NO VEGETATION, NORMAL PERICARDIUM.
- IVC NORMAL
- CONCLUSION: MILD PR, FAIR LV FUNCTION.

Cardiologist



3 Mahatma Gandhi Marg, Gandhi Nagar Mod Tonk Road, Jaipur (Raj.) Ph.: 0141-2710661

www.aakritilabs.com

CIN NO.: U85195RJ2004PTC019563

Name

: Mr. RAM JI LAL MEENA

Age/Gender: 53 Y/Male

Patient ID : 012301280017

BarcodeNo: 10074576

Referred By: Self

Registration No: 50903

Registered

: 28/Jan/2023 09:38AM

Analysed

: 28/Jan/2023 12:14PM

Reported

: 28/Jan/2023 12:14PM

Panel

: Medi Wheel (ArcoFemi

Healthcare Ltd)

USG: WHOLE ABDOMEN (Male)

LIVER

: Is normal in size, shape and echogenecity.

The IHBR and hepatic radicals are not dilated. No evidence of focal echopoor/echorich lesion seen.

Portal vein diameter and common bile duct appear normal.

GALL

: Is normal in size, shape and echotexture. Walls are smooth and

BLADDER regular with normal thickness. 14.7 mm size calculus seen in GB lumen.

PANCREAS : Is normal in size, shape and echotexture. Pancreatic duct is not dilated.

SPLEEN

:Is normal in size, shape and echogenecity. Spleenic hilum is not dilated.

KIDNEYS: Right Kidney:-Size: 111 x 52 mm, Left Kidney:-Size: 100 x 46 mm.

Bilateral Kidneys are normal in size, shape and echotexture, corticomedullary differentiation is fair and ratio appears normal.

Moderate dilatation of pelvi calyceal system and upper ureter is seen in right kidney.

9.7 mm size calculus seen in right upper ureter.

9.9 mm size calculus seen in lower calyx and 4.4 mm size calculus seen in

mid calyx of left kidney

URINARY: Bladder walls are smooth, regular and normal thickness.

BLADDER: No evidence of mass or stone in bladder lumen.

PROSTATE: Is normal in size, shape and echotexture,

measures: 36 x 26 x 25 mm, wt: 13 gms.

Its capsule is intact and no evidence of focal lesion.

SPECIFIC: No evidence of retroperitoneal mass or free fluid seen in peritoneal cavity.

No evidence of lymphadenopathy or mass lesion in retroperitoneum. Visualized bowel loop appear normal. Great vessels appear normal.

IMPRESSION :- Cholelithiasis

:- Moderate right side hydronephrosis with upper ureteric calculus

:- Left renal calculus

*** End Of Report ***

Page 1 of 1

Dr. Neera Mehta M.B.B.S., D.M.R.D. RMCNO.005807/14853



Aakriti Labs

3 Mahatma Gandhi Marg, Gandhi Nagar Mod Tonk Road, Jaipur (Raj.) Ph.: 0141-2710661

www.aakritilabs.com

CIN NO.: U85195RJ2004PTC019563

Name : MR, RAM JI LAL

Age/Sex

: 53 Yrs/ MALE

Ref.By

: AAKRITI LABS

Date

: 28 January 2023

RADIOGRAPH OF CHEST: PA VIEW

Soft tissue and bony cage are normal.

Both lungs are normal.

Both domes of diaphragm are normal in position and contour.

Hilar shadows are normal.

Mediastinum is central.

Both costo-phrenic angles are clear.

Cardiac size and shape are within normal limits.

IMPRESSION:

NO OBVIOUS ABNORMALITY.

DR. SHUBHAM SINGHAL CONSULTANT RADIOLOGIST