PID No.
 : IND166470
 Register On
 : 11/12/2021 9:22 AM

 SID No.
 : 421093944
 Collection On
 : 11/12/2021 12:40 PM

 Age / Sex
 : 45 Year(s) / Male
 Report On
 : 12/12/2021 8:19 AM

 Type
 : OP
 Printed On
 : 14/12/2021 5:17 PM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HAEMATOLOGY			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood/Electrical Impedance)	15.4	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood/Derived from Impedance)	45.6	%	42 - 52
RBC Count (EDTA Blood/Impedance Variation)	4.82	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (EDTA Blood/Derived from Impedance)	95.0	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood/Derived from Impedance)	32.0	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood/Derived from Impedance)	33.8	g/dL	32 - 36
RDW-CV (EDTA Blood/Derived from Impedance)	13.6	%	11.5 - 16.0
RDW-SD (EDTA Blood/Derived from Impedance)	45.22	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood/Impedance Variation)	6340	cells/cu.mm	4000 - 11000
Neutrophils (EDTA Blood/Impedance Variation & Flow Cytometry)	58.40	%	40 - 75
Lymphocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	31.29	%	20 - 45



VERIFIED BY



 PID No.
 : IND166470
 Register On
 : 11/12/2021 9:22 AM

 SID No.
 : 421093944
 Collection On
 : 11/12/2021 12:40 PM

 Age / Sex
 : 45 Year(s) / Male
 Report On
 : 12/12/2021 8:19 AM

 Type
 : OP
 Printed On
 : 14/12/2021 5:17 PM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Eosinophils (EDTA Blood/Impedance Variation & Flow Cytometry)	4.58	%	01 - 06
Monocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	5.44	%	02 - 10
Basophils (Blood/Impedance Variation & Flow Cytometry)	0.28	%	00 - 02
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	3.70	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	1.98	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.29	10^3 / μΙ	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.34	10^3 / μΙ	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.02	10^3 / μΙ	< 0.2
Platelet Count (EDTA Blood/Impedance Variation)	93.7	10^3 / μl	150 - 450
Remark: Verified on smear. Kindly correlate clinically	Advised follow up c	counts for confirmation	and monitoring.
MPV (EDTA Blood/Derived from Impedance)	12.71	fL	7.9 - 13.7
PCT (EDTA Blood/Automated Blood cell Counter)	0.12	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood/Manual Westergren Method)	10	mm/hr	0 - 15







 PID No.
 : IND166470
 Register On
 : 11/12/2021 9:22 AM

 SID No.
 : 421093944
 Collection On
 : 11/12/2021 12:40 PM

 Age / Sex
 : 45 Year(s) / Male
 Report On
 : 12/12/2021 8:19 AM

 Type
 : OP
 Printed On
 : 14/12/2021 5:17 PM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulfanilic Acid)	1.0	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.4	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.60	mg/dL	0.1 - 1.0
Total Protein (Serum/Biuret)	7.2	g/dL	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.3	g/dL	3.5 - 5.0
Globulin (Serum/Derived)	2.90	g/dL	2.3 - 3.5
A : G Ratio (Serum/Derived)	1.48		1.1 - 2.4
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC without P5P)	35	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC without P5P)	45	U/L	5 - 41
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	75	U/L	53 - 128
GGT(Gamma Glutamyl Transpeptidase) (Serum/Modified IFCC)	55	U/L	< 55



VERIFIED BY



 PID No.
 : IND166470
 Register On
 : 11/12/2021 9:22 AM

 SID No.
 : 421093944
 Collection On
 : 11/12/2021 12:40 PM

 Age / Sex
 : 45 Year(s) / Male
 Report On
 : 12/12/2021 8:19 AM

 Type
 : OP
 Printed On
 : 14/12/2021 5:17 PM

Ref. Dr : MediWheel

Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/Cholesterol oxidase/Peroxidase)	220	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol phosphate oxidase / peroxidase)	226	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the õusualö"circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	40	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	134.8	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	45.2	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	180.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220



VERIFIED BY



 PID No.
 : IND166470
 Register On
 : 11/12/2021 9:22 AM

 SID No.
 : 421093944
 Collection On
 : 11/12/2021 12:40 PM

 Age / Sex
 : 45 Year(s) / Male
 Report On
 : 12/12/2021 8:19 AM

 Type
 : OP
 Printed On
 : 14/12/2021 5:17 PM

Ref. Dr : MediWheel

<u>Investigation</u> <u>Observed</u> <u>Unit</u> <u>Biological</u> <u>Value</u> <u>Reference Interval</u>

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2. It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio 5.5 Optimal: < 3.3
(Serum/Calculated) Low Risk: 3.4 - 4.4
Average Risk: 4.5 - 7.1
Moderate Risk: 7.2 - 11.0

Triglyceride/HDL Cholesterol Ratio 5.7 Optimal: < 2.5 (TG/HDL) Mild to moderate risk: 2.5 - 5.0

(Serum/Calculated) High Risk: > 5.0

LDL/HDL Cholesterol Ratio 3.4 Optimal: 0.5 - 3.0 (Serum/Calculated) Borderline: 3.1 - 6.0 High Risk: > 6.0



VERIFIED BY



High Risk: > 11.0

 PID No.
 : IND166470
 Register On
 : 11/12/2021 9:22 AM

 SID No.
 : 421093944
 Collection On
 : 11/12/2021 12:40 PM

 Age / Sex
 : 45 Year(s) / Male
 Report On
 : 12/12/2021 8:19 AM

 Type
 : OP
 Printed On
 : 14/12/2021 5:17 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/ <i>HPLC</i>)	6.0	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 125.5 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.



VERIFIED BY



 PID No.
 : IND166470
 Register On
 : 11/12/2021 9:22 AM

 SID No.
 : 421093944
 Collection On
 : 11/12/2021 12:40 PM

 Age / Sex
 : 45 Year(s) / Male
 Report On
 : 12/12/2021 8:19 AM

 Type
 : OP
 Printed On
 : 14/12/2021 5:17 PM

Ref. Dr : MediWheel

<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
-	<u>Value</u>		Reference Interval

CLINICAL PATHOLOGY

PHYSICAL EXAMINATION

Colour Pale Yellow

(Urine)

Volume 30 mL

(Urine)

Appearance Clear Clear

(Urine)

CHEMICAL EXAMINATION

pH 6.0 4.6 - 8.0

(Urine)

Specific Gravity 1.020 1.003 - 1.030

(Urine)

Protein Negative Negative

(Urine)

Glucose Negative Negative

(Urine)

Ketones Negative Negative

(Urine)

(Urine)

Leukocytes Negative Negative

(Urine)

Nitrite Negative Negative

Neenu.M M.Sci BIO CHEMISTRY

VERIFIED BY

DR SRIJANA RAO S P
M D Pathology
Consultant Pathologist
KMC No :104646

 PID No.
 : IND166470
 Register On
 : 11/12/2021 9:22 AM


 SID No.
 : 421093944
 Collection On
 : 11/12/2021 12:40 PM

 Age / Sex
 : 45 Year(s) / Male
 Report On
 : 12/12/2021 8:19 AM

 Type
 : OP
 Printed On
 : 14/12/2021 5:17 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Bilirubin (Urine)	Negative		Negative
Blood (Urine)	Negative		Negative
Urobilinogen (Urine)	0.1	mg/dL	0.1 - 1.0
<u>Urine Microscopy Pictures</u>			
Pus Cells (Urine)	2-4	/hpf	0 - 2
Epithelial Cells (Urine)	2-3	/hpf	0 - 2
RBCs (Urine)	Nil	/hpf	0 - 1
Others (Urine)	Nil		Nil
Casts (Urine)	Nil		0 - 1
Crystals (Urine)	Nil		NIL
Bacteria	Nil		



VERIFIED BY

DR SRIJANA RAO S P
M D Pathology
Consultant Fathologist
KMC No : 104646

 PID No.
 : IND166470
 Register On
 : 11/12/2021 9:22 AM

 SID No.
 : 421093944
 Collection On
 : 11/12/2021 12:40 PM

 Age / Sex
 : 45 Year(s) / Male
 Report On
 : 12/12/2021 8:19 AM

 Type
 : OP
 Printed On
 : 14/12/2021 5:17 PM

Ref. Dr : MediWheel

Investigation	Observed Unit	<u>Biological</u>
	Value	Reference Interval

PHYSICAL EXAMINATION

Colour Brown

(Stool)

Consistency Semisolid Semi solid to solid

(Stool)

Mucus Absent Absent

(Stool)

Blood Absent Absent

(Stool)

CHEMICAL EXAMINATION

Reaction Alkaline Alkaline

(Stool)

Reducing Substances Negative Negative

(Stool)

<u>MICROSCOPIC EXAMINATION(STOOL</u> <u>COMPLETE)</u>

COMI LETE)

Ova Not Found Not Found

(Stool)

Cysts Not Found Not Found

(Stool)

Trophozoites Not Found Not found

(Stool)

Pus Cells 2-3 /hpf Nil

(Stool)



VERIFIED BY

DR SRIJANA RAO S P
M D Pathology
Consultant Fathologist
KMC No :104646

 PID No.
 : IND166470
 Register On
 : 11/12/2021 9:22 AM

 SID No.
 : 421093944
 Collection On
 : 11/12/2021 12:40 PM

 Age / Sex
 : 45 Year(s) / Male
 Report On
 : 12/12/2021 8:19 AM

 Type
 : OP
 Printed On
 : 14/12/2021 5:17 PM

Ref. Dr : MediWheel

 Investigation
 Observed Value
 Unit Value
 Biological Reference Interval

 RBCs
 Nil /hpf
 Nil

 (Stool)
 Stool
 Nil

 Others
 Bacteria Present
 Nil

(Stool)



VERIFIED BY

DR SRIJANA RAO S P
M D Pathology
Consultant Pathologist
KMC No :104646

 PID No.
 : IND166470
 Register On
 : 11/12/2021 9:22 AM

 SID No.
 : 421093944
 Collection On
 : 11/12/2021 12:40 PM

 Age / Sex
 : 45 Year(s) / Male
 Report On
 : 12/12/2021 8:19 AM

 Type
 : OP
 Printed On
 : 14/12/2021 5:17 PM

Ref. Dr : MediWheel

InvestigationObservedUnitBiologicalValueReference Interval

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING 'AB' 'Positive'

(EDTA Blood/Agglutination)



VERIFIED BY

DR SRIJANA RAO S P MD Pathology Consultant Pathologist KMC No :104646

 PID No.
 : IND166470
 Register On
 : 11/12/2021 9:22 AM

 SID No.
 : 421093944
 Collection On
 : 11/12/2021 12:40 PM

 Age / Sex
 : 45 Year(s) / Male
 Report On
 : 12/12/2021 8:19 AM

 Type
 : OP
 Printed On
 : 14/12/2021 5:17 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	12		
Glucose Fasting (FBS) (Plasma - F/GOD- POD)	89	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Urine Glucose - Fasting (Urine - F/GOD - POD)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD - POD)	92	mg/dL	70 - 140

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Blood Urea Nitrogen (BUN)	10	mg/dL	7.0 - 21
(Serum/ <i>Urease-GLDH</i>)			
Creatinine	0.9	mg/dL	0.9 - 1.3
(Serum/Modified Jaffe)			

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine

Uric Acid 5.1 mg/dL 3.5 - 7.2 (Serum/Uricase/Peroxidase)







APPROVED BY

-- End of Report --