
## Dr. Goyal'

### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Generato Physical Examination Tele: 0141-2293346, 4049787, 9887049787





नाम

शशांक गुप्ता

Name

SHASHANK GUPTA

कर्मचारी .क्ट्रक E.C. No.

182563

आरीकर्ता प्राधिकारी Issuing Authority



Cup!

धारक के हस्ताक्षर Signature Holder's

Heirer Carl

Dr. Piyesh Goyal M.B.S.S., D.M.R.D. M.B.S.S., D.M.R.D. RMC Reg. No.-0178 )

मिलने पर निम्मलिखित को लौटाएं अंचल कार्यालय, बड़ौदा अंचल-पांचवा तल, बड़ौदा भवन, आर सी दत्त रोड, अलकापुरी, बड़ौदा-३९००७, गुजरात फोन : 91 265 2316501

If found, please return to-Zonal Office, Baroda Zone, 5th Floor, Baroda Bhavan, R C Dutt Road, Alkapuri, Baroda-390007, Gujarat Phone: 91 265 2316501

B+ve


रक समूह / BLOOD Group

पहचान चिन्ह / Identification Marks

DR.GOYAL PATH LAB & IMAGING CENTER, JAIPUR
2036 / MR SHASHANK GUPTA / 29 Yrs / M/ Non Smoker
Heart Rate: 71 bpm / Tested On: 31-Aug-22 11:17:20 / HF 0.05 Hz - LF 35 Hz / Notch 50 Hz / Sn 1.00 Cm/mV / Sw 25 mm/s / Refd By: BOB

ECG





## Dr. Goyal

### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019 Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 31/08/2022 09:38:09

NAME :- Mr. SHASHANK GUPTA

Sex / Age :- Male

Sample Type :- EDTA

29 Yrs 3 Mon 15 Days

Company :- MediWheel

Patient ID: -12221942

Ref. By Dr:- BOB

Lab/Hosp:-

Sample Collected Time31/08/2022 10:08:55

Final Authentication: 31/08/2022 13:52:00

HAEMAT(	DLOGY
Value	Uni
	HAEMATO Value

Test Name	Value	Unit	Biological Ref Interval
BOB PACKAGE BELOW 40MALE			
HAEMOGARAM			
HAEMOGLOBIN (Hb)	15.0	g/dL	13.0 - 17.0
TOTAL LEUCOCYTE COUNT	4.91	/cumm	4.00 - 10.00
DIFFERENTIAL LEUCOCYTE COUNT	1.71	/ Cuillin	4.00 - 10.00
NEUTROPHIL	59.3	%	40.0 - 80.0
LYMPHOCYTE	32.6	%	20.0 - 40.0
EOSINOPHIL	4.9	%	1.0 - 6.0
MONOCYTE	3.0	%	2.0 - 10.0
BASOPHIL	0.2	%	0.0 - 2.0
NEUT#	2.92	10^3/uL	1.50 - 7.00
LYMPH#	1.60	10^3/uL	1.00 - 3.70
EO#	0.24	10^3/uL	0.00 - 0.40
MONO#	0.14	10^3/uL	0.00 - 0.70
BASO#	0.01	10^3/uL	0.00 - 0.10
TOTAL RED BLOOD CELL COUNT (RBC)	5.03	x10^6/uL	4.50 - 5.50
HEMATOCRIT (HCT)	44.80	%	40.00 - 50.00
MEAN CORP VOLUME (MCV)	89.2	fL	83.0 - 101.0
MEAN CORP HB (MCH)	29.8	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	33.4	g/dL	31.5 - 34.5
PLATELET COUNT	269	x10^3/uL	150 - 410
RDW-CV	13.9	%	11.6 - 14.0
MENTZER INDEX	17.73		

The Mentzer index is used to differentiate iron deficiency anemia from beta thalassemia trait. If a CBC indicates microcytic anemia, these are two of the most likely causes, making it necessary to distinguish between them.

If the quotient of the mean corpuscular volume divided by the red blood cell count is less than 13, thalassemia is more likely. If the result is greater than 13, then iron-deficiency anemia is more likely.

**AJAYSINGH** Technologist

Page No: 1 of 11



### Dr. Goya Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 31/08/2022 09:38:09

NAME :- Mr. SHASHANK GUPTA 29 Yrs 3 Mon 15 Days Ref. By Dr:- BOB

Patient ID: -12221942

Lab/Hosp:-

Sex / Age :- Male

Sample Type :- EDTA

Company:- MediWheel

Sample Collected Time31/08/2022 10:08:55

Final Authentication: 31/08/2022 13:52:00

#### **HAEMATOLOGY**

**Test Name** 

Unit Value

**Biological Ref Interval** 

Erythrocyte Sedimentation Rate (ESR)

28 H

mm/hr.

00 - 13

(ESR) Methodology: Measurment of ESR by cells aggregation.

instrument Name : Indepedent form Hematocrit value by Automated Analyzer (Roller-20)

Interpretation

: ESR test is a non-specific indicator ofinflammatory disease and abnormal protein states.

The test in used to detect, follow course of a certain disease (e.g-tuberculosis, rheumatic fever, myocardial infarction

Levels are higher in pregnency due to hyperfibrinogenaemia.

The "3-figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia {\figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia {\figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia {\figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia {\figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia {\figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia {\figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia {\figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia {\figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia {\figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia {\figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia {\figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia {\figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia {\figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia {\figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia {\figure ESR " x>100 value nearly always indicates serious disease such as a serious disease

AJAYSINGH Technologist

Page No: 2 of 11



# Dr. Goya

Path Lab & Imaging Centre

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019



Date

:- 31/08/2022 09:38:09

NAME :- Mr. SHASHANK GUPTA

Sex / Age :- Male

29 Yrs 3 Mon 15 Days

Company :- MediWheel

Patient ID: -12221942

Ref. By Dr:- BOB

Lab/Hosp:-

Sample Type :- EDTA, KOx/Na FLUORIDE-F, KSax/NpaeFCbl@RiexETPReB/R08/2022 10:08:55

Final Authentication: 31/08/2022 15:31:46

#### **HAEMATOLOGY**

Test Name

Value

Unit

**Biological Ref Interval** 

**BLOOD GROUP ABO** 

"B" POSITIVE

BLOOD GROUP ABO Methodology: Haemagglutination reaction Kit Name: Monoclonal agglutinating antibodies (Span clone).

FASTING BLOOD SUGAR (Plasma) Method:- GOD PAP

101.6

mg/dl

75.0 - 115.0

Impaired glucose tolerance (IGT) 111 - 125 mg/dL Diabetes Mellitus (DM) > 126 mg/dL

Instrument Name: Randox Rx Imola Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases

BLOOD SUGAR PP (Plasma)
Method:- GOD PAP

130.5

70.0 - 140.0

Instrument Name: Randox Rx Imola Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases.

URINE SUGAR (FASTING)

Nil

Nil

AJAYSINGH, MUKESHSINGH, SAPNA Technologist

Page No: 3 of 11



Dr. Piyush Goyal (D.M.R.D.) Dr. Chandrika Gupta

### Dr. Goyal Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 31/08/2022 09:38:09

NAME :- Mr. SHASHANK GUPTA

29 Yrs 3 Mon 15 Days

Patient ID :-12221942 Ref. By Dr:- BOB

Lab/Hosp:-

Company :- MediWheel Sample Type :- STOOL

Sex / Age :- Male

Sample Collected Time 31/08/2022 10:08:55

Final Authentication: 31/08/2022 11:46:42

**CLINICAL PATHOLOGY** 

**Test Name** 

Value

Unit

**Biological Ref Interval** 

STOOLANALYSIS

PHYSICAL EXAMINATION

MUCUS

BLOOD

MICROSCOPIC EXAMINATION

RBC's

WBC/HPF

OVA

CYSTS

OTHERS Collected Sample Received

/HPF

/HPF

SAPNA Technologist

Page No: 4 of 11



## Dr. Goyal

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019 Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 31/08/2022 09:38:09

NAME :- Mr. SHASHANK GUPTA

Sex / Age :- Male

29 Yrs 3 Mon 15 Days

Company :- MediWheel

Sample Type :- PLAIN/SERUM

Patient ID: -12221942

Ref. By Dr:- BOB

Lab/Hosp:-

Final Authentication: 31/08/2022 12:01:10

Sample Collected Time 31/08/2022 10:08:55 DIOCHEMICTOV

1 R	RIOCHEMI	STRY	
Test Name	Value	Unit	Biological Ref Interval
LIPID PROFILE			
TOTAL CHOLESTEROL Method:- Enzymatic Endpoint Method	115.22	mg/dl	Desirable <200 Borderline 200-239 High> 240
TRIGLYCERIDES Method:- GPO-PAP	46.06	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500
DIRECT HDL CHOLESTEROL  Method:- Direct clearance Method	30.20	mg/dl	Low < 40 High > 60
DIRECT LDL CHOLESTEROL Method:- Direct clearance Method	77.34	mg/dl	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190
VLDL CHOLESTEROL Method:-Calculated	9.21	mg/dl	0.00 - 80.00
T.CHOLESTEROL/HDL CHOLESTEROL RATIO Method:- Calculated	3.82		0.00 - 4.90
LDL / HDL CHOLESTEROL RATIO Method:- Calculated	2.56		0.00 - 3.50
TOTAL LIPID Method:-CALCULATED	325.05 L	mg/dl	400.00 - 1000.00

TOTAL CHOLESTEROL InstrumentName: Randox Rx Imola Interpretation: Cholesterol measurements are used in the diagnosis and treatments of lipid lipoprotein metabolism

\*FRIGLYCERIDES InstrumentName: Randox Rx Imola Interpretation: Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction

DHRECT HDLCHOLESTERO InstrumentName:Randox Rx Imola Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the meidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.

DHRECT LDL-CHOLESTEROLInstrumentName: Randox Rx Imola Interpretation: Accurate measurement of LDL-Cholesterol is of vital importance in therapies which focus on lipid uction to prevent atherosclerosis or reduce its progress and to avoid plaque rupture

TOTAL LIPID AND VLDL ARE CALCULATED

MUKESHSINGH

Page No: 5 of 11



## Dr. Goyal

### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 31/08/2022 09:38:09

NAME :- Mr. SHASHANK GUPTA

Sex / Age :- Male Company:- MediWheel

Sample Type :- PLAIN/SERUM

29 Yrs 3 Mon 15 Days

Final Authentication: 31/08/2022 12:01:10

#### **BIOCHEMISTRY**

Sample Collected Time 31/08/2022 10:08:55

Patient ID: -12221942

Ref. By Dr:- BOB

Lab/Hosp:-

Test Name	Value	Unit	Biological Ref Interval
LIVER PROFILE WITH GGT			
SERUM BILIRUBIN (TOTAL) Method:- Colorimetric method	0.82	mg/dl	Up to - 1.0 Cord blood <2 mg/dL Premature < 6 days <16mg/dL Full-term < 6 days= 12 mg/dL 1month - <12 months <2 mg/dL 1-19 years <1.5 mg/dL Adult - Up to - 1.2
SERUM BILIRUBIN (DIRECT) Method:-Colorimetric Method	0.31	mg/dL	Ref-(ACCP 2020) Adult - Up to 0.25 Newborn - <0.6 mg/dL >- 1 month - <0.2 mg/dL
SERUM BILIRUBIN (INDIRECT) Method:- Calculated	0.51	mg/dl	0.30-0.70
SGOT Method:- IFCC	20.8	U/L	Men- Up to - 37.0 Women - Up to - 31.0
SGPT Method:-IFCC	27.5	U/L	Men- Up to - 40.0 Women - Up to - 31.0
SERUM ALKALINE PHOSPHATASE Method:- AMP Buffer	62.70	IU/L	30.00 - 120.00
SERUM GAMMA GT Method:- IFCC	31.40	U/L	11.00 - 50.00
SERUM TOTAL PROTEIN Method:- Biuret Reagent	7.07	g/dl	6.40 - 8.30
SERUM ALBUMIN Method:- Bromocresol Green	4.39	g/dl	3.80 - 5.00
SERUM GLOBULIN Method:- CALCULATION	2.68	gm/dl	2.20 - 3.50
A/G RATIO	1.64		1.30 - 2.50

Total Bilirubin Methodology: Colorimetric method InstrumentName: Randox Rx Imola Interpretation An increase in bilirubin concentration in the serum occurs in toxic or infectious diseases of the liver e.g. hepatitis B or obstruction of the bile duct and in rhesus incompatible babies. High levels of unconjugated bilirubin indicate that too much haemoglobin is being destroyed or that the liver is not actively treating emoglobin it is receiving.

AST Aspartate Aminotransferase Methodology: IFCC InstrumentName:Randox Rx Imola Interpretation: Elevated levels of AST can signal myocardial infarction, hepatic disease, muscular dystrophy and organ damage. Although heart muscle is found to have the most activity of the enzyme, significant activity has also been seen in the brain, liver, gastric mucosa, adipose tissue and kidneys of humans.

SLT Alanine Aminotransferase Methodology: IFCCInstrumentName:Randox Rx Imola Interpretation: The enzyme ALT has been found to be in highest concentrations in the liver, with decreasing

concentrations found in kidney, heart, skeletal muscle, pancreas, spleen and lung tissue respectively. Elevated levels of the transaminases can indicate myocardial infarction, hepatic disease, muscular strophy and organ damage.

Alkaline Phosphatase Methodology:AMP Buffer InstrumentName:Randox Rx Imola Interpretation:Measurements of alkaline phosphatase are of use in the diagnosis, treatment and investigation of hepatibilary disease and in bone disease associated with increased osteoblastic activity. Alkaline phosphatase is also used in the diagnosis of parathyroid and intestinal disease.

TOTAL PROTEIN Methodology:Biaret Reagent InstrumentName:Randox Rx Imola Interpretation: Measurements obtained by this method are used in the aenosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

ALBUMIN (ALB) Methodology: Bromocresol Green InstrumentName:Randox Rx Imola Interpretation: Albumin measurements are used in the diagnosis and treatment of numerous diseases involving

MUKESHSINGH

Page No: 6 of 11



## Dr. Goyal's

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019 Tele: 0141-2293346, 4049787, 9887049787 Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com





Date

:- 31/08/2022 09:38:09

Sex / Age :- Male

NAME :- Mr. SHASHANK GUPTA

29 Yrs 3 Mon 15 Days

Company :- MediWheel

Sample Type :- PLAIN/SERUM

Patient ID :-12221942

Ref. By Dr:- BOB

Lab/Hosp:-

Sample Collected Time 31/08/2022 10:08:55

Final Authentication: 31/08/2022 12:01:10

**BIOCHEMISTRY** 

Test Name

Value Ur

**Biological Ref Interval** 

primarily the liver or kidneys. Globulin & A/G ratio is calculated.

Instrument Name Randox Rx Imola Interpretation: Elevations in GGT levels are seen earlier and more pronounced than those with other liver enzymes in cases of obstructive iaundice and metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or post-hepatic bilitary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal)

MUKESHSINGH

Page No: 7 of 11



## Dr. Goyal Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 31/08/2022 09:38:09

NAME :- Mr. SHASHANK GUPTA

Sex / Age :- Male

Sample Type :- PLAIN/SERUM

29 Yrs 3 Mon 15 Days

Company:- MediWheel

Lab/Hosp:-

Patient ID: -12221942

Ref. By Dr:- BOB

Final Authentication: 31/08/2022 12:01:10

Sample Collected Time 31/08/2022 10:08:55 **BIOCHEMISTRY** 

Unit **Biological Ref Interval** Value Test Name Men - 0.6-1.30 0.90 mg/dl SERUM CREATININE Women - 0.5-1.20 Men - 3.4-7.0 Women - 2.4-5.7 5.64 mg/dl SERUM URIC ACID Method:- Enzymatic colorimetric

MUKESHSINGH

Page No: 8 of 11



### Dr. Goya Path Lab & Imaging (

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 31/08/2022 09:38:09

NAME :- Mr. SHASHANK GUPTA

Sex / Age :- Male

29 Yrs 3 Mon 15 Days

Company:- MediWheel

Patient ID: -12221942

Ref. By Dr:- BOB

Lab/Hosp:-

Sample Type :- EDTA

Sample Collected Time31/08/2022 10:08:55

Final Authentication: 31/08/2022 13:52:00

**HAEMATOLOGY** 

Test Name

Value

**Biological Ref Interval** 

**GLYCOSYLATED HEMOGLOBIN (HbA1C)** 

5.6

%

Unit

Non-diabetic: < 5.7 Pre-diabetics: 5.7-6.4 Diabetics: = 6.5 or higher

ADA Target: 7.0

Action suggested: > 6.5

Instrument name: ARKRAY's ADAMS Lite HA 8380V, JAPAN.

Test Interpretation:

HIDAIC is formed by the condensation of glucose with n-terminal valine residue of each beta chain of HbA to form an unstable schiff base. It is the major fraction, constituting approximately 80% of HbA1c. Formation of glycated hemoglobin (GHb) is essentially irreversible and the concentration in the blood depends on both the lifespan of the red blood cells (RBC) (120 days) and the blood glucose concentration. The GHb concentration represents the integrated values for glucose overthe period of 6 to 8 weeks. GHb values are free of day to day glucose fluctuations and are unaffected by recent exercise or food ingestion. Concentration of plasmaglucose concentration in GHb depends on the time interval, with more recent values providing a larger contribution than earlier values. The interpretation of GHbdepends on RBC having a normal life span. Patients with hemolytic disease or other conditions with shortened RBC survival exhibit a substantial reduction of GHb.High GHb have been reported in iron deficiency anemia. GHb has been firmly established as an index of long term blood glucose concentrations and as a measureof the risk for the development of complications in patients with diabetes mellitus. The absolute risk of retinopathy and nephropathy are directly proportional to themean of HbA1C.Genetic variants (e.g. HbS trait, HbC trait), elevated HbF and chemically modified derivatives of hemoglobin can affect the accuracy of HbA1c measurements. The effects vary depending on the specific Hb vatiant or derivative and the specific HbA1c method.

Ref by ADA 2020

MEAN PLASMA GLUCOSE

Method:- Calculated Parameter

114

mg/dL

Non Diabetic < 100 mg/dL Prediabetic 100- 125 mg/dL Diabetic 126 mg/dL or Higher

**AJAYSINGH** Technologist

Page No: 9 of 11



# Dr. Goyal's Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019

Tele: 0141-2293346, 4049787, 9887049787

 $We b site: www.drgoyalspathlab.com \mid E{\text{-}mail: drgoyalpiyush@gmail.com}$ 



Date

:- 31/08/2022 09:38:09

NAME :- Mr. SHASHANK GUPTA Sex / Age :- Male

29 Yrs 3 Mon 15 Days

Company :- MediWheel

Sample Type :- URINE

Patient ID: -12221942

Ref. By Dr:- BOB

Lab/Hosp:-

Sample Collected Time 31/08/2022 10:08:55

Final Authentication: 31/08/2022 11:46:42

#### **CLINICAL PATHOLOGY**

Test Name	Value	Unit	Biological Ref Interval		
Urine Routine					
PHYSICAL EXAMINATION					
COLOUR	PALE YEI	LLOW	PALE YELLOW		
APPEARANCE	Clear		Clear		
CHEMICAL EXAMINATION					
REACTION(PH)	5.5		5.0 - 7.5		
SPECIFIC GRAVITY	1.010		1.010 - 1.030		
PROTEIN	NIL		NIL		
SUGAR	NIL		NIL		
BILIRUBIN	NEGATIV	E	NEGATIVE		
UROBILINOGEN	NORMAL		NORMAL		
KETONES	NEGATIV	E	NEGATIVE		
NITRITE	NEGATIV	E	NEGATIVE		
MICROSCOPY EXAMINATION					
RBC/HPF	NIL	/HPF	NIL		
WBC/HPF	2-3	/HPF	2-3		
EPITHELIAL CELLS	1-2	/HPF	2-3		
CRYSTALS/HPF	ABSENT		ABSENT		ABSENT
CAST/HPF	ABSENT		ABSENT		ABSENT
AMORPHOUS SEDIMENT	ABSENT		ABSENT		
BACTERIAL FLORA	ABSENT		ABSENT		
YEAST CELL	ABSENT	*	ABSENT		
OTHER	ABSENT				

SAPNA Technologist

Page No: 10 of 11



## Dr. Goya Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019

Tele: 0141-2293346, 4049787, 9887049787

 $Website: www.drgoyalspathlab.com \mid E{\text{-}mail: drgoyalpiyush@gmail.com}$ 



Date

:- 31/08/2022 09:38:09

NAME :- Mr. SHASHANK GUPTA

Sex / Age :- Male

29 Yrs 3 Mon 15 Days

Company:- MediWheel Sample Type :- PLAIN/SERUM Patient ID: -12221942

Ref. By Dr:- BOB

Lab/Hosp:-

Sample Collected Time 31/08/2022 10:08:55 TMMIINOASSAV

Final Authentication: 31/08/2022 12:44:56

П	MI	VI	U	N	O.	15	SA	1)

		LOUISI	
Test Name	Value	Unit	Biological Ref Interval
TOTAL THYROID PROFILE			
SERUM TOTAL T3 Method:- Chemiluminescence(Competitive immunoassay)	1.290	ng/ml	0.970 - 1.690
SERUM TOTAL T4 Method:- Chemiluminescence(Competitive immunoassay)	7.840	ug/dl	5.530 - 11.000
SERUM TSH ULTRA Method:- Enhanced Chemiluminescence Immunoassay	2.593	μIU/mL	0.400 - 4.649

interpretation: Triiodothyronine (T3) contributes to the maintenance of the euthyroid state. A decrease in T3 concentration of up to 50% occurs in a variety of clinical situations, including acute and chronic disease. Although T3 results alone cannot be used to diagnose hypothyroidism, T3 concentration may be more sensitive than thyroxine (T4) for hyperthyroidism. Consequently, the total T3 assay can be used in conjunction with other assays to aid in the differential diagnosis of thyroid disease. T3 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, Free T3 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake, or T4 uptake can be used with the total T3 result to calculate the free T3 index and estimate the concentration of free T3.

Interpretation: The measurement of Total T4 aids in the differential diagnosis of thyroid disease. While >99.9% of T4 is protein-bound, primarily to thyroxine-binding globulin (TBG), it is the free fraction that is biologically active. In most patients, the total T4 concentration is a good indicator of thyroid status. T4 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, free T4 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake may be used with the total T4 result to calculate the free T4 index (FT4I) and estimate the concentration of free T4. Some drugs and some nonthyroidal patient conditions are known to alter TT4 concentrations in vivo.

Interpretation: TSH stimulates the production of thyroxine (T4) and triiodothyronine (T3) by the thyroid gland. The diagnosis of overt hypothyroidism by the finding of a low total T4 or free T4 concentration is readily confirmed by a raised TSH concentration. Measurement of low or undetectable TSH concentrations may assist the diagnosis of hyperthyroidism, where concentrations of T4 and T3 are elevated and TSH secretion is suppressed. These have the advantage of discriminating between the concentrations of TSH observed in thyrotoxicosis, compared with the low, but detectable, concentrations that occur in subclinical hyperthyroidism. The performance of this assay has not been established for neonatal specimens. Some drugs and some nonthyroidal patient conditions are known to alter TSH concentrations in vivo.

#### INTERPRETATION

PREGNANCY	REFERENCE RANGE FOR TSH IN uIU/mL (As per American Thyroid
	Association)
1st Trimester	0.10-2.50
2nd Trimester	0.20-3.00
3rd Trimester	0.30-3.00

\*\*\* End of Report \*\*\*

NARENDRAKUMAR Technologist

Page No: 11 of 11





## Dr. Goya Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur Tele: 0141-2293346, 4049787, 9887049787

 $Website: www.drgoyalspathlab.com \mid \textbf{E-mail:drgoyalpiyush@gmail.com}$ 



Final Authentication: 31/08/2022 11:37:22

:- 31/08/2022 09:38:09

NAME :- Mr. SHASHANK GUPTA

Sex / Age :- Male

Company:- MediWheel

Patient ID: -12221942 Ref. By Doctor:-BOB

Lab/Hosp:-

BOB PACKAGE BELOW 40MALE

#### X RAY CHEST PA VIEW:

Both lung fields appears clear.

Bronchovascular markings appear normal.

Trachea is in midline.

Both the hilar shadows are normal.

Both the C.P.angles is clear.

Both the domes of diaphragm are normally placed.

Bony cage and soft tissue shadows are normal.

Heart shadows appear normal.

Impression: - Normal Study

(Please correlate clinically and with relevant further investigations)

\*\*\* End of Report \*\*\*

Page No: 1 of 1

Dr. Poon Gupta
MBBS, MD (Radio Diagnosis)
RMC, No. 32495

Dr. Ashish Choudhary

Dr. Rathod Hetali Amrutlal MBBS, M.D. (Radio-Diagnosis) RMC No. 17163

Transcript by.

BILAL

Dr. Piyush Goyal (D.M.R.D.)

Dr. Piyush Goyal M.B.B.S., D.M.R.D. RMC Reg No. 017996

MBBS, MD (Radio Diagnosis) Fetal Medicine Consultant FMF ID - 260517 | RMC No 22430



# Dr. Goyal's

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New S Tele: 0141-2293346, 4049787, 9887049781

Website : www.drgoyalspathlab.com | E-mail

Patient ID: -12221942 Ref. By Doctor:-BOB

Lab/Hosp:-



Final Authentication: 31/08/2022 11:16:38

:- 31/08/2022 09:38:09 :- Mr. SHASHANK GUPTA

NAME Sex / Age :- Male

MediWheel Company :-

BOB PACKAGE BELOW 40MALE

### **USG WHOLE ABDOMEN**

Liver is of normal size. Echo-texture is normal. No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is of normal size. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape. Echotexture is normal. No focal lesion is seen.

Kidneys are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. No focal lesion is seen. Collecting system does not show any dilatation.

Left kidney showing a calculus of size 4.6mm seen at mid calyx.

Urinary bladder is well distended and showing smooth wall with normal thickness. Urinary bladder does not show any calculus or mass lesion.

Prostate is normal in size(12.9cc) with normal echo-texture and outline.

No enlarged nodes are visualised. No retro-peritoneal lesion is identified

No significant free fluid is seen in peritoneal cavity.

#### IMPRESSION:

\*Small left renal calculus. Needs clinical correlation for further evaluation

\*\*\* End of Report \*\*\*

ANITASHARMA







# Dr. Goyal's Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 31/08/2022 09:38:09

NAME :- Mr. SHASHANK GUPTA

Sex / Age :- Male

Company :- . · MediWheel

Patient ID :-12221942

Ref. By Doctor:-BOB

Lab/Hosp:-

Final Authentication: 31/08/2022 11:16:38

BOB PACKAGE BELOW 40MALE 2D ECHO OPTION TMT (ADULT/CHILD)

2D-ECHOCARDIOGRAPHY M.MODE WITH DOPPLER STUDY:

FAIR TRANSTHORACIC ECHOCARIDIOGRAPHIC WINDOW MORPHOLOGY:

		NORI	ΜΔΙ	TRICUS	PID VALVE		NORWAL		
MITRAL VALV				DIIIMO	PULMONARY VALVE			NORMAL	
AORTIC VALV	E	NOR		FOLIVIC	North Chart		-		
		M.MODE	EXAMITATION:	•	1	IVS-D	06	mm	
AO	22.	mm	LA	28	Mm		_	_	
		mm	LVID	45	Mm	LVSD	28	mm	
IVS-S	14			14	Mm	RV		mm	
LVPW-D	08	mm	LVPW-S	14			_		
		mm	EDV		MI	LVVS		ml	
RVWT		111111		21112.44		ABSENT			
LVEF	75%			RWMA		ABSENT			
				СН	AMBERS:				
			RA			NORMAL			
	MODA	# A I	IKA						

		D.A.	NORMAL
LA ·	NORMAL	RA	NORMAL
11/	NORMAL	RV	NORMAL
PERICARDIUM		NORMAL	

Enter interest						COLOU	R DOPPLER:			
		MITR	AL VAI	VE					Mm/hg	
VELOCITY	0.9	7	m/se	С	PEAK GRADIENT					
A VELOCITY	0.6	4	m/se	c	MEAN	GRADIENT			Mm/hg	
MVA BY PHT	_	T.	Cm2		MVA B	Y PLANIME	TRY		Cm2	
	0.01						ABSENT			
MITRAL REGURGITATION	ON	A O D 7	TIC VA	IVE			7.			
a .						PEAK GRA	DIENT		mm/h	g
PEAK VELOCITY	(	).82		1117 500					mm/hg	
AR VMAX				m/se	c .	MEAN GR	ADIENI	,8		
AORTIC REGURGITATI	ON.					ABSENT				
AONTO NECESSION		TRICU	SPID \	/ALVI	E				mı	m/hg
PEAK VELOCITY		0.36		m	/sec	PEAK GR	ADIENT			
				m	/sec	MEAN G	RADIENT		mı	m/hg
MEAN VELOCITY					.,	_	12.5			
VMax VELOCITY							1			
	TATION					ABSENT				
TRICUSPID REGURGI	IATION	DIII	MONA	RY V	ALVE	it is		.*		1 . 0
		FUL	0.9			M/sec.	PEAK GRADIE	T		Mm/hg
PEAK VELOCITY			- 0.5			1	MEAN GRADIE	NT		Mm/hg
MEAN VALOCITY										
PULMONARY REGUE	RGITATI	ON				1	ABSENT			

**ANITASHARMA** 

Page No: 1 of 2

Dr. Piyush Goyal M.B.B.S., D.M.R.D. RMC Reg No. 017996 Dr. Poonam MBBS, MD (Radio Dage RMC No. 32495

Dr. Ashish Choudhary

MBBS, MD (Radio Diagnosis) Fetal Medicine Consultant FMF ID - 260517 | RMC No 22430 Dr. Rathod Hetali Amrutlal MBBS, M.D. (Radio-Diagnosis) RMC No. 17163

Transcript by.



## Dr. Goyal's Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 31/08/2022 09:38:09

NAME :- Mr. SHASHANK GUPTA

Sex / Age :- Male

Company :- MediWheel

Patient ID: -12221942

Ref. By Doctor:-BOB

Lab/Hosp:-

Final Authentication: 31/08/2022 11:16:38

#### Impression--

- 1. Normal LV size & contractility
- 2. No RWMA, LVEF 75%.
- 3. Normal cardiac chamber.
- 4. Normal valve
- 5. No clot, no vegetation, no pericardial effusion.

(Cardiologist)

\*\*\* End of Report \*\*\*

**ANITASHARMA** 

Page No: 2 of 2