2774588 MR. KRISHNA KANT 7/22/2023 10:26:03 AM

31 Years Male

Rate	61	. Sinus rhythm	normal P axis, V-rate 50- 99
		. ST elev, probable normal early repol pattern	ST elevation, age<55
PR	145		
QRSD	91		
QT	392		
QTc	395		

--AXIS--

P	46
QRS	74
T	21

- NORMAL ECG -

12 Lead; Standard Placement

Unconfirmed Diagnosis



Device: Speed: 25 mm/sec Limb: 10 mm/mV Chest: 10.0 mm/mV F 60~ 0.15-100 Hz 100B CL P?

NAME	**MR KRISHNA KANT**	**STUDY DATE**	**22/07/2023 10:02AM**
AGE / SEX	**31 y / M**	**HOSPITAL NO.**	**MH002774588**
ACCESSION NO.	**NM9084663**	**MODALITY**	**US**
REPORTED ON	**22/07/2023 10:24AM**	**REFERRED BY**	**Health Check MHD**

2D ECHOCARDIOGRAPHY REPORT

Findings:

	End diastole	End systole
IVS thickness (cm)	**1.0**	**1.2**
Left Ventricular Dimension (cm)	**4.6**	**2.7**
Left Ventricular Posterior Wall thickness (cm)	**1.0**	**1.2**

Aortic Root Diameter (cm)	**2.5**
Left Atrial Dimension (cm)	**2.7**
Left Ventricular Ejection Fraction (%)	**55%**

LEFT VENTRICLE	:	Normal in size. No RWMA. LVEF=55%
RIGHT VENTRICLE	:	Normal in size. Normal RV function.
LEFT ATRIUM	:	Normal in size
RIGHT ATRIUM	:	Normal in size
MITRAL VALVE	:	Trace MR
AORTIC VALVE	:	Normal
TRICUSPID VALVE	:	Trace TR (PASP ~21mmHg)
PULMONARY VALVE	:	Normal
MAIN PULMONARY ARTERY & ITS BRANCHES	:	Appears normal.
INTERATRIAL SEPTUM	:	Intact.
INTERVENTRICULAR SEPTUM	:	Intact.
PERICARDIUM	:	No pericardial effusion or thickening

DOPPLER STUDY

VALVE	Peak Velocity (cm/sec)	Maximum P.G. (mmHg)	Mean P. G. (mmHg)	Regurgitation	Stenosis
MITRAL	E=67 A=54	-	-	Trace	Nil
AORTIC	111	-	-	Nil	Nil
TRICUSPID	-	N	N	Trace	Nil
PULMONARY	69	N	N	Nil	Nil

SUMMARY & INTERPRETATION:

- No LV regional wall motion abnormality with LVEF =55%

NAME	MR KRISHNA KANT	STUDY DATE	22/07/2023 10:02AM
AGE / SEX	31 y / M	HOSPITAL NO.	MH002774588
ACCESSION NO.	NM9084663	MODALITY	US
REPORTED ON	22/07/2023 10:24AM	REFERRED BY	Health Check MHD

- Normal sized RA/RV/LV/LA with no chamber hypertrophy. Normal RV function.
- Trace MR
- Trace TR (PASP ~21mmHg)
- Normal mitral inflow pattern.
- IVC normal in size, >50% collapse with inspiration, suggestive of normal RA pressure.
- No clot/ no vegetation/ no pericardial effusion.

Please correlate clinically.

.
.

Dr. Samanjoy Mukherjee MBBS, MD, General Medicine, DM(Cardiology) DMC No.12194
Consultant (Cardiology)

********End Of Report********

Name	:	MR KRISHNA KANT	**Age**	:	31 Yr(s) Sex :Male
Registration No	:	MH002774588	**Lab No**	:	31230700858
Patient Episode	:	H03000055275	**Collection Date :**		22 Jul 2023 09:26
Referred By	:	HEALTH CHECK MHD	**Reporting Date :**		22 Jul 2023 12:38
Receiving Date	:	22 Jul 2023 12:12			

Department of Transfusion Medicine (Blood Bank)

BLOOD GROUPING, RH TYPING & ANTIBODY SCREEN (TYPE & SCREEN)
Specimen-Blood

Blood Group & Rh Typing (Agglutinaton by gel/tube technique)

Blood Group & Rh typing	B Rh(D) Positive

Antibody Screening (Microtyping in gel cards using reagent red cells)

Cell Panel I	NEGATIVE
Cell Panel II	NEGATIVE
Cell Panel III	NEGATIVE
Autocontrol	NEGATIVE

Final Antibody Screen Result	Negative

Technical Note:
ABO grouping and Rh typing is done by cell and serum grouping by microplate / gel technique. Antibody screening is done using a 3 cell panel of reagent red cells coated with Rh, Kell,Duffy,Kidd, Lewis, P,MNS,Lutheran and Xg antigens using gel technique.

----------END OF REPORT----------

Dr Himanshu Lamba

Name	:	MR KRISHNA KANT	**Age**	:	31 Yr(s) Sex :Male
Registration No	:	MH002774588	**Lab No**	:	32230708076
Patient Episode	:	H03000055275	**Collection Date :**		22 Jul 2023 09:25
Referred By	:	HEALTH CHECK MHD	**Reporting Date :**		22 Jul 2023 12:30
Receiving Date	:	22 Jul 2023 09:46			

BIOCHEMISTRY

Specimen: EDTA Whole blood

HbA1c (Glycosylated Hemoglobin)	4.2	%	As per American Diabetes Association(ADA) 2010 [4.0-6.5] HbA1c in % Non diabetic adults : < 5.6 % Prediabetes (At Risk) : 5.7 % - 6.4 % Diabetic Range : > 6.5 %
Methodology			High-Performance Liquid Chromatography(HPLC)

Use :
1.Monitoring compliance and long-term blood glucose level control in patients with diabetes.
2.Index of diabetic control (direct relationship between poor control and development of complications).
3. Predicting development and progression of diabetic microvascular complications.

Limitations :
1. A1C values may be falsely elevated or decreased in those with chronic kidney disease.
2.False elevations may be due in part to analytical interference from carbamylated hemoglobin formed in the presence of elevated concentrations of urea, with some assays.
3. False decreases in measured A1C may occur with hemodialysis and altered red cell turnover, especially in the setting of erythropoietin treatment

References : Rao.L.V.,Michael snyder.L.(2021).Wallach's Interpretation of Diagnostic Tests. 11th Edition. Wolterkluwer. NaderRifai,Andrea Rita Horvath,Carl T.wittwer. (2018)Teitz Text book of Clinical Chemistry and Molecular Diagnostics.First edition,Elsevier,South Asia.

Name	:	MR KRISHNA KANT	**Age**	:	31 Yr(s) Sex :Male
Registration No	:	MH002774588	**Lab No**	:	32230708076
Patient Episode	:	H03000055275	**Collection Date :**		22 Jul 2023 09:25
Referred By	:	HEALTH CHECK MHD	**Reporting Date :**		22 Jul 2023 11:43
Receiving Date	:	22 Jul 2023 09:52			

BIOCHEMISTRY

THYROID PROFILE, Serum — **Specimen Type : Serum**

T3 - Triiodothyronine (ECLIA)	0.82	ng/ml	[0.70-2.04]
T4 - Thyroxine (ECLIA)	5.51	µg/dl	[4.60-12.00]
Thyroid Stimulating Hormone (ECLIA)	3.560	µIU/mL	[0.340-4.250]

Note : TSH levels are subject to circadian variation, reaching peak levels between 2-4.a.m.and at a minimum between 6-10 pm.Factors such as change of seasons hormonal fluctuations,Ca or Fe supplements,high fibre diet,stress and illness affect TSH results.
* References ranges recommended by the American Thyroid Association
1) Thyroid. 2011 Oct;21(10):1081-125.PMID .21787128
2) http://www.thyroid-info.com/articles/tsh-fluctuating.html

Lipid Profile (Serum)

TOTAL CHOLESTEROL (CHOD/POD)	178	mg/dl	[<200] Moderate risk:200-239 High risk:>240
TRIGLYCERIDES (GPO/POD)	**216 #**	**mg/dl**	**[<150]** Borderline high:151-199 High: 200 - 499 Very high:>500
HDL - CHOLESTEROL (Direct)	42	mg/dl	[30-60]
Methodology: Homogenous Enzymatic			
VLDL - Cholesterol (Calculated)	**43 #**	**mg/dl**	**[10-40]**
(CALCULATED)LDL- CHOLESTEROL	93	mg/dl	[<100] Near/Above optimal-100-129 Borderline High:130-159

Name	:	MR KRISHNA KANT	**Age**	:	31 Yr(s) Sex :Male
Registration No	:	MH002774588	**Lab No**	:	32230708076
Patient Episode	:	H03000055275	**Collection Date :**		22 Jul 2023 09:25
Referred By	:	HEALTH CHECK MHD	**Reporting Date :**		22 Jul 2023 11:39
Receiving Date	:	22 Jul 2023 09:52			

BIOCHEMISTRY

T.Chol/HDL.Chol ratio	4.2	High Risk:160-189 <4.0 Optimal 4.0-5.0 Borderline >6 High Risk
LDL.CHOL/HDL.CHOL Ratio	2.2	<3 Optimal 3-4 Borderline >6 High Risk

Note:
Reference ranges based on ATP III Classifications.
Recommended to do fasting Lipid Profile after a minimum of 8 hours of overnight fasting.

Technical Notes:
Lipid profile is a panel of blood tests that serves as initial broad medical screening tool for abnormalities in lipids, the results of these tests can identify certain genetic diseases and determine approximate risks for cardiovascular disease, certain forms of pancreatitis and other diseases.

Test Name	Result	Unit	Biological Ref. Interval
LIVER FUNCTION TEST (Serum)			
BILIRUBIN-TOTAL (mod.J Groff)**	1.10	mg/dl	[0.10-1.20]
BILIRUBIN - DIRECT (mod.J Groff)	**0.31 #**	**mg/dl**	**[<0.2]**
BILIRUBIN - INDIRECT (mod.J Groff)	0.79	mg/dl	[0.20-1.00]
SGOT/ AST (P5P,IFCC)	30.00	IU/L	[5.00-37.00]
SGPT/ ALT (P5P,IFCC)	**58.00 #**	**IU/L**	**[10.00-50.00]**
ALP (p-NPP,kinetic)*	63	IU/L	[45-135]
TOTAL PROTEIN (mod.Biuret)	7.5	g/dl	[6.0-8.2]
SERUM ALBUMIN (BCG-dye)	4.7	g/dl	[3.5-5.0]
SERUM GLOBULIN (Calculated)	2.8	g/dl	[1.8-3.4]
ALB/GLOB (A/G) Ratio	1.68		[1.10-1.80]

Name	:	MR KRISHNA KANT	**Age**	:	31 Yr(s) Sex :Male
Registration No	:	MH002774588	**Lab No**	:	32230708076
Patient Episode	:	H03000055275	**Collection Date :**		22 Jul 2023 09:25
Referred By	:	HEALTH CHECK MHD	**Reporting Date :**		22 Jul 2023 11:39
Receiving Date	:	22 Jul 2023 09:52			

BIOCHEMISTRY

Note:
**NEW BORN:Vary according to age (days), body wt & gestation of baby
*New born: 4 times the adult value

Technical Notes:
Liver function test aids in diagnosis of various pre hepatic, hepatic and post hepatic causes of dysfunction like hemolytic anemia's, viral and alcoholic hepatitis and cholestasis of obstructive causes.

Test Name	Result	Unit	Biological Ref. Interval
KIDNEY PROFILE (Serum)			
BUN (Urease/GLDH)	12.00	mg/dl	[8.00-23.00]
SERUM CREATININE (mod.Jaffe)	1.00	mg/dl	[0.80-1.60]
SERUM URIC ACID (mod.Uricase)	6.9	mg/dl	[3.5-7.2]
SERUM CALCIUM (NM-BAPTA)	9.8	mg/dl	[8.6-10.0]
SERUM PHOSPHORUS (Molybdate, UV)	3.7	mg/dl	[2.3-4.7]
SERUM SODIUM (ISE)	138.0	mmol/l	[134.0-145.0]
SERUM POTASSIUM (ISE)	4.25	mmol/l	[3.50-5.20]
SERUM CHLORIDE (ISE / IMT)	101.6	mmol/L	[95.0-105.0]
eGFR	99.8	ml/min/1.73sq.m	[>60.0]

Technical Note
eGFR which is primarily based on Serum Creatinine is a derivation of CKD-EPI 2009 equation normalized to1.73 sq.m BSA and is not applicable to individuals below 18 years. eGFR tends to be less accurate when Serum Creatinine estimation is indeterminate e.g. patients at extremes of muscle mass, on unusual diets etc. and samples with severe Hemolysis / Icterus / Lipemia.

----------END OF REPORT----------

Dr. Neelam Singal
CONSULTANT BIOCHEMISTRY

Name	:	MR KRISHNA KANT	**Age**	:	31 Yr(s) Sex :Male
Registration No	:	MH002774588	**Lab No**	:	32230708077
Patient Episode	:	H03000055275	**Collection Date :**		22 Jul 2023 09:26
Referred By	:	HEALTH CHECK MHD	**Reporting Date :**		22 Jul 2023 15:43
Receiving Date	:	22 Jul 2023 13:39			

BIOCHEMISTRY

Specimen Type : Plasma

PLASMA GLUCOSE – PP

Plasma GLUCOSE – PP (Hexokinase)	101	mg/dl	[70–140]

Note : Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying, brisk glucose absorption , post exercise

Specimen Type : Serum/Plasma

Plasma GLUCOSE-Fasting (Hexokinase)	90	mg/dl	[70–100]

----------END OF REPORT----------

Dr. Neelam Singal
CONSULTANT BIOCHEMISTRY

Name	:	MR KRISHNA KANT	**Age**	:	31 Yr(s) Sex :Male
Registration No	:	MH002774588	**Lab No**	:	33230704842
Patient Episode	:	H03000055275	**Collection Date :**		22 Jul 2023 09:27
Referred By	:	HEALTH CHECK MHD	**Reporting Date :**		22 Jul 2023 12:51
Receiving Date	:	22 Jul 2023 09:46			

HAEMATOLOGY

ERYTHROCYTE SEDIMENTATION RATE (Automated) Specimen-Whole Blood

ESR	5.0	mm/1sthour	[0.0-10.0]

Interpretation :

Erythrocyte sedimentation rate (ESR) is a non-specific phenomena and is clinically useful in the diagnosis and monitoring of disorders associated with an increased production of acute phase reactants (e.g. pyogenic infections, inflammation and malignancies). The ESR is increased in pregnancy from about the 3rd month and returns to normal by the 4th week postpartum.

ESR is influenced by age, sex, menstrual cycle and drugs (eg. corticosteroids, contraceptives).

It is especially low (0 -1mm) in polycythemia, hypofibrinogenemia or congestive cardiac failure and when there are abnormalities of the red cells such as poikilocytosis, spherocytosis or sickle cells.

Test Name	Result	Unit	Biological Ref. Interval
COMPLETE BLOOD COUNT (EDTA Blood)			
WBC Count (Flow cytometry)	8600	/cu.mm	[4000-10000]
RBC Count (Impedence)	5.02	million/cu.mm	[4.50-5.50]
Haemoglobin (SLS Method)	16.2	g/dL	[13.0-17.0]
Haematocrit (PCV)	45.2	%	[40.0-50.0]
(RBC Pulse Height Detector Method)			
MCV (Calculated)	90.0	fL	[83.0-101.0]
MCH (Calculated)	**32.3 #**	**pg**	**[25.0-32.0]**
MCHC (Calculated)	**35.8 #**	**g/dL**	**[31.5-34.5]**
Platelet Count (Impedence)	296000	/cu.mm	[150000-410000]
RDW-CV (Calculated)	13.9	%	[11.6-14.0]
DIFFERENTIAL COUNT			
Neutrophils (Flowcytometry)	69.9	%	[40.0-80.0]
Lymphocytes (Flowcytometry)	21.6	%	[20.0-40.0]

Name	:	MR KRISHNA KANT	**Age**	:	31 Yr(s) Sex :Male
Registration No	:	MH002774588	**Lab No**	:	33230704842
Patient Episode	:	H03000055275	**Collection Date :**		22 Jul 2023 09:27
Referred By	:	HEALTH CHECK MHD	**Reporting Date :**		22 Jul 2023 11:04
Receiving Date	:	22 Jul 2023 09:46			

HAEMATOLOGY

Monocytes (Flowcytometry)	6.2	%	[2.0-10.0]
Eosinophils (Flowcytometry)	2.1	%	[1.0-6.0]
Basophils (Flowcytometry)	**0.2 #**	**%**	**[1.0-2.0]**
IG	0.00	%	
Neutrophil Absolute(Flouroscence flow cytometry)	6.0	/cu mm	[2.0-7.0]x10³
Lymphocyte Absolute(Flouroscence flow cytometry)	1.9	/cu mm	[1.0-3.0]x10³
Monocyte Absolute(Flouroscence flow cytometry)	0.5	/cu mm	[0.2-1.2]x10³
Eosinophil Absolute(Flouroscence flow cytometry)	0.2	/cu mm	[0.0-0.5]x10³
Basophil Absolute(Flouroscence flow cytometry)	0.0	/cu mm	[0.0-0.1]x10³

Complete Blood Count is used to evaluate wide range of health disorders, including anemia, infection, and leukemia. Abnormal increase or decrease in cell counts as revealed may indicate that an underlying medical condition that calls for further evaluation.

----------END OF REPORT----------

Lakshita Singh

Dr.Lakshita singh

Name	:	MR KRISHNA KANT	**Age**	:	31 Yr(s) Sex :Male
Registration No	:	MH002774588	**Lab No**	:	38230701527
Patient Episode	:	H03000055275	**Collection Date :**		22 Jul 2023 09:26
Referred By	:	HEALTH CHECK MHD	**Reporting Date :**		22 Jul 2023 14:41
Receiving Date	:	22 Jul 2023 11:54			

CLINICAL PATHOLOGY

Test Name	Result	Biological Ref. Interval
ROUTINE URINE ANALYSIS		
MACROSCOPIC DESCRIPTION		
Colour (Visual)	PALE YELLOW	(Pale Yellow - Yellow)
Appearance (Visual)	CLEAR	
CHEMICAL EXAMINATION		
Reaction[pH] (Reflectancephotometry(Indicator Method))	6.0	(5.0-9.0)
Specific Gravity (Reflectancephotometry(Indicator Method))	1.010	(1.003-1.035)
Bilirubin	Negative	NEGATIVE
Protein/Albumin (Reflectance photometry(Indicator Method)/Manual SSA)	Negative	(NEGATIVE-TRACE)
Glucose (Reflectance photometry (GOD-POD/Benedict Method))	NOT DETECTED	(NEGATIVE)
Ketone Bodies (Reflectance photometry(Legal's Test)/Manual Rotheras)	NOT DETECTED	(NEGATIVE)
Urobilinogen Reflactance photometry/Diazonium salt reaction	NORMAL	(NORMAL)
Nitrite Reflactance photometry/Griess test	NEGATIVE	NEGATIVE
Leukocytes Reflactance photometry/Action of Esterase	NIL	NEGATIVE
BLOOD (Reflectance photometry(peroxidase))	NIL	NEGATIVE
MICROSCOPIC EXAMINATION (Manual)	**Method: Light microscopy on centrifuged urine**	
WBC/Pus Cells	1-2 /hpf	(4-6)
Red Blood Cells	NIL	(1-2)
Epithelial Cells	1-2 /hpf	(2-4)
Casts	NIL	(NIL)
Crystals	NIL	(NIL)
Bacteria	NIL	
Yeast cells	NIL	

Interpretation:

Name	:	MR KRISHNA KANT	**Age**	:	31 Yr(s) Sex :Male
Registration No	:	MH002774588	**Lab No**	:	38230701527
Patient Episode	:	H03000055275	**Collection Date :**		22 Jul 2023 09:26
Referred By	:	HEALTH CHECK MHD	**Reporting Date :**		22 Jul 2023 14:41
Receiving Date	:	22 Jul 2023 11:54			

CLINICAL PATHOLOGY

URINALYSIS-Routine urine analysis assists in screening and diagnosis of various metabolic , urological, kidney and liver disorders
Protein: Elevated proteins can be an early sign of kidney disease. Urinary protein excretion can also be temporarily elevated by strenuous exercise, orthostatic proteinuria,dehydration, urina tract infections and acute illness with fever
Glucose: Uncontrolled diabetes mellitus can lead to presence of glucose in urine.
Other causes include pregnancy, hormonal disturbances, liver disease and certain medications.
Ketones: Uncontrolled diabetes mellitus can lead to presence of ketones in urine.
Ketones can also be seen in starvation, frequent vomiting, pregnancy and strenuous exercise.
Blood: Occult blood can occur in urine as intact erythrocytes or haemoglobin, which can occur in various urological, nephrological and bleeding disorders.
Leukocytes: An increase in leukocytes is an indication of inflammation in urinary tract or kidneys Most Common cause is bacterial urinary tract infection.
Nitrite: Many bacteria give positive results when their number is high. Nitrite concentration duri infection increases with length of time the urine specimen is retained in bladder prior to collection.
pH: The kidneys play an important role in maintaining acid base balance of the body. Conditions of the body producing acidosis/alkalosis or ingestion of certain type of food can affect the pH of urine.
Specific gravity: Specific gravity gives an indication of how concentrated the urine is. Increased Specific gravity is seen in conditions like dehydration, glycosuria and proteinuria while decrease Specific gravity is seen in excessive fluid intake, renal failure and diabetes insipidus.
Bilirubin: In certain liver diseases such as biliary obstruction or hepatitis, bilirubin gets excreted in urine.
Urobilinogen: Positive results are seen in liver diseases like hepatitis and cirrhosis and in case of hemolytic anemia.

----------END OF REPORT----------

Dr. Asha Preethi V.S.
CONSULTANT PATHOLOGY

NAME	**MR KRISHNA KANT**	**STUDY DATE**	**22/07/2023 10:52AM**
AGE / SEX	**31 y / M**	**HOSPITAL NO.**	**MH002774588**
ACCESSION NO.	**R5844056**	**MODALITY**	**US**
REPORTED ON	**22/07/2023 11:19AM**	**REFERRED BY**	**Health Check MHD**

USG WHOLE ABDOMEN

Results:

Liver is normal in size (~13.4 cm) and **shows grade I fatty changes**. No focal intra-hepatic lesion is detected. Intra-hepatic biliary radicals are not dilated. Portal vein is normal in calibre.

Gall bladder appears echofree with normal wall thickness.
Common bile duct is normal in calibre.

Pancreas is normal in size and echopattern.

Spleen is normal in size (~10.2 cm) and echopattern.

Both kidneys are normal in position, size (RK ~ 8.6 x 3.0 cm and LK ~ 9.2 x 3.7 cm) and outline. Cortico-medullary differentiation of both kidneys is maintained. Central sinus echoes are compact. No focal lesion or calculus seen. Bilateral pelvicalyceal systems are not dilated.

Urinary bladder is partially distended.

Prostate is normal in size, shape and echopattern (vol ~15.5 cc).

No significant free fluid is detected.

IMPRESSION: Grade I fatty liver.

Kindly correlate clinically

Dr. Pankaj Saini MD, DHA DMC No.15796
CONSULTANT RADIOLOGIST

********End Of Report********

NAME	**MR KRISHNA KANT**	**STUDY DATE**	**22/07/2023 9:19AM**
AGE / SEX	**31 y / M**	**HOSPITAL NO.**	**MH002774588**
ACCESSION NO.	**R5844057**	**MODALITY**	**CR**
REPORTED ON	**22/07/2023 9:46AM**	**REFERRED BY**	**Health Check MHD**

X-RAY CHEST - PA VIEW

Results:

Visualized lung fields appear clear.

Both hilar shadows appear normal.

Cardiothoracic ratio is within normal limits.

Both hemidiaphragmatic outlines appear normal.

Both costophrenic angles are clear.

Kindly correlate clinically.

Aarushi

Dr. Aarushi MBBS, MD, DNB DMC N0.03291
CONSULTANT RADIOLOGIST

********End Of Report********