

SUBURBAN DIAGNOSTICS - MALAD WEST

Patient Name: LIPIKA MISHRA
Patient ID: 2233019837

Date and Time: 26th Nov 22 8:47 AM

Age **41** **11** **9**
years months days

Gender **Female**


Heart Rate **90bpm**

Patient Vitals

BP: NA
Weight: NA
Height: NA
Pulse: NA
Spo2: NA
Resp: NA
Others: _____

Measurements

QRSD: 86ms
QT: 384ms
QTc: 469ms
PR: 124ms
P-R-T: 31° 21° 26°



ECG Within Normal Limits: Sinus Rhythm, Normal Axis. Please correlate clinically.

REPORTED BY

DR SONALI HONRAO
MD (General Medicine)
Physician
2001/04/1882



CID : 2233019837
Name : Mrs Lipika Mishra
Age / Sex : 41 Years/Female
Ref. Dr :
Reg. Location : Malad West Main Centre

Reg. Date : 26-Nov-2022
Reported : 26-Nov-2022/10:43

USG WHOLE ABDOMEN

LIVER:

The liver is normal in size, shape and smooth margins. It shows normal parenchymal echo pattern. The intra hepatic biliary and portal radical appear normal. No evidence of any intra hepatic cystic or solid lesion seen. The main portal vein and CBD appears normal.

GALL BLADDER:

The gall bladder is physiologically distended and appears normal. No evidence of gall stones or mass lesions seen.

PANCREAS:

The pancreas is well visualised and appears normal. No evidence of solid or cystic mass lesion.

KIDNEYS:

Both the kidneys are normal in size, shape and echotexture.

Non obstructive calculus measuring 4.7 mm is seen in upper calyx of right kidney. no calculus is seen in right kidney.

No evidence of any hydronephrosis or mass lesion seen.

Right kidney measures 8.5 x 3.8 cm.

Left kidney 9.9 x 3.8 cm.

SPLEEN:

The spleen is normal in size and echotexture. No evidence of focal lesion is noted.

There is no evidence of any lymphadenopathy or ascites.

URINARY BLADDER:

The urinary bladder is well distended and reveal no intraluminal abnormality.

UTERUS:

The uterus is anteverted and appears normal. It measures 7.8 x 4.2 x 2.8 cm in size. The endometrial thickness is 3.3 mm.

OVARIES:

Both the ovaries are well visualised and appears normal.

There is no evidence of any ovarian or adnexal mass seen.

Right ovary = 3.5 x 1.8 cm. Left ovary = 2.3 x 1.8 cm.



Use a QR Code Scanner
Application To Scan the Code

CID : 2233019837
Name : Mrs Lipika Mishra
Age / Sex : 41 Years/Female
Ref. Dr :
Reg. Location : Malad West Main Centre

Reg. Date : 26-Nov-2022
Reported : 26-Nov-2022/10:43

IMPRESSION:-

***NON OBSTRUCTIVE CALCULUS IN RIGHT KIDNEY.
NO OTHER SIGNIFICANT ABNORMALITY IS SEEN.***

Suggestion: Clinicopathological correlation.

Note : Investigations have their limitations. Solitary radiological investigations never confirm the final diagnosis. They only help in diagnosing the disease in correlation to clinical symptoms and other related tests. USG is known to have inter-observer variations. Further / Follow-up imaging may be needed in some case for confirmation of findings. Patient has been explained in detail about the USG findings including its limitations and need for further imaging if clinically indicated. Please interpret accordingly. All the possible precaution have been taken under covid-19 pandemic.

-----End of Report-----

This report is prepared and physically checked by DR SUNIL before dispatch.

Dr. Sunil Bhutka
DMRD DNB
MMC REG NO:2011051101



Use a QR Code Scanner
Application To Scan the Code

CID : 2233019837
Name : Mrs Lipika Mishra
Age / Sex : 41 Years/Female
Ref. Dr :
Reg. Location : Malad West Main Centre

Reg. Date : 26-Nov-2022
Reported : 26-Nov-2022/10:43



CID : 2233019837
Name : MRS.LIPIKA MISHRA
Age / Gender : 33 Years / Female
Consulting Dr. : -
Reg. Location : Malad West (Main Centre)

Collected : 26-Nov-2022 / 08:16
Reported : 26-Nov-2022 / 11:42

Use a QR Code Scanner
Application To Scan the Code

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE

CBC (Complete Blood Count), Blood

PARAMETER	RESULTS	BIOLOGICAL REF RANGE	METHOD
<u>RBC PARAMETERS</u>			
Haemoglobin	10.7	12.0-15.0 g/dL	Spectrophotometric
RBC	4.93	3.8-4.8 mil/cmm	Elect. Impedance
PCV	33.5	36-46 %	Calculated
MCV	67.9	80-100 fl	Measured
MCH	21.7	27-32 pg	Calculated
MCHC	32.0	31.5-34.5 g/dL	Calculated
RDW	17.7	11.6-14.0 %	Calculated
<u>WBC PARAMETERS</u>			
WBC Total Count	10760	4000-10000 /cmm	Elect. Impedance
<u>WBC DIFFERENTIAL AND ABSOLUTE COUNTS</u>			
Lymphocytes	35.6	20-40 %	
Absolute Lymphocytes	3830.6	1000-3000 /cmm	Calculated
Monocytes	2.9	2-10 %	
Absolute Monocytes	312.0	200-1000 /cmm	Calculated
Neutrophils	49.5	40-80 %	
Absolute Neutrophils	5326.2	2000-7000 /cmm	Calculated
Eosinophils	11.9	1-6 %	
Absolute Eosinophils	1280.4	20-500 /cmm	Calculated
Basophils	0.1	0.1-2 %	
Absolute Basophils	10.8	20-100 /cmm	Calculated
Immature Leukocytes	-		

WBC Differential Count by Absorbance & Impedance method/Microscopy.

PLATELET PARAMETERS

Platelet Count	449000	150000-400000 /cmm	Elect. Impedance
MPV	9.6	6-11 fl	Measured
PDW	16.4	11-18 %	Calculated



Use a QR Code Scanner
Application To Scan the Code

CID : 2233019837
Name : MRS.LIPIKA MISHRA
Age / Gender : 33 Years / Female
Consulting Dr. : -
Reg. Location : Malad West (Main Centre)

Collected : 26-Nov-2022 / 08:16
Reported : 26-Nov-2022 / 11:51

R
E
P
O
R
T

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE

<u>PARAMETER</u>	<u>RESULTS</u>	<u>BIOLOGICAL REF RANGE</u>	<u>METHOD</u>
GLUCOSE (SUGAR) FASTING, Fluoride Plasma	91.0	Non-Diabetic: < 100 mg/dl Impaired Fasting Glucose: 100-125 mg/dl Diabetic: >/= 126 mg/dl	Hexokinase
GLUCOSE (SUGAR) PP, Fluoride Plasma PP/R	105.4	Non-Diabetic: < 140 mg/dl Impaired Glucose Tolerance: 140-199 mg/dl Diabetic: >/= 200 mg/dl	Hexokinase
BILIRUBIN (TOTAL), Serum	0.51	0.1-1.2 mg/dl	Colorimetric
BILIRUBIN (DIRECT), Serum	0.23	0-0.3 mg/dl	Diazo
BILIRUBIN (INDIRECT), Serum	0.28	0.1-1.0 mg/dl	Calculated
TOTAL PROTEINS, Serum	8.8	6.4-8.3 g/dL	Biuret
ALBUMIN, Serum	4.2	3.5-5.2 g/dL	BCG
GLOBULIN, Serum	4.6	2.3-3.5 g/dL	Calculated
A/G RATIO, Serum	0.9	1 - 2	Calculated
SGOT (AST), Serum	28.7	5-32 U/L	NADH (w/o P-5-P)
SGPT (ALT), Serum	27.1	5-33 U/L	NADH (w/o P-5-P)
GAMMA GT, Serum	333.4	3-40 U/L	Enzymatic
ALKALINE PHOSPHATASE, Serum	154.9	35-105 U/L	Colorimetric
BLOOD UREA, Serum	25.1	12.8-42.8 mg/dl	Kinetic
BUN, Serum	11.7	6-20 mg/dl	Calculated
CREATININE, Serum	0.72	0.51-0.95 mg/dl	Enzymatic



Use a QR Code Scanner
Application To Scan the Code

CID : 2233019837
Name : MRS.LIPIKA MISHRA
Age / Gender : 33 Years / Female
Consulting Dr. : -
Reg. Location : Malad West (Main Centre)

Collected : 26-Nov-2022 / 11:47
Reported : 26-Nov-2022 / 15:00

eGFR, Serum	99	>60 ml/min/1.73sqm	Calculated
URIC ACID, Serum	6.4	2.4-5.7 mg/dl	Enzymatic
Urine Sugar (Fasting)	Absent	Absent	
Urine Ketones (Fasting)	Absent	Absent	
Urine Sugar (PP)	Absent	Absent	
Urine Ketones (PP)	Absent	Absent	

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West
*** End Of Report ***



Anupa

Dr. ANUPA DIXIT
M.D.(PATH)
Consultant Pathologist & Lab
Director



CID : 2233019837
Name : MRS.LIPIKA MISHRA
Age / Gender : 33 Years / Female
Consulting Dr. : -
Reg. Location : Malad West (Main Centre)

Collected : 26-Nov-2022 / 08:16
Reported : 26-Nov-2022 / 11:42

Use a QR Code Scanner
Application To Scan the Code

**AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE
GLYCOSYLATED HEMOGLOBIN (HbA1c)**

PARAMETER	RESULTS	BIOLOGICAL REF RANGE	METHOD
Glycosylated Hemoglobin (HbA1c), EDTA WB - CC	6.1	Non-Diabetic Level: < 5.7 % Prediabetic Level: 5.7-6.4 % Diabetic Level: >= 6.5 %	HPLC
Estimated Average Glucose (eAG), EDTA WB - CC	128.4	mg/dl	Calculated

Intended use:

- In patients who are meeting treatment goals, HbA1c test should be performed at least 2 times a year
- In patients whose therapy has changed or who are not meeting glycemic goals, it should be performed quarterly
- For microvascular disease prevention, the HbA1c goal for non pregnant adults in general is Less than 7%.

Clinical Significance:

- HbA1c, Glycosylated hemoglobin or glycated hemoglobin, is hemoglobin with glucose molecule attached to it.
- The HbA1c test evaluates the average amount of glucose in the blood over the last 2 to 3 months by measuring the percentage of glycosylated hemoglobin in the blood.

Test Interpretation:

- The HbA1c test evaluates the average amount of glucose in the blood over the last 2 to 3 months by measuring the percentage of Glycosylated hemoglobin in the blood.
- HbA1c test may be used to screen for and diagnose diabetes or risk of developing diabetes.
- To monitor compliance and long term blood glucose level control in patients with diabetes.
- Index of diabetic control, predicting development and progression of diabetic micro vascular complications.

Factors affecting HbA1c results:

Increased in: High fetal hemoglobin, Chronic renal failure, Iron deficiency anemia, Splenectomy, Increased serum triglycerides, Alcohol ingestion, Lead/opiate poisoning and Salicylate treatment.

Decreased in: Shortened RBC lifespan (Hemolytic anemia, blood loss), following transfusions, pregnancy, ingestion of large amount of Vitamin E or Vitamin C and Hemoglobinopathies

Reflex tests: Blood glucose levels, CGM (Continuous Glucose monitoring)

References: ADA recommendations, AACC, Wallach's interpretation of diagnostic tests 10th edition.

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West

*** End Of Report ***



J. Thakker

Dr.JYOT THAKKER

Pathologist & AVP(Medical Services)



CID : 2233019837
Name : MRS.LIPIKA MISHRA
Age / Gender : 33 Years / Female
Consulting Dr. : -
Reg. Location : Malad West (Main Centre)

Collected : 26-Nov-2022 / 08:16
Reported : 26-Nov-2022 / 13:35

Use a QR Code Scanner
Application To Scan the Code

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE
URINE EXAMINATION REPORT

<u>PARAMETER</u>	<u>RESULTS</u>	<u>BIOLOGICAL REF RANGE</u>	<u>METHOD</u>
<u>PHYSICAL EXAMINATION</u>			
Color	Yellow	Pale Yellow	-
Reaction (pH)	5.0	4.5 - 8.0	Chemical Indicator
Specific Gravity	1.020	1.001-1.030	Chemical Indicator
Transparency	Slight hazy	Clear	-
Volume (ml)	40	-	-
<u>CHEMICAL EXAMINATION</u>			
Proteins	1+	Absent	pH Indicator
Glucose	Absent	Absent	GOD-POD
Ketones	Absent	Absent	Legals Test
Blood	Trace	Absent	Peroxidase
Bilirubin	Absent	Absent	Diazonium Salt
Urobilinogen	Normal	Normal	Diazonium Salt
Nitrite	Absent	Absent	Griess Test
<u>MICROSCOPIC EXAMINATION</u>			
Leukocytes(Pus cells)/hpf	5-6	0-5/hpf	
Red Blood Cells / hpf	Occasional	0-2/hpf	
Epithelial Cells / hpf	15-20		
Casts	Absent	Absent	
Crystals	Absent	Absent	
Amorphous debris	Absent	Absent	
Bacteria / hpf	+(>20/hpf)	Less than 20/hpf	
Others	-		

Interpretation: The concentration values of Chemical analytes corresponding to the grading given in the report are as follows:

- Protein:(1+ ~25 mg/dl, 2+ ~75 mg/dl, 3+ ~ 150 mg/dl, 4+ ~ 500 mg/dl)
- Glucose:(1+ ~ 50 mg/dl, 2+ ~100 mg/dl, 3+ ~300 mg/dl,4+ ~1000 mg/dl)
- Ketone:(1+ ~5 mg/dl, 2+ ~15 mg/dl, 3+ ~ 50 mg/dl, 4+ ~ 150 mg/dl)

Reference: Pack insert



MC-2111



J. Thakker

Dr.JYOT THAKKER
M.D. (PATH), DPB
Pathologist & AVP(Medical Services)



Use a QR Code Scanner
Application To Scan the Code

CID : 2233019837
Name : MRS.LIPIKA MISHRA
Age / Gender : 33 Years / Female
Consulting Dr. : -
Reg. Location : Malad West (Main Centre)

Collected : 26-Nov-2022 / 08:16
Reported : 26-Nov-2022 / 13:35

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West
*** End Of Report ***



Use a QR Code Scanner
Application To Scan the Code

CID : 2233019837
Name : MRS.LIPIKA MISHRA
Age / Gender : 33 Years / Female
Consulting Dr. : -
Reg. Location : Malad West (Main Centre)

Collected : 26-Nov-2022 / 08:16
Reported : 26-Nov-2022 / 11:42

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE
BLOOD GROUPING & Rh TYPING

<u>PARAMETER</u>	<u>RESULTS</u>
ABO GROUP	B
Rh TYPING	POSITIVE

NOTE: Test performed by automated column agglutination technology (CAT) which is more sensitive than conventional methods.

Specimen: EDTA Whole Blood and/or serum

Clinical significance:

ABO system is most important of all blood group in transfusion medicine

Limitations:

- ABO blood group of new born is performed only by cell (forward) grouping because allo antibodies in cord blood are of maternal origin.
- Since A & B antigens are not fully developed at birth, both Anti-A & Anti-B antibodies appear after the first 4 to 6 months of life. As a result, weaker reactions may occur with red cells of newborns than of adults.
- Confirmation of newborn's blood group is indicated when A & B antigen expression and the isoagglutinins are fully developed at 2 to 4 years of age & remains constant throughout life.
- Cord blood is contaminated with Wharton's jelly that causes red cell aggregation leading to false positive result
- The Hh blood group also known as Oh or Bombay blood group is rare blood group type. The term Bombay is used to refer the phenotype that lacks normal expression of ABH antigens because of inheritance of hh genotype.

References:

1. Denise M Harmening, Modern Blood Banking and Transfusion Practices- 6th Edition 2012. F.A. Davis company. Philadelphia
2. AABB technical manual

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West

*** End Of Report ***



M Jain

Dr.MILLU JAIN
M.D.(PATH)
Pathologist



CID : 2233019837
Name : MRS.LIPIKA MISHRA
Age / Gender : 33 Years / Female
Consulting Dr. : -
Reg. Location : Malad West (Main Centre)

Collected : 26-Nov-2022 / 08:16
Reported : 26-Nov-2022 / 14:12

Use a QR Code Scanner
Application To Scan the Code

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE
LIPID PROFILE

PARAMETER	RESULTS	BIOLOGICAL REF RANGE	METHOD
CHOLESTEROL, Serum	215.2	Desirable: <200 mg/dl Borderline High: 200-239mg/dl High: >/=240 mg/dl	CHOD-POD
TRIGLYCERIDES, Serum	259.6	Normal: <150 mg/dl Borderline-high: 150 - 199 mg/dl High: 200 - 499 mg/dl Very high:>/=500 mg/dl	GPO-POD
HDL CHOLESTEROL, Serum	36.4	Desirable: >60 mg/dl Borderline: 40 - 60 mg/dl Low (High risk): <40 mg/dl	Homogeneous enzymatic colorimetric assay
NON HDL CHOLESTEROL, Serum	178.8	Desirable: <130 mg/dl Borderline-high:130 - 159 mg/dl High:160 - 189 mg/dl Very high: >/=190 mg/dl	Calculated
LDL CHOLESTEROL, Serum	143.5	Optimal: <100 mg/dl Near Optimal: 100 - 129 mg/dl Borderline High: 130 - 159 mg/dl High: 160 - 189 mg/dl Very High: >/= 190 mg/dl	Colorimetric
VLDL CHOLESTEROL, Serum	35.3	< /= 30 mg/dl	Calculated
CHOL / HDL CHOL RATIO, Serum	5.9	0-4.5 Ratio	Calculated
LDL CHOL / HDL CHOL RATIO, Serum	3.9	0-3.5 Ratio	Calculated

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West
*** End Of Report ***



J. Thakker

Dr.JYOT THAKKER
M.D. (PATH), DPB
Pathologist & AVP(Medical Services)



Use a QR Code Scanner
Application To Scan the Code

CID : 2233019837
Name : MRS.LIPIKA MISHRA
Age / Gender : 33 Years / Female
Consulting Dr. : -
Reg. Location : Malad West (Main Centre)

Collected : 26-Nov-2022 / 08:16
Reported : 26-Nov-2022 / 11:51

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE
THYROID FUNCTION TESTS

<u>PARAMETER</u>	<u>RESULTS</u>	<u>BIOLOGICAL REF RANGE</u>	<u>METHOD</u>
Free T3, Serum	4.7	3.5-6.5 pmol/L	ECLIA
Free T4, Serum	17.1	11.5-22.7 pmol/L First Trimester:9.0-24.7 Second Trimester:6.4-20.59 Third Trimester:6.4-20.59	ECLIA
sensitiveTSH, Serum	2.08	0.35-5.5 microIU/ml First Trimester:0.1-2.5 Second Trimester:0.2-3.0 Third Trimester:0.3-3.0	ECLIA



Use a QR Code Scanner
Application To Scan the Code

CID : 2233019837
Name : MRS.LIPIKA MISHRA
Age / Gender : 33 Years / Female
Consulting Dr. : -
Reg. Location : Malad West (Main Centre)

Collected : 26-Nov-2022 / 08:16
Reported : 26-Nov-2022 / 11:51

Interpretation:

A thyroid panel is used to evaluate thyroid function and/or help diagnose various thyroid disorders.

Clinical Significance:

- 1)TSH Values between high abnormal upto15 microIU/ml should be correlated clinically or repeat the test with new sample as physiological factors can give falsely high TSH.
- 2)TSH values may be trasiently altered becuase of non thyroidal illness like severe infections,liver disease, renal and heart severe burns, trauma and surgery etc.

TSH	FT4 / T4	FT3 / T3	Interpretation
High	Normal	Normal	Subclinical hypothyroidism, poor compliance with thyroxine, drugs like amiodarone, Recovery phase of non-thyroidal illness, TSH Resistance.
High	Low	Low	Hypothyroidism, Autoimmune thyroiditis, post radio iodine Rx, post thyroidectomy, Anti thyroid drugs, tyrosine kinase inhibitors & amiodarone, amyloid deposits in thyroid, thyroid tumors & congenital hypothyroidism.
Low	High	High	Hyperthyroidism, Graves disease, toxic multinodular goiter, toxic adenoma, excess iodine or thyroxine intake, pregnancy related (hyperemesis gravidarum, hydatiform mole)
Low	Normal	Normal	Subclinical Hyperthyroidism, recent Rx for Hyperthyroidism, drugs like steroids & dopamine), Non thyroidal illness.
Low	Low	Low	Central Hypothyroidism, Non Thyroidal Illness, Recent Rx for Hyperthyroidism.
High	High	High	Interfering anti TPO antibodies, Drug interference: Amiodarone, Heparin, Beta Blockers, steroids & anti epileptics.

Diurnal Variation:TSH follows a diurnal rhythm and is at maximum between 2 am and 4 am , and is at a minimum between 6 pm and 10 pm. The variation is on the order of 50 to 206%. Biological variation:19.7%(with in subject variation)

Reflex Tests:Anti thyroid Antibodies,USG Thyroid ,TSH receptor Antibody. Thyroglobulin, Calcitonin

Limitations:

1. Samples should not be taken from patients receiving therapy with high biotin doses (i.e. >5 mg/day) until atleast 8 hours following the last biotin administration.
2. Patient samples may contain heterophilic antibodies that could react in immunoassays to give falsely elevated or depressed results. this assay is designed to minimize interference from heterophilic antibodies.

Reference:

- 1.O.koulouri et al. / Best Practice and Research clinical Endocrinology and Metabolism 27(2013)
- 2.Interpretation of the thyroid function tests, Dayan et al. THE LANCET . Vol 357
- 3.Tietz ,Text Book of Clinical Chemistry and Molecular Biology -5th Edition
- 4.Biological Variation:From principles to Practice-Callum G Fraser (AACC Press)

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West
*** End Of Report ***



Anupa

Dr. ANUPA DIXIT
M.D.(PATH)
Consultant Pathologist & Lab
Director

CID#	: 2233019837	SID#	: 177805596769
Name	: MRS.LIPIKA MISHRA	Registered	: 26-Nov-2022 / 08:07
Age / Gender	: 33 Years/Female	Collected	: 26-Nov-2022 / 08:07
Consulting Dr.	: -	Reported	: 26-Nov-2022 / 16:07
Reg.Location	: Malad West (Main Centre)	Printed	: 26-Nov-2022 / 16:12

PHYSICAL EXAMINATION REPORT

History and Complaints:

C/O IN HANDS AND JOINT PAIN

EXAMINATION FINDINGS:

Height (cms):	147 CMS	Weight (kg):	76.9 KGS
Temp (0c):	AFEBRILE	Skin:	NAD
Blood Pressure (mm/hg):	120/80	Nails:	NAD
Pulse:	90 /MIN	Lymph Node:	NOT PALPABLE

Systems

Cardiovascular: NAD
Respiratory: NAD
Genitourinary: NAD
GI System: NAD
CNS: NAD

IMPRESSION:

ADVICE:

CHIEF COMPLAINTS:

- | | |
|-----------------------------|----|
| 1) Hypertension: | NO |
| 2) IHD | NO |
| 3) Arrhythmia | NO |
| 4) Diabetes Mellitus | NO |
| 5) Tuberculosis | NO |

CENTRAL PROCESSING LAB: 2nd Floor, Aston, Sundervan Complex, Above Mercedes Showroom, Andheri West - 400053

HEALTHLINE - MUMBAI: 022-6170-0000 | **OTHER CITIES:** 1800-266-4343

For Feedback - customerservice@suburbandiagnosics.com | www.suburbandiagnosics.com

CID# : 2233019837
Name : MRS.LIPIKA MISHRA
Age / Gender : 33 Years/Female
Consulting Dr. : -
Reg.Location : Malad West (Main Centre)


SID# : 177805596769
Registered : 26-Nov-2022 / 08:07
Collected : 26-Nov-2022 / 08:07
Reported : 26-Nov-2022 / 16:07
Printed : 26-Nov-2022 / 16:12


- | | |
|--|--------------|
| 6) Asthama | NO |
| 7) Pulmonary Disease | NO |
| 8) Thyroid/ Endocrine disorders | NO |
| 9) Nervous disorders | NO |
| 10) GI system | NO |
| 11) Genital urinary disorder | NO |
| 12) Rheumatic joint diseases or symptoms | NO |
| 13) Blood disease or disorder | NO |
| 14) Cancer/lump growth/cyst | NO |
| 15) Congenital disease | NO |
| 16) Surgeries | LSCS IN 2014 |
| 17) Musculoskeletal System | NO |

PERSONAL HISTORY:

- | | |
|---------------|--------|
| 1) Alcohol | NO |
| 2) Smoking | NO |
| 3) Diet | MIXED |
| 4) Medication | NOVA D |

*** End Of Report ***




Dr.Sonali Honrao
MD physician
Sr. Manager-Medical Services
(Cardiology)