Name PID No. SID No. Age / Sex Type Ref. Dr	<ul> <li>: Mrs. SARASWATHI M</li> <li>: MED121503956</li> <li>: 122016342</li> <li>: 38 Year(s) / Female</li> <li>: OP</li> <li>: MediWheel</li> </ul>	Collection On : Report On :	26/11/2022 9:01 AM 26/11/2022 9:59 AM 26/11/2022 6:00 PM 05/12/2022 5:35 PM	MEDALL
Investigation		<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> Reference Interval
BLOOD GROUPING AND Rh TYPING (EDTA Blood/Agglutination) INTERPRETATION: Reconfirm the Blood gr		'B' 'Positive'	re blood transfusion	
<u>Complet</u>	e Blood Count With - ESR			
Haemogi (EDTA Bl	lobin ood/Spectrophotometry)	12.5	g/dL	12.5 - 16.0
Packed C	Cell Volume(PCV)/Haematocrit	37.9	%	37 - 47
RBC Co (EDTA Bl	unt ood/Impedance Variation)	4.43	mill/cu.mm	4.2 - 5.4
Mean Co	orpuscular Volume(MCV) ood/Derived from Impedance)	85.7	fL	78 - 100
	orpuscular Haemoglobin(MCH)	28.2	pg	27 - 32
concentr	orpuscular Haemoglobin ation(MCHC) ood/Derived from Impedance)	33.0	g/dL	32 - 36
RDW-C		14.0	%	11.5 - 16.0
RDW-SI (EDTA Bl	<b>)</b> ood/Derived from Impedance)	41.99	fL	39 - 46
	ukocyte Count (TC) ood/Impedance Variation)	5500	cells/cu.mm	4000 - 11000
Neutropl (EDTA Blo <i>Cytometry</i> )	ood/Impedance Variation & Flow	53.2	%	40 - 75
Lympho (EDTA Blo <i>Cytometry</i> )	ood/Impedance Variation & Flow	34.9	%	20 - 45
Eosinopl (EDTA Ble Cytometry)	ood/Impedance Variation & Flow	3.9	%	01 - 06



APPROVED BY

The results pertain to sample tested.

Page 1 of 7

Name	: Mrs. SARASWATHI M		
PID No.	: MED121503956	Register On	: 26/11/2022 9:01 AM
SID No.	: 122016342	Collection On	: 26/11/2022 9:59 AM
Age / Sex	: 38 Year(s) / Female	Report On	: 26/11/2022 6:00 PM
Туре	: OP	Printed On	: 05/12/2022 5:35 PM
Ref. Dr	: MediWheel		

•	20/11/2022 3.01 AM	
:	26/11/2022 9:59 AM	
:	26/11/2022 6:00 PM	MEI
	05/12/2022 5·25 DM	

ALL

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> Reference Interval
Monocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	7.3	%	01 - 10
Basophils (EDTA Blood/Impedance Variation & Flow Cytometry)	0.7	%	00 - 02
INTERPRETATION: Tests done on Automated	l Five Part cell count	er. All abnormal results a	re reviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	2.93	10^3 / µl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	1.92	10^3 / µl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.21	10^3 / µl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.40	10^3 / µl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.04	10^3 / µl	< 0.2
Platelet Count (EDTA Blood/Impedance Variation)	255	10^3 / µl	150 - 450
MPV (EDTA Blood/Derived from Impedance)	9.5	fL	8.0 - 13.3
PCT (EDTA Blood/Automated Blood cell Counter)	0.24	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Blood/Automated - Westergren method)	31	mm/hr	< 20
BUN / Creatinine Ratio	23.5		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	114.7	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125

Diabetic:  $\geq 126$ 



APPROVED BY

The results pertain to sample tested.

Page 2 of 7

Name : Mrs. SARASWATHI M			
PID No. : MED121503956	Register On : 26/	11/2022 9:01 AM	M
SID No. : 122016342	Collection On : 26	/11/2022 9:59 AM	
Age / Sex : 38 Year(s) / Female	Report On : 26	/11/2022 6:00 PM	MEDALL
Type : OP	Printed On : 05	/12/2022 5:35 PM	
Ref. Dr : MediWheel			
Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
<b>INTERPRETATION:</b> Factors such as ty blood glucose level.	pe, quantity and time of food i	ntake, Physical activity, Ps	ychological stress, and drugs can influence
Glucose, Fasting (Urine) (Urine - F/GOD - POD)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD-PAP)	126.8	mg/dL	70 - 140
<b>INTERPRETATION:</b> Factors such as type, quantity and time of Fasting blood glucose level may be higher resistance, Exercise or Stress, Dawn Pher	er than Postprandial glucose, be	cause of physiological sur	ge in Postprandial Insulin secretion, Insulin
Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	15.3	mg/dL	7.0 - 21
Creatinine (Serum/ <i>Modified Jaffe</i> )	0.65	mg/dL	0.6 - 1.1
<b>INTERPRETATION:</b> Elevated Creatini ingestion of cooked meat, consuming Prosuch as cefoxitin, cefazolin, ACE inhibito etc.	otein/ Creatine supplements, Di	abetic Ketoacidosis, proloi	
Uric Acid (Serum/ <i>Enzymatic</i> )	2.8	mg/dL	2.6 - 6.0
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	0.46	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.16	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.30	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/ <i>Modified IFCC)</i>	14.1	U/L	5 - 40
SGPT/ALT (Alanine Aminotransfe (Serum/ <i>Modified IFCC</i> )	prase) 15.7	U/L	5 - 41



APPROVED BY

The results pertain to sample tested.

Page 3 of 7

Name	: Mrs. SARASWATHI M			
PID No.	: MED121503956	Register On	: 26/11/2022 9:01 AM	C
SID No.	: 122016342	<b>Collection On</b>	: 26/11/2022 9:59 AM	
Age / Sex	: 38 Year(s) / Female	Report On	: 26/11/2022 6:00 PM	MED
Туре	: OP	Printed On	: 05/12/2022 5:35 PM	

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	17.1	U/L	< 38
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	69.0	U/L	42 - 98
Total Protein (Serum/Biuret)	7.04	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.22	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	2.82	gm/dL	2.3 - 3.6
A : G RATIO (Serum/Derived)	1.50		1.1 - 2.2
Lipid Profile			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	145.4	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/ <i>GPO-PAP with ATCS</i> )	86.4	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

**INTERPRETATION:** The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual\_circulating level of triglycerides during most part of the day.

HDL Cholesterol	48.9	mg/dL	Optimal(Negative Risk Factor): >=
(Serum/Immunoinhibition)			60

Borderline: 50 - 59 High Risk: < 50



APPROVED BY

ALL

The results pertain to sample tested.

Ref. Dr

: MediWheel

Page 4 of 7

Name	: Mrs. SARASWATHI M		
PID No.	: MED121503956	Register On : 26/11/2022 9:01 AM	$\mathbf{C}$
SID No.	: 122016342	Collection On : 26/11/2022 9:59 AM	
Age / Sex	: 38 Year(s) / Female	Report On : 26/11/2022 6:00 PM	MEDALL
Туре	: OP	Printed On : 05/12/2022 5:35 PM	
Ref. Dr	: MediWheel		

#### Investigation **Observed** <u>Unit</u> **Biological Value** Reference Interval 79.2 Optimal: < 100LDL Cholesterol mg/dL Above Optimal: 100 - 129 (Serum/Calculated) Borderline: 130 - 159 High: 160 - 189 Very High: >=190 < 30 17.3 mg/dL VLDL Cholesterol (Serum/Calculated) 96.5 Optimal: < 130 Non HDL Cholesterol mg/dL Above Optimal: 130 - 159 (Serum/Calculated) Borderline High: 160 - 189 High: 190 - 219 Very High: $\geq 220$

**INTERPRETATION:** 1.Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio (Serum/ <i>Calculated</i> )	3		Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
Triglyceride/HDL Cholesterol Ratio (TG/HDL) (Serum/ <i>Calculated</i> )	1.8		Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
LDL/HDL Cholesterol Ratio (Serum/Calculated)	1.6		Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0
<u>Glycosylated Haemoglobin (HbA1c)</u>			
HbA1C (Whole Blood/ <i>HPLC</i> )	6.0	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

**INTERPRETATION:** If Diabetes - Good control : 6.1 - 7.0 %, Fair control : 7.1 - 8.0 %, Poor control >= 8.1 %



APPROVED BY

The results pertain to sample tested.

Page 5 of 7

Name	: Mrs. SARASWATHI M			
PID No.	: MED121503956	Register On : 2	26/11/2022 9:01 AM	M
SID No.	: 122016342	Collection On :	26/11/2022 9:59 AM	
Age / Sex	: 38 Year(s) / Female	Report On :	26/11/2022 6:00 PM	MEDALL
Туре	: OP	Printed On	05/12/2022 5:35 PM	
Ref. Dr	: MediWheel	-		
<u>Investiga</u>	ation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
Estimate (Whole Blo	d Average Glucose	125.5	mg/dL	
HbA1c pro- control as Condition: hypertrigh Condition: ingestion, <i>THYRO</i>	compared to blood and urinary gluco s that prolong RBC life span like Iron yceridemia,hyperbilirubinemia,Drugs s that shorten RBC survival like acute Pregnancy, End stage Renal disease ID PROFILE / TFT	ose determinations. h deficiency anemia, V s, Alcohol, Lead Poiso e or chronic blood loss can cause falsely low 1	Vitamin B12 & Folate de ning, Asplenia can give s, hemolytic anemia, He HbA1c.	falsely elevated HbA1C values. moglobinopathies, Splenomegaly,Vitamin E
	odothyronine) - Total nemiluminescent Immunometric Assay	1.69	ng/ml	0.7 - 2.04
<b>INTERPH</b> Comment Total T3 v		n like pregnancy, drug	gs, nephrosis etc. In such	n cases, Free T3 is recommended as it is
	oxine) - Total nemiluminescent Immunometric Assay	9.58	µg/dl	4.2 - 12.0
<b>INTERPI</b> Comment Total T4 v		n like pregnancy, druş	gs, nephrosis etc. In such	n cases, Free T4 is recommended as it is
	yroid Stimulating Hormone) memiluminescent Immunometric Assay	1.27	µIU/mL	0.35 - 5.50
INTERPI Reference 1 st trimes 2 nd trimes 3 rd trimes (Indian Th Comment 1.TSH refu 2.TSH Lev be of the c	erence range during pregnancy deper	, reaching peak levels as influence on the me	between 2-4am and at a asured serum TSH conc	



APPROVED BY

The results pertain to sample tested.

Page 6 of 7

Name	: Mrs. SARASWATHI M	
PID No.	: MED121503956	<b>Register On</b> : 26/11/2022 9:01 AM
SID No.	: 122016342	Collection On : 26/11/2022 9:59 AM
Age / Sex	: 38 Year(s) / Female	<b>Report On</b> : 26/11/2022 6:00 PM
Туре	: OP	Printed On : 05/12/2022 5:35 PM

Ref. Dr

: MediWheel



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> Reference Interval
<u>Urine Analysis - Routine</u>			
COLOUR (Urine)	pale yellow		Yellow to Amber
APPEARANCE (Urine)	Clear		Clear
Protein (Urine/Protein error of indicator)	Negative		Negative
Glucose (Urine/GOD - POD)	Negative		Negative
Pus Cells (Urine/Automated <sup>-</sup> Flow cytometry)	Occasional	/hpf	NIL
Epithelial Cells (Urine/Automated <sup>-</sup> Flow cytometry )	Occasional	/hpf	NIL
RBCs (Urine/Automated ~Flow cytometry )	NIL	/hpf	NIL
Casts (Urine/Automated <sup>-</sup> Flow cytometry )	NIL	/hpf	NIL
Crystals (Urine/Automated <sup>-</sup> Flow cytometry )	NIL	/hpf	NIL
Others (Urine)	NIL		

**INTERPRETATION:** Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.



APPROVED BY

-- End of Report --

The results pertain to sample tested.



PRECISION DIAGNOSTICS

- A MEDALL COMPANY

Name	Mrs.SARASWATHI M	ID	MED121503956
Age & Gender	38/FEMALE	Visit Date	26/11/2022
Ref Doctor Name	MediWheel		

## **ULTRA SOUND SCAN**

## WHOLE ABDOMEN

# Liver is normal in size and shows homogenously increased parenchymal echoes with no focal abnormality.

There is no intra or extra hepatic biliary ductal dilatation. Portal vein and IVC are normal.

**Gall bladder** is normal sized and smooth walled. No evidence of calculi. Wall thickness is normal.

Pancreas shows a normal configuration and echotexture. Pancreatic duct is normal.

Spleen is normal in size and echotexture.

**Bilateral kidneys** are normal in size, shape and position. Cortical echoes are normal bilaterally. There is no calculus or calyceal dilatation.

**Right kidney** measures 10.2 x 4.7 cm. **Left kidney** measures 10.1 x 4.8 cm. Ureters are not dilated. No abnormality is seen in the region of the **adrenal glands**. No para aortic lymphadenopathy is seen.

**Urinary bladder** is smooth walled and uniformly transonic. No intravesical mass or calculus.

**Uterus i**s anteverted, and measures 8.4 x 4.6 x 3.8 cm. **Seedling fibroid measuring 1.7 x 1.5 cm is noted in the posterior wall of uterus.** Endometrial thickness is 7 mm.

### **REPORT DISCLAIMER** also have 7. Results of the test are influenced by the various factors such as sensitivity, specificity of the

- This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2.The results reported here in are subject to interpretation by qualified medical professionals only.
- 3.Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital, its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
  - 9.Liability is limited to the extend of amount billed.

procedures of the tests, quality of the samples and drug interactions etc.,

- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11.Disputes, if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



A MEDALL COMPANY

NameMrs.SARASWATHI MIDMED121503956Age & Gender38/FEMALEVisit Date26/11/2022Ref Doctor NameMediWheel

**Right ovary** measures 3.0 x 1.8 cm.

Left ovary measures 2.9 x 1.8 cm.

No significant mass or cyst is seen in the ovaries.

Parametria are free.

Iliac fossae are normal.

There is no free or loculated peritoneal fluid.

## **IMPRESSION:**

- Grade I fatty liver.
- Seedling uterine fibroid.

·Vignut

Dr. D. Vignesh Kumar MBBS, DNB(RD) Consultant Radiologist

### REPORT DISCLAIMER

- 1. This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3.Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its ruthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- procedures of the tests, quality of the samples and drug interactions etc., 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety partial or selective interpretation may lead to false opinion.
- 11.Disputes, if any, with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.