PID No. : MED111017418 : 422019565 SID No.

Age / Sex : 58 Year(s) / Female

Type : OP

Ref. Dr : MediWheel



Collection On : 12/03/2022 11:01 AM

Report On : 13/03/2022 3:58 PM

Printed On : 14/03/2022 6:31 PM



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HAEMATOLOGY			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood/Spectrophotometry)	12.7	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood/Derived from Impedance)	41.0	%	37 - 47
RBC Count (EDTA Blood/Impedance Variation)	4.75	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (EDTA Blood/Derived from Impedance)	86.0	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood/Derived from Impedance)	26.8	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood/Derived from Impedance)	31.1	g/dL	32 - 36
RDW-CV (EDTA Blood/Derived from Impedance)	13.9	%	11.5 - 16.0
RDW-SD (EDTA Blood/Derived from Impedance)	41.84	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood/Impedance Variation)	9200	cells/cu.mm	4000 - 11000
Neutrophils (EDTA Blood/Impedance Variation & Flow Cytometry)	69.4	%	40 - 75
Lymphocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	24.0	%	20 - 45





 PID No.
 : MED111017418
 Register On
 : 12/03/2022 10:16 AM

 SID No.
 : 422019565
 Collection On
 : 12/03/2022 11:01 AM

Printed On

: 14/03/2022 6:31 PM

Type : OP

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Eosinophils (EDTA Blood/Impedance Variation & Flow Cytometry)	1.4	%	01 - 06
Monocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	4.6	%	01 - 10
Basophils (Blood/Impedance Variation & Flow Cytometry)	0.6	%	00 - 02
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	6.38	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	2.21	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.13	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.42	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.06	10^3 / μl	< 0.2
Platelet Count (EDTA Blood/Impedance Variation)	269	10^3 / μl	150 - 450
MPV (EDTA Blood/Derived from Impedance)	11.6	fL	8.0 - 13.3
PCT (EDTA Blood/Automated Blood cell Counter)	0.31	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood/Modified Westergren)	10	mm/hr	< 30





 PID No.
 : MED111017418
 Register On
 : 12/03/2022 10:16 AM

 SID No.
 : 422019565
 Collection On
 : 12/03/2022 11:01 AM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation BIOCHEMISTRY	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulfanilic Acid)	0.4	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.1	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.3	mg/dL	0.1 - 1.0
Total Protein (Serum/Biuret)	7.1	gm/dL	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.3	gm/dL	3.5 - 5.2
Globulin (Serum/ <i>Derived</i>)	2.8	gm/dL	2.3 - 3.6
A: G Ratio (Serum/Derived)	1.5		1.1 - 2.2
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC Kinetic)	22	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC / Kinetic)	22	U/L	5 - 41
Alkaline Phosphatase (SAP) (Serum/IFCC Kinetic)	109	U/L	53 - 141
GGT(Gamma Glutamyl Transpeptidase) (Serum/SZASZ standarised IFCC)	15	U/L	< 38

: 14/03/2022 6:31 PM



VERIFIED BY



 PID No.
 : MED111017418
 Register On
 : 12/03/2022 10:16 AM

 SID No.
 : 422019565
 Collection On
 : 12/03/2022 11:01 AM

Printed On

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/Cholesterol oxidase/Peroxidase)	179	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol phosphate oxidase / peroxidase)	138	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

: 14/03/2022 6:31 PM

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the "usual" circulating level of triglycerides during most part of the day.

r			
HDL Cholesterol (Serum/Immunoinhibition)	43	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/Calculated)	108.4	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	27.6	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	136.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220



VERIFIED BY



 PID No.
 : MED111017418
 Register On
 : 12/03/2022 10:16 AM

 SID No.
 : 422019565
 Collection On
 : 12/03/2022 11:01 AM

Printed On

Type : OP

Ref. Dr : MediWheel



InvestigationObservedUnitBiologicalValueReference Interval

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2. It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

: 14/03/2022 6:31 PM

Total Cholesterol/HDL Cholesterol Ratio 4.2 Optimal: < 3.3 (Serum/Calculated) Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1

Moderate Risk: 7.2 - 11.0 High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio 3.2 Optimal: < 2.5

(TG/HDL) Mild to moderate risk: 2.5 - 5.0

(Serum/Calculated) High Risk: > 5.0

LDL/HDL Cholesterol Ratio 2.5 Optimal: 0.5 - 3.0

(Serum/Calculated)
Borderline: 3.1 - 6.0
High Risk: > 6.0



VERIFIED BY



Printed On

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/ <i>HPLC</i>)	6.8	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

: 14/03/2022 6:31 PM

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 148.46 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.



VERIFIED BY



PID No. : MED111017418 Register On : 12/03/2022 10:16 AM

Printed On

Type : OP

Ref. Dr : MediWheel

MEDALI

<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	Value		Reference Interval

: 14/03/2022 6:31 PM

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.03 ng/mL 0.4 - 1.81

(Serum/CMIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Thyroxine) - Total 5.94 μg/dL 4.2 - 12.0

(Serum/CMIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 3.62 µIU/mL 0.35 - 5.50

(Serum/Chemiluminescent Microparticle

Immunoassay(CMIA))

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values&lt;0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



 PID No.
 : MED111017418
 Register On
 : 12/03/2022 10:16 AM

 SID No.
 : 422019565
 Collection On
 : 12/03/2022 11:01 AM

 Age / Sex
 : 58 Year(s) / Female
 Report On
 : 13/03/2022 3:58 PM



Ref. Dr : MediWheel

<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	Value		Reference Interval

CLINICAL PATHOLOGY

PHYSICAL EXAMINATION

Colour (Urine)	Pale Yellow		
Appearance (Urine)	Clear		Clear
Volume (Urine)	20	mL	
<u>CHEMICAL EXAMINATION(Automated-Urineanalyser)</u>			
pH (Urine/AUTOMATED URINANALYSER)	5.0		4.5 - 8.0
Specific Gravity (Urine)	1.025		1.002 - 1.035
Ketones (Urine)	Negative		Negative
Urobilinogen	0.2		0.2 - 1.0

(Urine/AUTOMATED URINANALYSER)

Nitrite Negative (Urine/AUTOMATED URINANALYSER)

Bilirubin Negative Negative

(Urine/AUTOMATED URINANALYSER)

Negative

Protein Negative (Urine) Negative



(Urine/AUTOMATED URINANALYSER)



Negative

PID No. : MED111017418 Register On : 12/03/2022 10:16 AM : 422019565 SID No. Collection On : 12/03/2022 11:01 AM Age / Sex : 58 Year(s) / Female Report On

Printed On

Type : OP

Ref. Dr : MediWheel

Investigation Glucose (Urine)	Observed Value Negative	<u>Unit</u>	Biological Reference Interval Negative
Leukocytes (Urine)	Negative	leuco/uL	Negative
MICROSCOPY(URINE DEPOSITS)			
Pus Cells (Urine/Flow cytometry)	3-4	/hpf	3-5
Epithelial Cells (Urine)	2-3	/hpf	1-2
RBCs (Urine/Flow cytometry)	Nil	/hpf	2-3
Others (Urine)	Nil		Nil
Casts (Urine/Flow cytometry)	Nil	/hpf	0 - 1
Crystals (Urine)	Nil		NIL

: 13/03/2022 3:58 PM

: 14/03/2022 6:31 PM





Printed On

Type : OP

Ref. Dr : MediWheel



InvestigationObservedUnitBiologicalValueReference Interval

: 14/03/2022 6:31 PM

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING 'B' 'Positive'

(EDTA Blood/Agglutination)





Printed On

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	8.8		6 - 22
Glucose Fasting (FBS) (Plasma - F/GOD - POD)	101	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

: 14/03/2022 6:31 PM

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose Fasting - Urine	Negative		Negative
(Urine - F)			
Glucose Postprandial (PPBS)	105	mg/dL	70 - 140

(Plasma - PP/GOD - POD)

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Glucose Postprandial - Urine (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease-GLDH)	7	mg/dL	7.0 - 21
Creatinine (Serum/Jaffe Kinetic)	0.8	mg/dL	0.6 - 1.1

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine etc.

Uric Acid 4.5 mg/dL 2.6 - 6.0

(Serum/Uricase/Peroxidase)



VERIFIED BY



APPROVED BY

-- End of Report --

Name	MANJULA DEVI N	ID	MED111017418
Age & Gender	58/FeMale		12-03-2022 00:00:00
Ref Doctor Name	MediWheel		



2 D ECHOCARDIOGRAPHIC STUDY

M mode measurement:

AORTA : 2.2cms

LEFT ATRIUM : 2.4cms

AVS :----

LEFT VENTRICLE (DIASTOLE) : 4.8cms

(SYSTOLE) : 3.3cms

VENTRICULAR SEPTUM (DIASTOLE) : 1.0cms

(SYSTOLE) : 1.3cms

POSTERIOR WALL (DIASTOLE) : 1.0cms

(SYSTOLE) : 1.5cms

EDV : 107ml

ESV : 43ml

FRACTIONAL SHORTENING : 32%

EJECTION FRACTION : 60%

EPSS :---

RVID : 1.91cms

DOPPLER MEASUREMENTS:

MITRAL VALVE : E' 0.92 m/s A' 0.79 m/s NO MR

AORTIC VALVE : 1.18 m/s NO AR

TRICUSPID VALVE : E' 1.90 m/s A' - m/s NO TR

PULMONARY VALVE : 0.57 m/s NO PR

Name	MANJULA DEVI N	ID	MED111017418
Age & Gender	58/FeMale		12-03-2022 00:00:00
Ref Doctor Name	MediWheel		



2D ECHOCARDIOGRAPHY FINDINGS:

Left ventricle : Normal size, Normal systolic function.

No regional wall motion abnormalities.

Left Atrium : Normal.

Right Ventricle : Normal.

Right Atrium : Normal.

Mitral valve : Normal, No mitral valve prolapsed.

Aortic valve : Normal, Trileaflet.

Tricuspid valve : Normal.

Pulmonary valve : Normal.

IAS : Intact.

IVS : Intact.

Pericardium : No pericardial effusion.

IMPRESSION:

- > NORMAL SIZED CARDIAC CHAMBERS.
- > NORMAL LV SYSTOLIC FUNCTION. EF: 60%.
- > NO REGIONAL WALL MOTION ABNORMALITIES.
- > NORMAL VALVES.
- > NO CLOTS / PERICARDIAL EFFUSION / VEGETATION.

DR. K.S. SUBRAMANI. MBBS, MD, DM (CARDIOLOGY) FESC, FICC SENIOR CONSULTANT INTERVENTIONAL CARDIOLOGIST Kss/vp $\,$

Name	MANJULA DEVI N	ID	MED111017418
Age & Gender	58/FeMale		12-03-2022 00:00:00
Ref Doctor Name	MediWheel		



- Note:

 * Report to be interpreted by qualified medical professional.

 * To be correlated with other clinical findings.

 * Parameters may be subjected to inter and intra observer variations.

Name	MANJULA DEVI N	ID	MED111017418
Age & Gender	58/FeMale	Visit Date	12-03-2022 00:00:00
Ref Doctor Name	MediWheel		



ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in size and shows diffuse mild fatty changes. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER is not visualized, consistent with h/o cholecystectomy.

CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN shows normal shape, size and echopattern. Spleen measures 9.9cms in long axis and 3.8cms in short axis.

No demonstrable Para -aortic lymphadenopathy.

KIDNEYS move well with respiration and have normal shape, size and echopattern. Cortico- medullary differentiations are well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney	9.7	1.1
Left Kidney	11.2	1.3

URINARY BLADDER shows normal shape and wall thickness. It has clear contents. No evidence of diverticula.

Prevoid: 180cc Postvoid: 20cc

UTERUS is not visualized, consistent with h/o hysterectomy.

OVARIES are not visualized. However no adnexal mass noted.

No evidence of ascites/pleural effusion.

IMPRESSION:

- > POST CHOLECYSTECTOMY & POST HYSTERECTOMY STATUS.
- > MILD FATTY CHANGES IN THE LIVER.

DR. H.K. ANAND

DR. MEERA S

Name	MANJULA DEVI N	ID	MED111017418
Age & Gender	58/FeMale		12-03-2022 00:00:00
Ref Doctor Name	MediWheel		



CONSULTANT RADIOLOGISTS

Name	MANJULA DEVI N	ID	MED111017418
Age & Gender	58/FeMale	Visit Date	12-03-2022 00:00:00
Ref Doctor Name	MediWheel		



X-ray mammogram (mediolateral oblique and craniocaudal views) followed by Sonomammography was performed.

MAMMOGRAPHY OF BOTH BREASTS

Both breasts show symmetrical fibro fatty tissue.

A well-defined lobulated soft tissue lesion is noted in the upper central of the left breast. .

No evidence of cluster micro calcification.

Subcutaneous fat deposition is within normal limits.

Left axillary lymph nodes noted.

SONOMAMMOGRAPHY OF BOTH BREASTS

Both breasts show normal predominantly fatty echopattern.

A well-defined, slightly lobulated hypoechoic lesion measuring about $3.0 \times 1.1 \text{cms}$ is noted in the

12 O' clock position (periareolar) in the left breast.

Color doppler shows no increase in the vascularity of the lesion. Features suggest fibroadenoma.

No evidence of focal solid / cystic areas in the right breast.

No evidence of ductal dilatation.

Few lymphnodes with maintained fatty hilum are noted in both axillae.

IMPRESSION:

> WELL-DEFINED SLIGHTLY LOBULATED HYPOECHOIC LESION IN THE UPPER CENTRAL OF THE LEFT BREAST SUGGESTS FIBROADENOMA.

ASSESSMENT: BI-RADS CATEGORY -2

DR. H.K. ANAND CONSULTANT RADIOLOGISTS DR. MEERA S

Name	MANJULA DEVI N	ID	MED111017418
Age & Gender	58/FeMale	Visit Date	12-03-2022 00:00:00
Ref Doctor Name	MediWheel		



BI-RADS CLASSIFICATION

CATEGORY	RESULT
0	Assessment incomplete. Need additional imaging evaluation
1	Negative. Routine mammogram in 1 year recommended.
2	Benign finding. Routine mammogram in 1 year recommended.
3	Probably benign finding. Short interval follow-up suggested.
4	Suspicious. Biopsy should be considered.
5	Highly suggestive of malignancy. Appropriate action should be taken.

Name	MANJULA DEVI N	ID	MED111017418
Age & Gender	58/FeMale		12-03-2022 00:00:00
Ref Doctor Name	MediWheel		



X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

Impression: Essentially normal study.