# Dr. Goyal's Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787


Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

# **General Physical Examination**

Date of Examination: 25 06 23	
Name: Ashok Kumay	Age: 34 Sex: male.
DOB: 01/07/1988.	
Referred By:	
Photo ID: Adhan ID#: a Hacker	
	Wt: <u>76</u> (Kg)
Chest (Expiration): (cm)	Abdomen Circumference: 86 (cm)
Blood Pressure: 106 14 mm Hg PR: 86 / mi	n RR: 17 / min Temp: Activite
BMI	
Eye Examination:	
Other:	No colocur blindness
Other:	Not Significant
On examination he/she appears physically and mental	ly fit: Yes / No
Signature Of Examine:	Name of Examinee:
Signature Medical Examiner :	Name Medical Examiner 8 D.M.R.O




Dr. Ply th Go, -MB.B.S., D.M.R.D. RMC Reg. No.-017808



Robburno

DR.GOYAL PATH LAB
568 / MR ASHOK KUMAR / 34 Yrs / M/ Non Smoker
Heart Rate : 75 bpm / Tested On : 25-Jun-23 11:53:03 / HF 0.05 Hz - LF 35 Hz / Notch 50 Hz / Sn 1.00 Cm/mV / Sw 25 mm/s / Refd By.: BOB

ECG





Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 25/06/2023 11:10:03 NAME :- Mr. ASHOK KUMAR

Patient ID :-12231467

Ref. By Dr:- BOB

Lab/Hosp:-

Sex / Age :- Male

34 Yrs 11 Mon 25 Days

Company:- MediWheel Sample Type :- EDTA

Sample Collected Time 25/06/2023 11:14:11

Final Authentication: 25/06/2023 12:54:28

# **HAEMATOLOGY**

Test Name	Value	Unit	Biological Ref Interval
HAEMOGARAM			
HAEMOGLOBIN (Hb)	14.2	g/dL	13.0 - 17.0
TOTAL LEUCOCYTE COUNT	5.10	/cumm	4.00 - 10.00
DIFFERENTIAL LEUCOCYTE COUNT			
NEUTROPHIL	64.1	%	40.0 - 80.0
LYMPHOCYTE	31.8	%	20.0 - 40.0
EOSINOPHIL	1.0	%	1.0 - 6.0
MONOCYTE	3.0	%	2.0 - 10.0
BASOPHIL	0.1	%	0.0 - 2.0
NEUT#	3.27	10^3/uL	1.50 - 7.00
LYMPH#	1.62	10^3/uL	1.00 - 3.70
EO#	0.05	10^3/uL	0.00 - 0.40
MONO#	0.15	10^3/uL	0.00 - 0.70
BASO#	0.01	10^3/uL	0.00 - 0.10
TOTAL RED BLOOD CELL COUNT (RBC)	5.15	x10^6/uL	4.50 - 5.50
HEMATOCRIT (HCT)	41.90	%	40.00 - 50.00
MEAN CORP VOLUME (MCV)	81.5 L	fL	83.0 - 101.0
MEAN CORP HB (MCH)	27.6	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	33.9	g/dL	31.5 - 34.5
PLATELET COUNT	197	x10^3/uL	150 - 410
RDW-CV	12.7	%	11.6 - 14.0
MENTZER INDEX	15.83		

The Mentzer index is used to differentiate iron deficiency anemia from beta thalassemia trait. If a CBC indicates microcytic anemia, these are two of the most likely causes, making it necessary to distinguish between them.

If the quotient of the mean corpuscular volume divided by the red blood cell count is less than 13, thalassemia is more likely. If the result is greater than 13, then iron-deficiency anemia is more likely.

**AJAYSINGH Technologist** 

Page No: 2 of 12



Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 25/06/2023 11:10:03

NAME :- Mr. ASHOK KUMAR

( KUMAR Ref. By Dr:- BOB

Sex / Age :- Male 34 Yrs 11 Mon 25 Days Lab/Hosp :-

Company :- MediWheel

Sample Type :- EDTA

Sample Collected Time 25/06/2023 11:14:11

Final Authentication: 25/06/2023 12:54:28

**HAEMATOLOGY** 

Test Name	Value	Unit	Biological Ref Interval
BOB PACKAGE BELOW 40MALE GLYCOSYLATED HEMOGLOBIN (HbA1C) Method:- HPLC	5.9	%	Non-diabetic: < 5.7 Pre-diabetics: 5.7-6.4 Diabetics: = 6.5 or higher ADA Target: 7.0 Action suggested: > 6.5

Patient ID: -12231467

Instrument name: ARKRAY's ADAMS Lite HA 8380V, JAPAN.

### Test Interpretation:

HbA1C is formed by the condensation of glucose with n-terminal valine residue of each beta chain of HbA to form an unstable schiff base. It is the major fraction, constituting approximately 80% of HbA1c. Formation of glycated hemoglobin (GHb) is essentially irreversible and the concentration in the blood depends on both the lifespan of the red blood cells (RBC) (120 days) and the blood glucose concentration. The GHb concentration represents the integrated values for glucose overthe period of 6 to 8 weeks. GHb values are free of day to day glucose fluctuations and are unaffected by recent exercise or food ingestion. Concentration of plasmaglucose concentration in GHb depends on the time interval, with more recent values providing a larger contribution than earlier values. The interpretation of GHbdepends on RBC having a normal life span. Patients with hemolytic disease or other conditions with shortened RBC survival exhibit a substantial reduction of GHb. High GHb have been reported in iron deficiency anemia. GHb has been firmly established as an index of long term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. The absolute risk of retinopathy and nephropathy are directly proportional to themean of HbA1c. Genetic variants (e.g. HbS trait, HbC trait), elevated HbF and chemically modified derivatives of hemoglobin can affect the accuracy of HbA1c measurements. The effects vary depending on the specific Hb vatiant or derivative and the specific HbA1c method.

Ref by ADA 2020

MEAN PLASMA GLUCOSE
Method:- Calculated Parameter

123

mg/dL

Non Diabetic < 100 mg/dL Prediabetic 100- 125 mg/dL Diabetic 126 mg/dL or Higher

AJAYSINGH Technologist

Page No: 1 of 12





Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 25/06/2023 11:10:03 NAME :- Mr. ASHOK KUMAR

**Erythrocyte Sedimentation Rate (ESR)** 

Patient ID: -12231467

Ref. By Dr:- BOB

34 Yrs 11 Mon 25 Days Sex / Age :- Male

Lab/Hosp:-

Company:-MediWheel Sample Type :- EDTA

Sample Collected Time 25/06/2023 11:14:11

Final Authentication: 25/06/2023 12:54:28

00 - 13

### HAEMATOLOGY

Test Name	Value Unit	Biological Ref Interval

mm/hr.

(ESR) Methodology: Measurment of ESR by cells aggregation.

Instrument Name : Indepedent form Hematocrit value by Automated Analyzer (Roller-20)

: ESR test is a non-specific indicator of inflammatory disease and abnormal protein states.

The test in used to detect, follow course of a certain disease (e.g-tuberculosis, rheumatic fever, myocardial infarction

09

Levels are higher in pregnency due to hyperfibrinogenaemia.

The "3-figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia (CBC) hetthedology: dTJc SlC Fluorescent Flow cytometry, HB SLS method, TRBC, PCV, PLT Hydrodynamically focused Impedance. and MCH, MCV, MCHC, MENTZER INDEX are calculated. InstrumentName: Sysmex 6 part fully automatic analyzer XN-L, Japan

**AJAYSINGH Technologist** 

Page No: 3 of 12





Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

:- 25/06/2023 11:10:03 Date NAME :- Mr. ASHOK KUMAR

Patient ID: -12231467 Ref. By Dr:- BOB

Sex / Age :- Male 34 Yrs 11 Mon 25 Days

Company:- MediWheel Sample Type :- PLAIN/SERUM

Sample Collected Time 25/06/2023 11:14:11 Final Authentication: 25/06/2023 12:23:26

# **BIOCHEMISTRY**

Lab/Hosp:-

Test Name	Value	Unit	Biological Ref Interval
LIPID PROFILE			*
TOTAL CHOLESTEROL Method:- Enzymatic Endpoint Method	155.52	mg/dl	Desirable <200 Borderline 200-239 High> 240
TRIGLYCERIDES Method:- GPO-PAP	107.10	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500
DIRECT HDL CHOLESTEROL Method:- Direct clearance Method	23.09	mg/dl	Low < 40 High > 60
DIRECT LDL CHOLESTEROL Method:- Direct clearance Method	114.58	mg/dl	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190
VLDL CHOLESTEROL Method:- Calculated	21.42	mg/dl	0.00 - 80.00
T.CHOLESTEROL/HDL CHOLESTEROL RATIO Method:- Calculated	<b>6.74</b> H		0.00 - 4.90
LDL / HDL CHOLESTEROL RATIO Method:- Calculated	<b>4.96</b> H		0.00 - 3.50
TOTAL LIPID Method:- CALCULATED	477.57	mg/dl	400.00 - 1000.00

TOTAL CHOLESTEROL InstrumentName: Randox Rx Imola Interpretation: Cholesterol measurements are used in the diagnosis and treatments of lipid lipoprotein metabolism

TRIGLYCERIDES InstrumentName: Randox Rx Imola Interpretation: Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction

DIRECT HDLCHOLESTERO InstrumentName:Randox Rx Imola Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.

DIRECT LDL-CHOLESTEROL Instrument Name: Randox Rx Imola Interpretation: Accurate measurement of LDL-Cholesterol is of vital importance in therapies which focus on lipid reduction to prevent atherosclerosis or reduce its progress and to avoid plaque rupture

TOTAL LIPID AND VLDL ARE CALCULATED

MUKESHSINGH

Page No: 4 of 12



Dr. Chandrika Gupta MBBS.MD (Path) RMC NO. 21021/008037



Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 25/06/2023 11:10:03

NAME :- Mr. ASHOK KUMAR

Patient ID :-12231467 Ref. By Dr:- BOB

Lab/Hosp :-

Sex / Age :- Male

Sample Type :- PLAIN/SERUM

34 Yrs 11 Mon 25 Days

Company :- MediWheel

Sample Collected Time 25/06/2023 11:14:11

Final Authentication: 25/06/2023 12:23:26

### **BIOCHEMISTRY**

	DIOCHEDITA	VIII.	
Test Name	Value	Unit	Biological Ref Interva
LIVER PROFILE WITH GGT SERUM BILIRUBIN (TOTAL) Method:- Colorimetric method	0.92	mg/dl	Up to - 1.0 Cord blood <2 Premature < 6 days <16 Full-term < 6 days= 12 1month - <12 months <2 1-19 years <1.5 Adult - Up to - 1.2
SERUM BILIRUBIN (DIRECT) Method:- Colorimetric Method	0.32	mg/dL	Ref-(ACCP 2020) Adult - Up to 0.25 Newborn - <0.6 >- 1 month - <0.2
SERUM BILIRUBIN (INDIRECT) Method:- Calculated	0.60	mg/dl	0.30-0.70
SGOT Method:- IFCC	<b>40.5</b> H	U/L	Men- Up to - 37.0 Women - Up to - 31.0
SGPT Method:- IFCC	<b>59.6</b> H	U/L	Men- Up to - 40.0 Women - Up to - 31.0
SERUM ALKALINE PHOSPHATASE Method:- AMP Buffer	94.80	IU/L	30.00 - 120.00
SERUM GAMMA GT Method:- IFCC	<b>62.70</b> H	U/L	11.00 - 50.00
SERUM TOTAL PROTEIN Method:- Biuret Reagent	7.27	g/dl	6.40 - 8.30
SERUM ALBUMIN Method:- Bromocresol Green	4.58	g/dl	3.80 - 5.00
SERUM GLOBULIN Method:- CALCULATION	2.69	gm/dl	2.20 - 3.50
A/G RATIO	1.70		1.30 - 2.50

Total BilirubinMethodology:Colorimetric method InstrumentName:Randox Rx Imola Interpretation An increase in bilirubin concentration in the serum occurs in toxic or infectious diseases of the liver e.g. hepatitis B or obstruction of the bile duct and in rhesus incompatible babies. High levels of unconjugated bilirubin indicate that too much haemoglobin is being destroyed or that the liver is not actively treating the haemoglobin it is receiving.

AST Aspartate Aminotransferase Methodology: IFCC InstrumentName:Randox Rx Imola Interpretation: Elevated levels of AST can signal myocardial infarction, hepatic disease, muscular dystrophy and organ damage. Although heart muscle is found to have the most activity of the enzyme, significant activity has also been seen in the brain, liver, gastric mucosa, adipose tissue and kidneys of humans.

ALT Alanine Aminotransferase Methodology: IFCCInstrumentName:Randox Rx Imola Interpretation: The enzyme ALT has been found to be in highest concentrations in the liver, with decreasing concentrations found in kidney, heart, skeletal muscle, pancreas, spleen and lung tissue respectively. Elevated levels of the transaminases can indicate myocardial infarction, hepatic disease, muscular dystrophy and organ damage.

Alkaline Phosphatase Methodology: AMP Buffer InstrumentName:Randox Rx Imola Interpretation: Measurements of alkaline phosphatase are of use in the diagnosis, treatment and investigation of hepatobilary disease and in bone disease associated with increased osteoblastic activity. Alkaline phosphatase is also used in the diagnosis of parathyroid and intestinal disease.

hepatobilary disease and in bone disease associated with increased osteoblastic activity. Alkaline phosphatase is also used in the diagnosis of parathyroid and intestinal disease.

TOTAL PROTEIN Methodology:Biuret Reagent InstrumentName:Randox Rx Imola Interpretation: Measurements obtained by this method are used in the

diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

ALBUMIN (ALB) Methodology: Bromocresol Green InstrumentName: Randox Rx Imola Interpretation: Albumin measurements are used in the diagnosis and treatment of numerous diseases involving primarily the liver or kidneys. Globulin & A/G ratio is calculated.

Instrument Name Randox Rx Imola Interpretation: Elevations in GGT levels are seen earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or post-hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal)

MUKESHSINGH

Page No: 5 of 12





Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 25/06/2023 11:10:03

NAME :- Mr. ASHOK KUMAR

KUMAR Ref. By Dr:- BOB

Sex / Age :- Male 34 Yrs 11 Mon 25 Days Lab/Hosp :-

Company :- MediWheel

Sample Type :- PLAIN/SERUM

Sample Collected Time 25/06/2023 11:14:11 Final Authentication: 25/06/2023 13:39:15

Patient ID: -12231467

### **IMMUNOASSAY**

Test Name	Value	Unit	Biological Ref Interval
TOTAL THYROID PROFILE			
SERUM TOTAL T3 Method:- Chemiluminescence(Competitive immunoassay)	1.158	ng/ml	0.970 - 1.690
SERUM TOTAL T4 Method:- Chemiluminescence(Competitive immunoassay)	7.364	ug/dl	5.530 - 11.000
SERUM TSH ULTRA Method:- Enhanced Chemiluminescence Immunoassay	4.470	μIU/mL	0.350 - 5.500

Interpretation: Triiodothyronine (T3) contributes to the maintenance of the euthyroid state. A decrease in T3 concentration of up to 50% occurs in a variety of clinical situations, including acute and chronic disease. Although T3 results alone cannot be used to diagnose hypothyroidism, T3 concentration may be more sensitive than thyroxine (T4) for hyperthyroidism. Consequently, the total T3 assay can be used in conjunction with other assays to aid in the differential diagnosis of thyroid disease. T3 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, Free T3 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake, or T4 uptake can be used with the total T3 result to calculate the free T3 index and estimate the concentration of free T3.

Interpretation: The measurement of Total T4 aids in the differential diagnosis of thyroid disease. While >99.9% of T4 is protein-bound, primarily to thyroxine-binding globulin (TBG), it is the free fraction that is biologically active. In most patients, the total T4 concentration is a good indicator of thyroid status. T4 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, free T4 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake may be used with the total T4 result to calculate the free T4 index (FT4I) and estimate the concentration of free T4. Some drugs and some nonthyroidal patient conditions are known to alter TT4 concentrations in vivo.

Interpretation: TSH stimulates the production of thyroxine (T4) and triiodothyronine (T3) by the thyroid gland. The diagnosis of overt hypothyroidism by the finding of a low total T4 or free T4 concentration is readily confirmed by a raised TSH concentration. Measurement of low or undetectable TSH concentrations may assist the diagnosis of hyperthyroidism, where concentrations of T4 and T3 are elevated and TSH secretion is suppressed. These have the advantage of discriminating between the concentrations of TSH observed in thyrotoxicosis, compared with the low, but detectable, concentrations that occur in subclinical hyperthyroidism. The performance of this assay has not been established for neonatal specimens. Some drugs and some nonthyroidal patient conditions are known to alter TSH concentrations in vivo.

### INTERPRETATION

PREGNANCY	REFERENCE RANGE FOR TSH IN uIU/mL (As per American Thyroid	
1st Trimester	Association) 0.10-2.50	
2nd Trimester	0.20-3.00	
3rd Trimester	0.30-3.00	

AJAYKUMAR Technologist

Page No: 6 of 12



**Dr. Chandrika Gupta** MBBS.MD ( Path ) RMC NO. 21021/008037



Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 25/06/2023 11:10:03 NAME :- Mr. ASHOK KUMAR

Patient ID :-12231467 Ref. By Dr:- BOB

34 Yrs 11 Mon 25 Days Lab/Hosp :-

Company:- MediWheel

Sex / Age :- Male

Sample Type :- URINE

**Test Name** 

Sample Collected Time 25/06/2023 11:14:11

**Biological Ref Interval** 

Final Authentication: 25/06/2023 12:12:15

Unit

# **CLINICAL PATHOLOGY** Value

			8		
Urine Routine					
PHYSICAL EXAMINATION					
COLOUR	PALE YELI	LOW	PALE YELLOW	PALE YELLOW	
APPEARANCE	Clear		Clear		
CHEMICAL EXAMINATION					
REACTION(PH) Method:- Reagent Strip(Double indicatior blue reaction)	6.0		5.0 - 7.5		
SPECIFIC GRAVITY Method:- Reagent Strip(bromthymol blue)	1.015		1.010 - 1.030		
PROTEIN Method:- Reagent Strip (Sulphosalicylic acid test)	NIL		NIL		
GLUCOSE Method:- Reagent Strip (Glu.Oxidase Peroxidase Benedict)	NIL		NIL		
BILIRUBIN  Method:- Reagent Strip (Azo-coupling reaction)	NEGATIVE		NEGATIVE		
UROBILINOGEN Method:- Reagent Strip (Modified ehrlich reaction)	NORMAL		NORMAL		
KETONES Method:- Reagent Strip (Sodium Nitropruside) Rothera's	NEGATIVE		NEGATIVE		
NITRITE  Method:- Reagent Strip (Diazotization reaction)	NEGATIVE		NEGATIVE		
<b>MICROSCOPY EXAMINATION</b>					
RBC/HPF	NIL	/HPF	NIL		
WBC/HPF	2-3	/HPF	2-3		
EPITHELIAL CELLS	1-2	/HPF	2-3		
CRYSTALS/HPF	ABSENT		ABSENT		
CAST/HPF	ABSENT		ABSENT		
AMORPHOUS SEDIMENT	ABSENT		ABSENT		
BACTERIAL FLORA	ABSENT		ABSENT		
YEAST CELL	ABSENT		ABSENT		
OTHER	ABSENT				

**VIJENDRAMEENA Technologist** 

Page No: 7 of 12





Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 25/06/2023 11:10:03

NAME :- Mr. ASHOK KUMAR

Patient ID :-12231467 Ref. By Dr:- BOB Sex / Age :- Male

34 Yrs 11 Mon 25 Days

Lab/Hosp :-

Company :- MediWheel

Sample Type :- KOx/Na FLUORIDE-F, PLAIN/SEARbible Collected Time 25/06/2023 11:14:11

Final Authentication: 25/06/2023 12:23:26

### **BIOCHEMISTRY**

Test Name	Value	Unit	Biological Ref Interval
FASTING BLOOD SUGAR (Plasma) Method:- GOD PAP	91.4	mg/dl	75.0 - 115.0
Impaired glucose tolerance (IGT)		111 - 125 mg/dL	
Diabetes Mellitus (DM)		> 126 mg/dL	

Instrument Name: Randox Rx Imola Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases.

SERUM CREATININE Method:- Colorimetric Method	0.87	mg/dl	Men - 0.6-1.30 Women - 0.5-1.20
SERUM URIC ACID Method:- Enzymatic colorimetric	6.94	mg/dl	Men - 3.4-7.0 Women - 2.4-5.7

MUKESHSINGH

Page No: 9 of 12





Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 25/06/2023 11:10:03

Patient ID: -12231467 NAME :- Mr. ASHOK KUMAR Ref. By Dr:- BOB

Sex / Age :- Male 34 Yrs 11 Mon 25 Days

Company :- MediWheel Sample Type :- EDTA, URINE

Sample Collected Time 25/06/2023 11:14:11

Final Authentication: 25/06/2023 12:54:28

**HAEMATOLOGY** 

**Test Name** Value Unit **Biological Ref Interval** 

Lab/Hosp:-

**BLOOD GROUP ABO** 

"B" NEGATIVE

BLOOD GROUP ABO Methodology: Haemagglutination reaction Kit Name: Monoclonal agglutinating antibodies (Span clone).

URINE SUGAR (FASTING) Collected Sample Received

Nil

Nil

AJAYSINGH, VIJENDRAMEENA **Technologist** 

Page No: 11 of 12





Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

:- 25/06/2023 11:10:03 Date

NAME :- Mr. ASHOK KUMAR

34 Yrs 11 Mon 25 Days

Ref. By Dr:- BOB

Patient ID: -12231467

Lab/Hosp:-

Company :- MediWheel Sample Type :- PLAIN/SERUM

Sex / Age :- Male

Sample Collected Time 25/06/2023 11:14:11

Final Authentication: 25/06/2023 12:23:26

**BIOCHEMISTRY** 

**Test Name** Value **Biological Ref Interval** Unit

**BLOOD UREA NITROGEN (BUN)** 

13.9

mg/dl

0.0 - 23.0

\*\*\* End of Report \*\*\*

MUKESHSINGH

Page No: 12 of 12





Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 25/06/2023 11:10:03

NAME :- Mr. ASHOK KUMAR

Sex / Age :- Male

34 Yrs 11 Mon 25 Days

Company :- MediWheel

Patient ID: -12231467 Ref. By Doctor:-BOB

Lab/Hosp :-

Final Authentication: 25/06/2023 13:06:31

**BOB PACKAGE BELOW 40MALE** 

# X RAY CHEST PA VIEW:

Both lung fields appears clear.

Bronchovascular markings appear normal.

Trachea is in midline.

Both the hilar shadows are normal.

Both the C.P.angles is clear.

Both the domes of diaphragm are normally placed.

Bony cage and soft tissue shadows are normal.

Heart shadows appear normal.

<u>Impression</u>:- Normal Study

(Please correlate clinically and with relevant further investigations)

\*\*\* End of Report \*\*\*

DR. R.P. BANSAL

M.B.B.S, MD RADIO-DIAGNOSIS SR. CONSULTANT RADIOLOGIST (RMC No. 006640 / 9402)
Page No: 1 of 1

Dr. Piyush Goyal (D.M.R.D.) BILAL



Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date :- 25/06/2023 11:10:03

NAME :- Mr. ASHOK KUMAR

Sex / Age :- Male 34 Yrs 11 Mon 25 Days

Company :- MediWheel

Patient ID :-12231467 Ref. By Doctor:-BOB

Lab/Hosp:-

Final Authentication: 25/06/2023 13:54:53

### **BOB PACKAGE BELOW 40MALE**

# **USG WHOLE ABDOMEN**

**Liver** is of normal size. Echo-texture is normal. No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

**Gall bladder** is of normal size. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

**Pancreas** is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape. Echotexture is normal. No focal lesion is seen.

**Kidneys** are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. No focal lesion is seen. Collecting system does not show any dilatation or calculus.

**Urinary bladder** is well distended and showing smooth wall with normal thickness. Urinary bladder does not show any calculus or mass lesion.

**Prostate** is normal in size with normal echo-texture and outline. No enlarged nodes are visualised. No retro-peritoneal lesion is identified No significant free fluid is seen in peritoneal cavity.

## **IMPRESSION:**

Normal study

Needs clinical correlation for further evaluation

\*\*\* End of Report \*\*\*

**AHSAN** 

Page No: 1 of 1

M.B.B.S., D.M.R.D. RMC Reg No. 017996 Dr. Poonam Gupta MBBS, MD (Radio Diagnosis) RMC No. 32495 Dr. Ashish Choudhary
MBBS, MD (Radio Diagnosis)
Fetal Medicine Consultant
FMF ID - 260517 | RMC No 22430

**Dr. Abhishek Jain**MBBS, DNB, (Radio-Diagnosis)
RMC No. 21687

Transcript by.



Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 25/06/2023 11:10:03

NAME :- Mr. ASHOK KUMAR

Sex / Age :- Male 34 Yrs 11 Mon 25 Days

Company :- MediWheel

Patient ID :-12231467 Ref. By Doctor:-BOB

Lab/Hosp :-

Final Authentication: 25/06/2023 13:54:53

BOB PACKAGE BELOW 40MALE 2D ECHO OPTION TMT (ADULT/CHILD)

# **2D-ECHOCARDIOGRAPHY M.MODE WITH DOPPLER STUDY:**

MITRAL VALVE	NORMAL	TRICUSPID VALVE NORM		
AORTIC VALVE	NORMAL	DI II MONARY VALVE	NORMAL	

TOTTIO VALV		INON	IVIAL	POLIVIONARY VALVE			INORMAL	
		M.MODE	<b>EXAMITATION:</b>					
AO	31	mm	LA	35	Mm	IVS-D	9	mm
VS-S	16	mm	LVID	33	Mm	LVSD	22	mm
LVPW-D	8	mm	LVPW-S	18	Mm	RV		mm
RVWT		mm	EDV		МІ	LVVS		ml
LVEF	62%			RWMA		ABSENT		

		CHA	AMBERS:	
LA	NORMAL	RA	NORMAL	
LV	NORMAL	RV	NORMAL	
PERICARDIUI	M	NORMAL	-	

### **COLOUR DOPPLER:** MITRAL VALVE E VELOCITY 0.88 m/sec PEAK GRADIENT Mm/hg A VELOCITY 0.62 m/sec MEAN GRADIENT Mm/hg MVA BY PHT Cm2 MVA BY PLANIMETRY MITRAL REGURGITATION ABSENT **AORTIC VALVE** PEAK VELOCITY 1.2 m/sec PEAK GRADIENT mm/hg AR VMAX m/sec MEAN GRADIENT mm/hg **AORTIC REGURGITATION** ABSENT TRICUSPID VALVE PEAK VELOCITY m/sec PEAK GRADIENT mm/hg MEAN VELOCITY m/sec MEAN GRADIENT mm/hg VMax VELOCITY TRICUSPID REGURGITATION ABSENT **PULMONARY VALVE** PEAK VELOCITY 0.98 M/sec. PEAK GRADIENT Mm/hg MEAN VALOCITY MEAN GRADIENT Mm/hg PULMONARY REGURGITATION ABSENT

Page No: 1 of 2

**AHSAN** 

Dr. Piyush Goyal M.B.B.S., D.M.R.D. RMC Reg No. 017996 **Dr. Poonam Gupta**MBBS, MD (Radio Diagnosis)
RMC No. 32495

Dr. Ashish Choudhary
MBBS, MD (Radio Diagnosis)
Fetal Medicine Consultant
FMF ID - 260517 | RMC No 22430

**Dr. Abhishek Jain**MBBS, DNB, (Radio-Diagnosis)
RMC No. 21687

Transcript by.



Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

**ii a ia a a a a a** a a a i ia a i ia **a** ia

Date :- 25/06/2023 11:10:03

NAME :- Mr. ASHOK KUMAR

Sex / Age :- Male

34 Yrs 11 Mon 25 Days

Company :- MediWheel

Patient ID :-12231467 Ref. By Doctor:-BOB

Lab/Hosp :-

Final Authentication: 25/06/2023 13:54:53

# Impression--

- 1. Normal LV size & contractility
- 2. No RWMA, LVEF 62 %.
- 3. Normal cardiac chamber.
- 4. Normal valve
- 5. No clot, no vegetation, no pericardial effusion. (Cardiologist)

\*\*\* End of Report \*\*\*

Page No: 2 of 2

AHSAN

