

Age / Gender: 32 years / Male

Patient ID: 15576

Source: MEDI WHEEL

Referral: SELF

Collection Time : Jan 14, 2023, 11:29 a.m. **Reporting Time :** Jan 14, 2023, 02:10 p.m.

Sample ID:

			000700022	
Test Description	Value(s)	Reference Range	Unit	
CBC; Complete Blood Count				
Hemoglobin (Hb)* Method : Cynmeth Photometric Measurement	15.0	13.5 - 18.0	gm/dL	
Erythrocyte (RBC) Count* Method : Electrical Impedence	5.34	4.7 - 6.0	mil/cu.mm	
Packed Cell Volume (PCV)* Method : Calculated	43	42 - 52	%	
Mean Cell Volume (MCV)* Method : Electrical Impedence	81	78 - 100	fL	
Mean Cell Haemoglobin (MCH)* Method : Calculated	28	27 - 31	pg	
Mean Corpuscular Hb Concn. (MCHC)* Method : Calculated	35	32 - 36	gm/dL	
Red Cell Distribution Width (RDW)* Method : Electrical Impedence	13.5	11.5 - 14.0	%	
Total Leucocytes (WBC) Count* Method : Electrical Impedence	6300	4000-10000	cell/cu.mm	
Neutrophils* Method : VCSn Technology	61	40 - 80	%	
Lymphocytes* Method : VCSn Technology	33	20 - 40	%	
Monocytes* Method : VCSn Technology	5	2 - 10	%	
Eosinophils* Method : VCSn Technology	1	1 - 6	%	
Basophils	0	0 - 1		
Platelet Count* Method : Electrical Impedence	2.58	1.5 - 4.5	10^3/ul	
Mean Platelet Volume (MPV)* Method : Electrical Impedence	7.0	7.2 - 11.7	fL	

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Approved by





Age / Gender: 32 years / Male

Patient ID: 15576

Source: MEDI WHEEL

Referral: SELF

Collection Time : Jan 14, 2023, 11:29 a.m. **Reporting Time :** Jan 14, 2023, 02:10 p.m.

Sample ID:

Test Description	Value(s)	Reference Range	Unit
PCT*	0.18	0.2 - 0.5	%
Method : Calculated	447	0.0.470	0/
PDW* Method : Calculated	14.7	9.0 - 17.0	%

Tests done on Automated Three Part Cell Counter. (WBC, RBC, Platelet count by impedance method, colorimetric method for Hemoglobin, WBC differential by flow cytometry using laser technology other parameters are calculated). All Abnormal Haemograms are reviewed confirmed microscopically.

Esr, Erythrocyte Sedimentation Rate

Esr, Erythrocyte Sedimentation Rate (Westergren)

20

0-10

mm/hr

Interpretation:

- It indicates presence and intensity of an inflammatory process. It does not diagnose a specific disease. Changes in the ESR are more significant than the abnormal results of a single test.
- It is a prognostic test and used to monitor the course or response to treatment of diseases like tuberculosis, bacterial endocarditis, acute rheumatic fever, rheumatoid arthritis, SLE, Hodgkins disease, temporal arteritis and polymyalgia rheumatica.
- It is also increased in pregnancy, multiple myeloma, menstruation, and hypothyroidism.

U	r	n	е	К	0	u	tı	n	е

Colour*	Yellow		
Volume*	15	-	ml
Transparency (Appearance)*	Clear	Clear	
Reaction (pH)*	5.0	4.5 - 8	
Specific Gravity*	1.025	1.010 - 1.030	

Chemical Examination (Automated Dipstick Method) Urine

Urine Glucose*	Negative	Negative
Urine Protein*	Negative	Negative
Urine Ketone*	Negative	Negative

Dr.CH.Deepthi Chandrika M.D. Pathology

Reg.No.APCM/FMR/77174

Approved by

Scan to Validate



— Flower us —



Age / Gender: 32 years / Male

Patient ID: 15576

Source: MEDI WHEEL

Referral: SELF

Collection Time : Jan 14, 2023, 11:29 a.m.

Reporting Time : Jan 14, 2023, 02:10 p.m.

Sample ID :

Test Description	Value(s)	Reference Range	Unit
Dlood*	Nogotivo	Negative	
Blood*	Negative	Negative	
Bilirubin*	Negative	Negative	
Nitrite*	Negative	Negative	
Leucocytes*	Negative	Negative	
Jrobilinogen*	Normal	Normal	
Microscopic Examination Urine			
Pus Cells (WBCs)*	2-3	0 - 5	/hpf
Epithelial Cells*	1-2	0 - 4	/hpf
Red blood Cells*	Absent	Absent	/hpf
Crystals*	Absent	Absent	
Cast*	Absent	Absent	
Bacteria*	Absent	Absent	

Stool Complete Exam

Blood Group & Rh Type

Blood Grouping & Rh Typing

"B" + (POSITIVE)

Method : Forward and Reverse By Tube Method

Methodology

This is done by forward and reverse grouping by tube Agglutination method.

Interpretation

Newborn baby does not produce ABO antibodies until 3 to 6 months of age. So the blood group of the Newborn baby is done by ABO antigen grouping (forward grouping) only, antibody grouping (reverse grouping) is not required. Confirmation of the New-born's blood group is indicated when the A and B antigen expression and the isoagglutinins are fully developed (2–4 years).

Fasting - Glucose

Dr.CH.Deepthi Chandrika M.D. Pathology

Reg.No.APCM/FMR/77174

Approved by

Scan to Validate



m - 2, — 15mer 22777 🕓 f 🔮



Age / Gender: 32 years / Male

Patient ID: 15576

Source: MEDI WHEEL

Referral: SELF

Collection Time : Jan 14, 2023, 11:29 a.m. **Reporting Time :** Jan 14, 2023, 02:10 p.m.

Sample ID :

Test Description	Value(s)	Reference Range	Unit
Glucose Fasting*	82	Normal: 70-100	mg/dL
Method : Plasma, Hexokinase		Impaired Fasting Glucose (IFG):	
		100-125	
		Diabetes Mellitus: >= 126	
		(On more than one occasion)	
		(American Diabetes Association	
		guidelines 2017)	
Post Prandial Blood Sugar			
Blood Glucose-Post Prandial*	100	80-140	mg/dL
Method : Plasma - P, Hexokinase			· ·
Fasting Urine Sugar			
Fasting Urine Sugar	NEGATIVE	NEGATIVE -	
Post Prandial Urine Sugar			
Post Prandial Urine Sugar	NEGATIVE		
HBA1C (Glycosylated Haemoglobin)			
Glyco Hb (HbA1C)	5.3	Non-Diabetic: <=5.9	%
Method : EDTA Whole blood,HPLC		Pre Diabetic:6.0-6.4	
		Diabetic: >=6.5	
Estimated Average Glucose :	105.41		mg/dL
Interpretations			

- 1. HbA1C has been endorsed by clinical groups and American Diabetes Association guidelines 2017 for diagnosing diabetes using a cut off point of 6.5%
- 2. Low glycated haemoglobin in a non diabetic individual are often associated with systemic inflammatory diseases, chronic anaemia (especially severe iron deficiency and haemolytic), chronic renal failure and liver diseases. Clinical correlation suggested.

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Approved by





Age / Gender: 32 years / Male

Patient ID: 15576

Source: MEDI WHEEL

Referral: SELF

Collection Time : Jan 14, 2023, 11:29 a.m. **Reporting Time :** Jan 14, 2023, 02:10 p.m.

Sample ID :

Test Description Value(s) Reference Range Unit

3. In known diabetic patients, following values can be considered as a tool for monitoring the glycemic control.

Excellent control-6-7 %
Fair to Good control – 7-8 %
Unsatisfactory control – 8 to 10 %

Poor Control – More than 10 %

Thyroid Function Test (TFT)			
TRI-IODO THYRONINE (T3)	174	60 - 181	ng/dL
Method : CLIA			
TOTAL THYROXINE (T4)	8.3	4.2 - 12.0	ug/dL
Method : CLIA			
THYROID STIMULATING HORMONE (TSH)	1.60	0.46 – 8.10 : 1 Yrs – 5 Yrs	uIU/mL
Method : CLIA		0.36 – 5.80 : 6 Yrs – 18 Yrs	
		0.35 – 5.50 : >18 Yrs	
		Pregnancy Ranges	
		1st Trimester :0.1 - 2.5	
		2nd Trimester :0.2 - 3.0	
		3rd Trimester:0.3 - 3.0	

Comments:

IF NOT ON DRUGS SUGGESTED FT3 & FT4 ESTIMATION

Please correlate with clinical conditions.

Note: Serum T3, T4 and TSH form the three components of thyroid screening panel, useful in diagnosing various disorders of the thyroid gland. Primary Hypothyroidism is accompanied by depressed serum T3 and T4 values and elevated serum TSH levels. Although elevated TSH levels are nearly always indicative of Primary Hypothyroidism, rarely they can from TSH secreting pituitary tumors (Secondary hyperthyroidism)To confirm diagnosis - evaluate FT3 and FT4.

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Approved by

Scan to Validate



— Flower us —

© f © ©



Age / Gender: 32 years / Male

Patient ID: 15576

Source: MEDI WHEEL

Referral: SELF

Collection Time : Jan 14, 2023, 11:29 a.m. **Reporting Time :** Jan 14, 2023, 02:10 p.m.

Sample ID :

Test Description	Value(s)	Reference Range	Unit
<u>Lipid Profile</u>			
Cholesterol-Total Method : Serum, Cholesterol oxidase esterase, peroxidase	219	Desirable: <= 200 Borderline High: 201-239 High: > 239 Ref: The National Cholesterol Education Program (NCEP) Adult Treatment Panel III Report.	mg/dL
Triglycerides Method : Serum, Enzymatic, endpoint	109	Normal: < 150 Borderline High: 150-199 High: 200-499 Very High: >= 500	mg/dL
Cholesterol-HDL Direct Method : Serum, Direct measure-PEG	42	<40: Low 40 - 60: Optimal > 60: Desirable	mg/dL
LDL Cholesterol Method : Serum	155.20	Optimal: < 100 Near optimal/above optimal: 100-129 Borderline high: 130-159 High: 160-189 Very High: >= 190	mg/dL
Non - HDL Cholesterol, Serum Method : calculated	177	Desirable: < 130 mg/dL Borderline High: 130-159mg/dL High: 160-189 mg/dL Very High: > or = 190 mg/dL	mg/dL
VLDL Cholesterol Method : calculated	21.80	6 - 38	mg/dL
CHOL/HDL RATIO Method : calculated	5.21	3.5 - 5.0	ratio

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Odepter

Approved by





Age / Gender: 32 years / Male

Patient ID: 15576

Source: MEDI WHEEL

Referral: SELF

Collection Time: Jan 14, 2023, 11:29 a.m.

Reporting Time: Jan 14, 2023, 02:10 p.m.

Sample ID :

Test Description	Value(s)	Reference Range	Unit
LDL/HDL RATIO	3.70	Desirable / low risk - 0.5 -3.0	ratio
Method : calculated		Low/ Moderate risk - 3.0- 6.0	
		Elevated / High risk - > 6.0	
HDL/LDL RATIO	0.27	Desirable / low risk - 0.5 -3.0	ratio
Method : calculated	V.=.	Low/ Moderate risk - 3.0- 6.0	
		Elevated / High risk - > 6.0	
Note: 8-10 hours fasting sample is required.			
KIDNEY FUNCTION TEST			
Urea *	16	15- 50	mg/dL
Method : Serum			
Blood Urea Nitrogen-BUN*	7.48	7 - 24	mg/dL
Method : Serum, Urease			
Uric Acid*	7.7	3.5 - 7.2	mg/dL
Method : Serum, Uricase/POD			
Creatinine*	0.62	0.7 - 1.3	mg/dL
Method : Serum, Jaffe IDMS			
Liver Funtion Test (LFT) with GGT			
Bilirubin - Total	0.5	0.3 - 1.2	mg/dL
Method : Serum, Jendrassik Grof			
Bilirubin - Direct	0.2	Adults and Children: < 0.2	mg/dL
Method : Serum, Diazotization			
Bilirubin - Indirect	0.30	0.1 - 1.0	mg/dL
Method : Serum, Calculated			
SGOT	26	< 50	U/L
Method : Serum, UV with P5P, IFCC 37 degree			
SGPT	15	< 50	U/L
Method : Serum, UV with P5P, IFCC 37 degree			

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Odepter

Approved by





Age / Gender: 32 years / Male

Patient ID: 15576

Source: MEDI WHEEL

Referral: SELF

Sample ID:

Collection Time: Jan 14, 2023, 11:29 a.m.

Reporting Time : Jan 14, 2023, 02:10 p.m.

Test Description	Value(s)	Reference Range	Unit
SGOT/SGPT	1.73	0.7 - 1.4	ratio
Method : calculated			
GGT-Gamma Glutamyl Transpeptidase	-	< 55	U/L
Method : Serum, G-glutamyl-carboxy-nitoanilide			
Alkaline Phosphatase-ALPI	78	30-120	U/L
Method : Serum, PNPP, AMP Buffer, IFCC 37 degree			
Total Protein	8.0	6.6 - 8.3	g/dL
Method : Serum, Biuret, reagent blank end point			
Albumin	4.4	Adults: 3.5 - 5.2	g/dL
Method : Serum, Bromcresol purple			
Globulin	3.6	1.8 - 3.6	g/dL
Method : Calculated			
A/G Ratio	1.22	1.2 - 2.2	ratio
Method : Calculated			

END OF REPORT

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Odepter

Reg. No. APCIVI/FIVIR///I

Approved by

