

DIAGNOSTIC REPORT



Cert. No. MC-2351



CLIENT CODE : C000138383

CLIENT'S NAME AND ADDRESS :
ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd
24 SCO, SECTOR 11 D
CHANDIGARH, 160011
PUNJAB, INDIA
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956

PATIENT NAME : RAM CHARAN

PATIENT ID : RAMCM24127080

ACCESSION NO : 0080WC011145 **AGE :** 52 Years **SEX :** Male

ABHA NO :

DRAWN : **RECEIVED :** 28/03/2023 08:54

REPORTED : 28/03/2023 15:29

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

MEDI WHEEL FULL BODY HEALTH CHECK UP ABOVE 40 MALE

BLOOD COUNTS,EDTA WHOLE BLOOD

HEMOGLOBIN (HB)	14.3	13.0 - 17.0	g/dL
METHOD : CYANMETHEMOGLOBIN METHOD			
RED BLOOD CELL (RBC) COUNT	4.78	4.5 - 5.5	mil/ μ L
WHITE BLOOD CELL (WBC) COUNT	7.00	4.0 - 10.0	thou/ μ L
PLATELET COUNT	171	150 - 410	thou/ μ L

RBC AND PLATELET INDICES

HEMATOCRIT (PCV)	42.4	40.0 - 50.0	%
MEAN CORPUSCULAR VOLUME (MCV)	88.8	83.0 - 101.0	fL
METHOD : DERIVED PARAMETER FROM RBC HISTOGRAM			
MEAN CORPUSCULAR HEMOGLOBIN (MCH)	30.0	27.0 - 32.0	pg
METHOD : CALCULATED PARAMETER			
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION (MCHC)	33.8	31.5 - 34.5	g/dL
METHOD : CALCULATED PARAMETER			
RED CELL DISTRIBUTION WIDTH (RDW)	13.6	11.6 - 14.0	%
METHOD : CALCULATED PARAMETER			
MENTZER INDEX	18.6		
MEAN PLATELET VOLUME (MPV)	11.8	High 6.8 - 10.9	fL
METHOD : DERIVED PARAMETER FROM PLATELET HISTOGRAM			

WBC DIFFERENTIAL COUNT

NEUTROPHILS	46	40 - 80	%
METHOD : LIGHT ABSORBANCE OF CYTCHEMICAL STAINED CELLS IMPEDENCE			
LYMPHOCYTES	44	High 20 - 40	%
METHOD : LIGHT ABSORBANCE OF CYTCHEMICAL STAINED CELLS IMPEDENCE			
MONOCYTES	6	2.0 - 10.0	%
METHOD : LIGHT ABSORBANCE OF CYTCHEMICAL STAINED CELLS IMPEDENCE			
EOSINOPHILS	4	1.0 - 6.0	%
BASOPHILS	0	0 - 1	%
METHOD : LIGHT ABSORBANCE OF CYTCHEMICAL STAINED CELLS IMPEDENCE			
ABSOLUTE NEUTROPHIL COUNT	3.22	2.0 - 7.0	thou/ μ L
ABSOLUTE LYMPHOCYTE COUNT	3.08	High 1.0 - 3.0	thou/ μ L
ABSOLUTE MONOCYTE COUNT	0.42	0.2 - 1.0	thou/ μ L
ABSOLUTE EOSINOPHIL COUNT	0.28	0.02 - 0.50	thou/ μ L
ABSOLUTE BASOPHIL COUNT	0	Low 0.02 - 0.10	thou/ μ L



Scan to View Details



Scan to View Report

DIAGNOSTIC REPORT



Cert. No. MC-2351



CLIENT CODE : C000138383

CLIENT'S NAME AND ADDRESS :
 ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
 F-703, LADO SARAI, MEHRAULI
 SOUTH WEST DELHI
 NEW DELHI 110030
 DELHI INDIA
 8800465156

SRL Ltd
 24 SCO, SECTOR 11 D
 CHANDIGARH, 160011
 PUNJAB, INDIA
 Tel : 9111591115, Fax :
 CIN - U74899PB1995PLC045956

PATIENT NAME : RAM CHARAN

PATIENT ID : RAMCM24127080

ACCESSION NO : 0080WC011145 **AGE :** 52 Years **SEX :** Male

ABHA NO :

DRAWN : **RECEIVED :** 28/03/2023 08:54

REPORTED : 28/03/2023 15:29

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

METHOD : CALCULATED PARAMETER

NEUTROPHIL LYMPHOCYTE RATIO (NLR) 1.0

METHOD : CALCULATED PARAMETER

ERYTHROCYTE SEDIMENTATION RATE (ESR),WHOLE BLOOD

E.S.R 05 0 - 14 mm at 1 hr

METHOD : MODIFIED WESTERGRN

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD

HBA1C 5.3 Non-diabetic Adult < 5.7 %

Pre-diabetes 5.7 - 6.4
 Diabetes diagnosis: > or = 6.5
 Therapeutic goals: < 7.0
 Action suggested : > 8.0
 (ADA Guideline 2021)

ESTIMATED AVERAGE GLUCOSE(EAG) 105.4 < 116.0 mg/dL

GLUCOSE FASTING,FLUORIDE PLASMA

FBS (FASTING BLOOD SUGAR) 104 74 - 106 mg/dL

METHOD : HEXOKINASE

GLUCOSE, POST-PRANDIAL, PLASMA

PPBS(POST PRANDIAL BLOOD SUGAR) 112 Non-Diabetes mg/dL

70 - 140

METHOD : HEXOKINASE

LIPID PROFILE, SERUM

CHOLESTEROL, TOTAL 192 < 200 Desirable mg/dL

200 - 239 Borderline High
 >/= 240 High

METHOD : CHOLESTEROL OXIDASE, ESTERASE,PEROXIDASE

TRIGLYCERIDES 166 **High** < 150 Normal mg/dL

150 - 199 Borderline High
 200 - 499 High
 >/= 500 Very High

METHOD : ENZYMATIC ASSAY

HDL CHOLESTEROL 45 < 40 Low mg/dL

>/=60 High

METHOD : DIRECT MEASURE - PEG



Scan to View Details



Scan to View Report

DIAGNOSTIC REPORT



Cert. No. MC-2351



CLIENT CODE : C000138383

CLIENT'S NAME AND ADDRESS :
 ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
 F-703, LADO SARAI, MEHRAULI
 SOUTH WEST DELHI
 NEW DELHI 110030
 DELHI INDIA
 8800465156

SRL Ltd
 24 SCO, SECTOR 11 D
 CHANDIGARH, 160011
 PUNJAB, INDIA
 Tel : 9111591115, Fax :
 CIN - U74899PB1995PLC045956

PATIENT NAME : RAM CHARAN

PATIENT ID : RAMCM24127080

ACCESSION NO : 0080WC011145 **AGE :** 52 Years **SEX :** Male

ABHA NO :

DRAWN : **RECEIVED :** 28/03/2023 08:54

REPORTED : 28/03/2023 15:29

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
CHOLESTEROL LDL		114	High < 100 Optimal 100 - 129 Near or above optimal 130 - 159 Borderline High 160 - 189 High >/= 190 Very High	mg/dL
METHOD : CHOLESTEROL OXIDASE, ESTERASE, PEROXIDASE				
NON HDL CHOLESTEROL		147	High Desirable: Less than 130 Above Desirable: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very high: > or = 220	mg/dL
METHOD : CALCULATED PARAMETER				
VERY LOW DENSITY LIPOPROTEIN		33.2	Desirable value :	mg/dL
METHOD : CALCULATED PARAMETER				
CHOL/HDL RATIO		4.3	3.3-4.4 Low Risk 4.5-7.0 Average Risk 7.1-11.0 Moderate Risk > 11.0 High Risk	
METHOD : CALCULATED PARAMETER				
LDL/HDL RATIO		2.5	0.5 - 3.0 Desirable/Low Risk 3.1 - 6.0 Borderline/Moderate Risk >6.0 High Risk	
METHOD : CALCULATED PARAMETER				



Scan to View Details



Scan to View Report



CLIENT CODE : C000138383

CLIENT'S NAME AND ADDRESS :
ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd
24 SCO, SECTOR 11 D
CHANDIGARH, 160011
PUNJAB, INDIA
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956

PATIENT NAME : RAM CHARAN

PATIENT ID : RAMCM24127080

ACCESSION NO : 0080WC011145 **AGE :** 52 Years **SEX :** Male

ABHA NO :

DRAWN : **RECEIVED :** 28/03/2023 08:54

REPORTED : 28/03/2023 15:29

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

Interpretation(s)

- Cholesterol levels help assess the patient risk status and to follow the progress of patient under treatment to lower serum cholesterol concentrations.
- Serum Triglyceride (TG) are a type of fat and a major source of energy for the body. Both quantity and composition of the diet impact on plasma triglyceride concentrations. Elevations in TG levels are the result of overproduction and impaired clearance. High TG are associated with increased risk for CAD (Coronary artery disease) in patients with other risk factors, such as low HDL-C, some patient groups with elevated apolipoprotein B concentrations, and patients with forms of LDL that may be particularly atherogenic.
- HDL-C plays a crucial role in the initial step of reverse cholesterol transport, this considered to be the primary atheroprotective function of HDL
- LDL -C plays a key role in causing and influencing the progression of atherosclerosis and, in particular, coronary sclerosis. The majority of cholesterol stored in atherosclerotic plaques originates from LDL, thus LDL-C value is the most powerful clinical predictor.
- Non HDL cholesterol: Non-HDL-C measures the cholesterol content of all atherogenic lipoproteins, including LDL hence it is a better marker of risk in both primary and secondary prevention studies. Non-HDL-C also covers, to some extent, the excess ASCVD risk imparted by the sdLDL, which is significantly more atherogenic than the normal large buoyant particles, an elevated non-HDL-C indirectly suggests greater proportion of the small, dense variety of LDL particles

Serum lipid profile is measured for cardiovascular risk prediction. Lipid Association of India recommends LDL-C as primary target and Non HDL-C as co-primary treatment target.

Risk Stratification for ASCVD (Atherosclerotic cardiovascular disease) by Lipid Association of India

Risk Category	
Extreme risk group	A. CAD with > 1 feature of high risk group B. CAD with > 1 feature of Very high risk group or recurrent ACS (within 1 year) despite LDL-C < or = 50 mg/dl or polyvascular disease
Very High Risk	1. Established ASCVD 2. Diabetes with 2 major risk factors or evidence of end organ damage 3. Familial Homozygous Hypercholesterolemia
High Risk	1. Three major ASCVD risk factors. 2. Diabetes with 1 major risk factor or no evidence of end organ damage. 3. CKD stage 3B or 4. 4. LDL >190 mg/dl 5. Extreme of a single risk factor. 6. Coronary Artery Calcium - CAC >300 AU. 7. Lipoprotein a >= 50mg/dl 8. Non stenotic carotid plaque
Moderate Risk	2 major ASCVD risk factors
Low Risk	0-1 major ASCVD risk factors
Major ASCVD (Atherosclerotic cardiovascular disease) Risk Factors	
1. Age > or = 45 years in males and > or = 55 years in females	3. Current Cigarette smoking or tobacco use
2. Family history of premature ASCVD	4. High blood pressure
5. Low HDL	

Newer treatment goals and statin initiation thresholds based on the risk categories proposed by LAI in 2020.

Risk Group	Treatment Goals		Consider Drug Therapy	
	LDL-C (mg/dl)	Non-HDL (mg/dl)	LDL-C (mg/dl)	Non-HDL (mg/dl)
Extreme Risk Group Category A	<50 (Optional goal <OR = 30)	< 80 (Optional goal <OR = 60)	>OR = 50	>OR = 80



Scan to View Details



Scan to View Report

DIAGNOSTIC REPORT



Cert. No. MC-2351



CLIENT CODE : C000138383

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd
24 SCO, SECTOR 11 D
CHANDIGARH, 160011
PUNJAB, INDIA
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956

PATIENT NAME : RAM CHARAN

PATIENT ID : RAMCM24127080

ACCESSION NO : 0080WC011145 **AGE :** 52 Years **SEX :** Male

ABHA NO :

DRAWN : **RECEIVED :** 28/03/2023 08:54

REPORTED : 28/03/2023 15:29

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

Extreme Risk Group Category B	<OR = 30	<OR = 60	> 30	>60
Very High Risk	<50	<80	>OR= 50	>OR= 80
High Risk	<70	<100	>OR= 70	>OR= 100
Moderate Risk	<100	<130	>OR= 100	>OR= 130
Low Risk	<100	<130	>OR= 130*	>OR= 160

*After an adequate non-pharmacological intervention for at least 3 months.

References: Management of Dyslipidaemia for the Prevention of Stroke: Clinical Practice Recommendations from the Lipid Association of India. Current Vascular Pharmacology, 2022, 20, 134-155.

LIVER FUNCTION PROFILE, SERUM

BILIRUBIN, TOTAL METHOD : DIAZONIUM ION, BLANKED (ROCHE)	0.48	UPTO 1.2	mg/dL
BILIRUBIN, DIRECT METHOD : DIAZOTIZATION	0.13	0.00 - 0.30	mg/dL
BILIRUBIN, INDIRECT METHOD : CALCULATED PARAMETER	0.35	0.00 - 0.60	mg/dL
TOTAL PROTEIN METHOD : BIURET	7.3	6.6 - 8.7	g/dL
ALBUMIN METHOD : BROMOCRESOL GREEN	4.6	3.97 - 4.94	g/dL
GLOBULIN METHOD : CALCULATED PARAMETER	2.7	2.0 - 4.0 Neonates - Pre Mature: 0.29 - 1.04	g/dL
ALBUMIN/GLOBULIN RATIO METHOD : CALCULATED PARAMETER	1.7	1.0 - 2.0	RATIO
ASPARTATE AMINOTRANSFERASE (AST/SGOT)	19	0 - 40	U/L
ALANINE AMINOTRANSFERASE (ALT/SGPT) METHOD : UV WITHOUT PYRIDOXAL-5 PHOSPHATE	24	0 - 41	U/L
ALKALINE PHOSPHATASE METHOD : PNPP - AMP BUFFER	88	40 - 129	U/L
GAMMA GLUTAMYL TRANSFERASE (GGT) METHOD : GAMMA GLUTAMYL CARBOXY 4NITROANILIDE	21	8 - 61	U/L
LACTATE DEHYDROGENASE METHOD : LACTATE -PYRUVATE	168	135 - 225	U/L

BLOOD UREA NITROGEN (BUN), SERUM

BLOOD UREA NITROGEN METHOD : UREASE - UV	9	6 - 20	mg/dL
---	---	--------	-------



DIAGNOSTIC REPORT



Cert. No. MC-2351



CLIENT CODE : C000138383

CLIENT'S NAME AND ADDRESS :
 ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
 F-703, LADO SARAI, MEHRAULI
 SOUTH WEST DELHI
 NEW DELHI 110030
 DELHI INDIA
 8800465156

SRL Ltd
 24 SCO, SECTOR 11 D
 CHANDIGARH, 160011
 PUNJAB, INDIA
 Tel : 9111591115, Fax :
 CIN - U74899PB1995PLC045956

PATIENT NAME : RAM CHARAN

PATIENT ID : RAMCM24127080

ACCESSION NO : 0080WC011145 **AGE :** 52 Years **SEX :** Male

ABHA NO :

DRAWN : **RECEIVED :** 28/03/2023 08:54

REPORTED : 28/03/2023 15:29

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

CREATININE, SERUM

CREATININE 0.88 0.70 - 1.20 mg/dL
 METHOD : ALKALINE PICRATE-KINETIC

BUN/CREAT RATIO

BUN/CREAT RATIO 10.23 5.00 - 15.00
 METHOD : CALCULATED PARAMETER

URIC ACID, SERUM

URIC ACID 5.6 3.4 - 7.0 mg/dL
 METHOD : URICASE, COLORIMETRIC

TOTAL PROTEIN, SERUM

TOTAL PROTEIN 7.3 6.6 - 8.7 g/dL
 METHOD : BIURET

ALBUMIN, SERUM

ALBUMIN 4.6 3.97 - 4.94 g/dL
 METHOD : BROMOCRESOL GREEN

GLOBULIN

GLOBULIN 2.7 2.0 - 4.0 g/dL
 Neonates -
 Pre Mature:
 0.29 - 1.04
 METHOD : CALCULATED PARAMETER

ELECTROLYTES (NA/K/CL), SERUM

SODIUM, SERUM 136 136 - 145 mmol/L
 METHOD : ISE INDIRECT

POTASSIUM, SERUM 4.08 3.5 - 5.1 mmol/L
 METHOD : ISE INDIRECT

CHLORIDE, SERUM 99 98 - 107 mmol/L
 METHOD : ISE INDIRECT



Scan to View Details



Scan to View Report

DIAGNOSTIC REPORT



Cert. No. MC-2351

CLIENT CODE : C000138383

CLIENT'S NAME AND ADDRESS :
ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd
24 SCO, SECTOR 11 D
CHANDIGARH, 160011
PUNJAB, INDIA
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956

PATIENT NAME : RAM CHARAN

PATIENT ID : RAMCM24127080

ACCESSION NO : 0080WC011145 **AGE :** 52 Years **SEX :** Male

ABHA NO :

DRAWN : **RECEIVED :** 28/03/2023 08:54

REPORTED : 28/03/2023 15:29

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

Interpretation(s)

Sodium	Potassium	Chloride
Decreased in: CCF,cirrhosis, vomiting, diarrhea, excessive sweating, salt-losing nephropathy,adrenal insufficiency, nephrotic syndrome, water intoxication, SIADH. Drugs: thiazides, diuretics, ACE inhibitors, chlorpropamide,carbamazepine,anti depressants (SSRI), antipsychotics.	Decreased in: Low potassium intake,prolonged vomiting or diarrhea, RTA types I and II, hyperaldosteronism, Cushing's syndrome,osmotic diuresis (e.g., hyperglycemia),alkalosis, familial periodic paralysis,trauma (transient).Drugs: Adrenergic agents, diuretics.	Decreased in: Vomiting, diarrhea, renal failure combined with salt deprivation, over-treatment with diuretics, chronic respiratory acidosis, diabetic ketoacidosis, excessive sweating, SIADH, salt-losing nephropathy, porphyria, expansion of extracellular fluid volume, adrenalinsufficiency, hyperaldosteronism,metabolic alkalosis. Drugs: chronic laxative,corticosteroids, diuretics.
Increased in: Dehydration (excessivesweating, severe vomiting or diarrhea),diabetes mellitus, diabetesinsipidus, hyperaldosteronism, inadequate water intake. Drugs: steroids, licorice,oral contraceptives.	Increased in: Massive hemolysis, severe tissue damage, rhabdomyolysis, acidosis, dehydration,renal failure, Addison' s disease, RTA type IV, hyperkalemic familial periodic paralysis. Drugs: potassium salts, potassium- sparing diuretics,NSAIDs, beta-blockers, ACE inhibitors, high-dose trimethoprim-sulfamethoxazole.	Increased in: Renal failure, nephrotic syndrome, RTA,dehydration, overtreatment with saline,hyperparathyroidism, diabetes insipidus, metabolic acidosis from diarrhea (Loss of HCO3-), respiratory alkalosis,hyperadrenocorticism. Drugs: acetazolamide,androgens, hydrochlorothiazide,salicylates.
Interferences: Severe lipemia or hyperproteinemi, if sodium analysis involves a dilution step can cause spurious results. The serum sodium falls about 1.6 mEq/L for each 100 mg/dL increase in blood glucose.	Interferences: Hemolysis of sample, delayed separation of serum, prolonged fist clenching during blood drawing, and prolonged tourniquet placement. Very high WBC/PLT counts may cause spurious. Plasma potassium levels are normal.	Interferences: Test is helpful in assessing normal and increased anion gap metabolic acidosis and in distinguishing hypercalcemia due to hyperparathyroidism (high serum chloride) from that due to malignancy (Normal serum chloride)

PHYSICAL EXAMINATION, URINE

COLOR PALE YELLOW
APPEARANCE CLEAR

CHEMICAL EXAMINATION, URINE

PH 5.5 4.7 - 7.5
METHOD : REFLECTANCE SPECTROPHOTOMETRY- DOUBLE INDICATOR METHOD

SPECIFIC GRAVITY 1.005 1.003 - 1.035
METHOD : REFLECTANCE SPECTROPHOTOMETRY (PKA CHANGE OF PRETREATED POLY ELECTROLYTES)

PROTEIN NOT DETECTED NOT DETECTED
METHOD : REFLECTANCE SPECTROPHOTOMETRY (PROTEIN-ERROR-OF-INDICATORS PRINCIPLE)

GLUCOSE NOT DETECTED NOT DETECTED
METHOD : REFLECTANCE SPECTROPHOTOMETRY(GLUCOSE OXIDAE/PEROXIDASE METHOD)

KETONES NOT DETECTED NOT DETECTED
METHOD : REFLECTANCE SPECTROPHOTOMETRY (SODIUM NITROPRUSSIDE REACTION)

BLOOD NOT DETECTED NOT DETECTED
METHOD : REFLECTANCE SPECTROPHOTOMETRY (PEROXIDASE METHOD)

BILIRUBIN NOT DETECTED NOT DETECTED



Scan to View Details



Scan to View Report

DIAGNOSTIC REPORT



Cert. No. MC-2351



CLIENT CODE : C000138383

CLIENT'S NAME AND ADDRESS :
 ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
 F-703, LADO SARAI, MEHRAULI
 SOUTH WEST DELHI
 NEW DELHI 110030
 DELHI INDIA
 8800465156

SRL Ltd
 24 SCO, SECTOR 11 D
 CHANDIGARH, 160011
 PUNJAB, INDIA
 Tel : 9111591115, Fax :
 CIN - U74899PB1995PLC045956

PATIENT NAME : RAM CHARAN

PATIENT ID : RAMCM24127080

ACCESSION NO : 0080WC011145 **AGE :** 52 Years **SEX :** Male

ABHA NO :

DRAWN : **RECEIVED :** 28/03/2023 08:54

REPORTED : 28/03/2023 15:29

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

METHOD : REFLECTANCE SPECTROPHOTOMETRY (DIAZO REACTION)				
UROBILINOGEN		NORMAL	NORMAL	
METHOD : REFLECTANCE SPECTROPHOTOMETRY - EHRlich REACTION				
NITRITE		NOT DETECTED	NOT DETECTED	
METHOD : REFLECTANCE SPECTROPHOTOMETRY, CONVERSION OF NITRATE TO NITRITE				
LEUKOCYTE ESTERASE		NOT DETECTED	NOT DETECTED	
MICROSCOPIC EXAMINATION, URINE				
RED BLOOD CELLS		NOT DETECTED	NOT DETECTED	/HPF
METHOD : MICROSCOPIC EXAMINATION				
PUS CELL (WBC'S)		1-2	0-5	/HPF
METHOD : MICROSCOPIC EXAMINATION				
EPITHELIAL CELLS		1-2	0-5	/HPF
METHOD : MICROSCOPIC EXAMINATION				
CASTS		NOT DETECTED		
CRYSTALS		NOT DETECTED		
METHOD : MICROSCOPIC EXAMINATION				
BACTERIA		NOT DETECTED	NOT DETECTED	
METHOD : MICROSCOPIC EXAMINATION				
YEAST		NOT DETECTED	NOT DETECTED	



DIAGNOSTIC REPORT



Cert. No. MC-2351



CLIENT CODE : C000138383

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd
24 SCO, SECTOR 11 D
CHANDIGARH, 160011
PUNJAB, INDIA
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956

PATIENT NAME : RAM CHARAN

PATIENT ID : RAMCM24127080

ACCESSION NO : 0080WC011145 **AGE :** 52 Years **SEX :** Male

ABHA NO :

DRAWN : **RECEIVED :** 28/03/2023 08:54

REPORTED : 28/03/2023 15:29

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

Interpretation(s)

The following table describes the probable conditions, in which the analytes are present in urine

Presence of	Conditions
Proteins	Inflammation or immune illnesses
Pus (White Blood Cells)	Urinary tract infection, urinary tract or kidney stone, tumors or any kind of kidney impairment
Glucose	Diabetes or kidney disease
Ketones	Diabetic ketoacidosis (DKA), starvation or thirst
Urobilinogen	Liver disease such as hepatitis or cirrhosis
Blood	Renal or genital disorders/trauma
Bilirubin	Liver disease
Erythrocytes	Urological diseases (e.g. kidney and bladder cancer, urolithiasis), urinary tract infection and glomerular diseases
Leukocytes	Urinary tract infection, glomerulonephritis, interstitial nephritis either acute or chronic, polycystic kidney disease, urolithiasis, contamination by genital secretions
Epithelial cells	Urolithiasis, bladder carcinoma or hydronephrosis, ureteric stents or bladder catheters for prolonged periods of time
Granular Casts	Low intratubular pH, high urine osmolality and sodium concentration, interaction with Bence-Jones protein
Hyaline casts	Physical stress, fever, dehydration, acute congestive heart failure, renal diseases
Calcium oxalate	Metabolic stone disease, primary or secondary hyperoxaluria, intravenous infusion of large doses of vitamin C, the use of vasodilator naftidrofuryl oxalate or the gastrointestinal lipase inhibitor orlistat, ingestion of ethylene glycol or of star fruit (Averrhoa carambola) or its juice
Uric acid	arthritis
Bacteria	Urinary infection when present in significant numbers & with pus cells.
Trichomonas vaginalis	Vaginitis, cervicitis or salpingitis

THYROID PANEL, SERUM

T3	110.5	80.0 - 200.0	ng/dL
METHOD : ECLIA			
T4	8.94	5.10 - 14.10	µg/dL
METHOD : ECLIA			
TSH (ULTRASENSITIVE)	4.450	High 0.270 - 4.200	µIU/mL
METHOD : SANDWICH (ECLIA)			



Scan to View Details



Scan to View Report

DIAGNOSTIC REPORT



CLIENT CODE : C000138383

CLIENT'S NAME AND ADDRESS :
ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd
24 SCO, SECTOR 11 D
CHANDIGARH, 160011
PUNJAB, INDIA
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956

Cert. No. MC-2351

PATIENT NAME : RAM CHARAN

PATIENT ID : RAMCM24127080

ACCESSION NO : 0080WC011145 **AGE :** 52 Years **SEX :** Male

ABHA NO :

DRAWN : **RECEIVED :** 28/03/2023 08:54

REPORTED : 28/03/2023 15:29

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

Interpretation(s)

Triiodothyronine T3 , Thyroxine T4, and Thyroid Stimulating Hormone TSH are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate. Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH. Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hyperthyroidism, TSH levels are low. Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3. Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Sr. No.	TSH	Total T4	FT4	Total T3	Possible Conditions
1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3) Post Thyroidectomy (4) Post Radio-Iodine treatment
2	High	Normal	Normal	Normal	(1)Subclinical Hypothyroidism (2) Patient with insufficient thyroid hormone replacement therapy (3) In cases of Autoimmune/Hashimoto thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical inflammation, drugs like amphetamines, Iodine containing drug and dopamine antagonist e.g. domperidone and other physiological reasons.
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre (3)Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4 replacement therapy (7) First trimester of Pregnancy
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism
6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent treatment for Hyperthyroidism
8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies

REF: 1. TIETZ Fundamentals of Clinical chemistry 2.Guidlines of the American Thyroid association during pregnancy and Postpartum, 2011.

NOTE: It is advisable to detect Free T3,FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

PHYSICAL EXAMINATION,STOOL

COLOUR	BROWN
CONSISTENCY	SEMI FORMED
MUCUS	ABSENT NOT DETECTED
VISIBLE BLOOD	ABSENT ABSENT
ADULT PARASITE	NOT DETECTED

METHOD : MICROSCOPIC EXAMINATION



DIAGNOSTIC REPORT

Patient Ref. No. 80000001398215



Cert. No. MC-2351



CLIENT CODE : C000138383

CLIENT'S NAME AND ADDRESS :ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156SRL Ltd
24 SCO, SECTOR 11 D
CHANDIGARH, 160011
PUNJAB, INDIA
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956**PATIENT NAME : RAM CHARAN**PATIENT ID : **RAMCM24127080**ACCESSION NO : **0080WC011145** AGE : 52 Years SEX : Male

ABHA NO :

DRAWN :

RECEIVED : 28/03/2023 08:54

REPORTED : 28/03/2023 15:29

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

CHEMICAL EXAMINATION,STOOL

STOOL PH 6.5

MICROSCOPIC EXAMINATION,STOOL

PUS CELLS	1-2		/hpf
RED BLOOD CELLS	NOT DETECTED	NOT DETECTED	/HPF
CYSTS	NOT DETECTED	NOT DETECTED	
OVA	NOT DETECTED		
LARVAE	NOT DETECTED	NOT DETECTED	
TROPHOZOITES	NOT DETECTED	NOT DETECTED	



Scan to View Details



Scan to View Report



Patient Ref. No. 80000001398215



Cert. No. MC-2351



CLIENT CODE : C000138383

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd
24 SCO, SECTOR 11 D
CHANDIGARH, 160011
PUNJAB, INDIA
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956

PATIENT NAME : RAM CHARAN

PATIENT ID : RAMCM24127080

ACCESSION NO : 0080WC011145 AGE : 52 Years SEX : Male

ABHA NO :

DRAWN :

RECEIVED : 28/03/2023 08:54

REPORTED : 28/03/2023 15:29

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

Interpretation(s)

Stool routine analysis is only a screening test for disorders of gastrointestinal tract like infection, malabsorption, etc. The following table describes the probable conditions, in which the analytes are present in stool.

PRESENCE OF	CONDITION
Pus cells	Pus in the stool is an indication of infection
Red Blood cells	Parasitic or bacterial infection or an inflammatory bowel condition such as ulcerative colitis
Parasites	Infection of the digestive system. Stool examination for ova and parasite detects presence of parasitic infestation of gastrointestinal tract. Various forms of parasite that can be detected include cyst, trophozoite and larvae. One negative result does not rule out the possibility of parasitic infestation. Intermittent shedding of parasites warrants examinations of multiple specimens tested on consecutive days. Stool specimens for parasitic examination should be collected before initiation of anti-diarrheal therapy or antiparasitic therapy. This test does not detect presence of opportunistic parasites like Cyclospora, Cryptosporidia and Isospora species. Examination of Ova and Parasite has been carried out by direct and concentration techniques.
Mucus	Mucus is a protective layer that lubricates, protects & reduces damage due to bacteria or viruses.
Charcot-Leyden crystal	Parasitic diseases.
Ova & cyst	Ova & cyst indicate parasitic infestation of intestine.
Frank blood	Bleeding in the rectum or colon.
Occult blood	Occult blood indicates upper GI bleeding.
Macrophages	Macrophages in stool are an indication of infection as they are protective cells.
Epithelial cells	Epithelial cells that normally line the body surface and internal organs show up in stool when there is inflammation or infection.
Fat	Increased fat in stool maybe seen in conditions like diarrhoea or malabsorption.
pH	Normal stool pH is slightly acidic to neutral. Breast-fed babies generally have an acidic stool.

ADDITIONAL STOOL TESTS :

- Stool Culture**:- This test is done to find cause of GI infection, make decision about best treatment for GI infection & to find out if treatment for GI infection worked.
- Fecal Calprotectin**: It is a marker of intestinal inflammation. This test is done to differentiate Inflammatory Bowel Disease (IBD) from Irritable Bowel Syndrome (IBS).
- Fecal Occult Blood Test (FOBT)**: This test is done to screen for colon cancer & to evaluate possible cause of unexplained anaemia.
- Clostridium Difficile Toxin Assay**: This test is strongly recommended in healthcare associated bloody or watery diarrhoea, due to overuse of broad spectrum antibiotics which alter the normal GI flora.
- Biofire (Film Array) GI PANEL**: In patients of Diarrhoea, Dysentery, Rice watery Stool, FDA approved, Biofire Film Array Test, (Real Time Multiplex PCR) is strongly recommended as it identifies organisms, bacteria, fungi, virus, parasite and other opportunistic pathogens, Vibrio cholera infections only in 3 hours. Sensitivity 96% & Specificity 99%.
- Rota Virus Immunoassay**: This test is recommended in severe gastroenteritis in infants & children associated with watery diarrhoea, vomiting & abdominal cramps. Adults are also affected. It is highly contagious in nature.



Scan to View Details



Scan to View Report



Patient Ref. No. 8000001398215



Cert. No. MC-2351



CLIENT CODE : C000138383

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd
24 SCO, SECTOR 11 D
CHANDIGARH, 160011
PUNJAB, INDIA
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956

PATIENT NAME : RAM CHARAN

PATIENT ID : RAMCM24127080

ACCESSION NO : 0080WC011145 AGE : 52 Years SEX : Male

ABHA NO :

DRAWN : RECEIVED : 28/03/2023 08:54

REPORTED : 28/03/2023 15:29

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

ABO GROUP TYPE B

METHOD : SLIDE AGGLUTINATION

RH TYPE POSITIVE

METHOD : SLIDE AGGLUTINATION

Interpretation(s)

BLOOD COUNTS, EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients A.-P. Yang, et al. International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope.

ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD-TEST DESCRIPTION :-

Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition. CRP is superior to ESR because it is more sensitive and reflects a more rapid change.

TEST INTERPRETATION

Increase in: Infections, Vasculitis, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy, Estrogen medication, Aging.

Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias, Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis).

In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm/hr(95 if anemic). ESR returns to normal 4th week post partum.

Decreased in: Polycythemia vera, Sickle cell anemia

LIMITATIONS

False elevated ESR : Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia

False Decreased : Poikilocytosis, (Sickle Cells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs(Quinine,

salicylates)

REFERENCE :

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th edition.

GLYCOSYLATED HEMOGLOBIN(HbA1C), EDTA WHOLE BLOOD-Used For:

1. Evaluating the long-term control of blood glucose concentrations in diabetic patients.

2. Diagnosing diabetes.

3. Identifying patients at increased risk for diabetes (prediabetes).

The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for well-controlled type 2 diabetic patients) to determine whether a patient's metabolic control has remained continuously within the target range.

1. eAG (Estimated average glucose) converts percentage HbA1c to mg/dl, to compare blood glucose levels.

2. eAG gives an evaluation of blood glucose levels for the last couple of months.

3. eAG is calculated as eAG (mg/dl) = 28.7 * HbA1c - 46.7

HbA1c Estimation can get affected due to :



Scan to View Details



Scan to View Report



Cert. No. MC-2351

CLIENT CODE : C000138383

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd
24 SCO, SECTOR 11 D
CHANDIGARH, 160011
PUNJAB, INDIA
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956

PATIENT NAME : RAM CHARAN

PATIENT ID : RAMCM24127080

ACCESSION NO : 0080WC011145 AGE : 52 Years SEX : Male

ABHA NO :

DRAWN : RECEIVED : 28/03/2023 08:54

REPORTED : 28/03/2023 15:29

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

1. Shortened Erythrocyte survival : Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss, hemolytic anemia) will falsely lower HbA1c test results. Fructosamine is recommended in these patients which indicates diabetes control over 15 days.

2. Vitamin C & E are reported to falsely lower test results. (possibly by inhibiting glycation of hemoglobin).

3. Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates addition are reported to interfere with some assay methods, falsely increasing results.

4. Interference of hemoglobinopathies in HbA1c estimation is seen in

a) Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c.
b) Heterozygous state detected (D10 is corrected for HbS & HbC trait.)
c) HbF > 25% on alternate platform (Boronate affinity chromatography) is recommended for testing of HbA1c. Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy

GLUCOSE FASTING, FLUORIDE PLASMA-TEST DESCRIPTION
Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and so that no glucose is excreted in the urine.

Increased in: Diabetes mellitus, Cushing's syndrome (10 - 15%), chronic pancreatitis (30%). Drugs: corticosteroids, phenytoin, estrogen, thiazides.
Decreased in: Pancreatic islet cell disease with increased insulin, insulinoma, adrenocortical insufficiency, hypopituitarism, diffuse liver disease, malignancy (adrenocortical, stomach, fibrosarcoma), infant of a diabetic mother, enzyme deficiency diseases (e.g. galactosemia), Drugs-insulin, ethanol, propranolol, sulfonylureas, tolbutamide, and other oral hypoglycemic agents.

NOTE: While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values), there is wide fluctuation within individuals. Thus, glycosylated hemoglobin (HbA1c) levels are favored to monitor glycemic control.
High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glycosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.
GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glycosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc. Additional test HbA1c

LIVER FUNCTION PROFILE, SERUM-
Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. **Elevated levels** result from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors & Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health. AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis.

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Paget's disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilson's disease.

GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc.

Total Protein also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease. Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

Albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc

BLOOD UREA NITROGEN (BUN), SERUM- Causes of Increased levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)

Causes of decreased level include Liver disease, SIADH.

CREATININE, SERUM- Higher than normal level may be due to:

- Blockage in the urinary tract, Kidney problems, such as kidney damage or failure, infection, or reduced blood flow, Loss of body fluid (dehydration), Muscle problems, such as breakdown of muscle fibers, Problems during pregnancy, such as seizures (eclampsia), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:

- Myasthenia Gravis, Muscular atrophy

URIC ACID, SERUM- Causes of Increased levels: Dietary (High Protein Intake, Prolonged Fasting, Rapid weight loss), Gout, Lesch nyhan syndrome, Type 2 DM, Metabolic syndrome

Causes of decreased levels: Low Zinc intake, OCP, Multiple Sclerosis

TOTAL PROTEIN, SERUM- is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin.

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease.

Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

ALBUMIN, SERUM-



DIAGNOSTIC REPORT

Patient Ref. No. 80000001398215



Cert. No. MC-2351



CLIENT CODE : C000138383

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd
24 SCO, SECTOR 11 D
CHANDIGARH, 160011
PUNJAB, INDIA
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956

PATIENT NAME : RAM CHARANPATIENT ID : **RAMCM24127080**ACCESSION NO : **0080WC011145** AGE : 52 Years SEX : Male

ABHA NO :

DRAWN : RECEIVED : 28/03/2023 08:54

REPORTED : 28/03/2023 15:29

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. **Low blood albumin levels (hypoalbuminemia) can be caused by:** Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.
ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same."

The test is performed by both forward as well as reverse grouping methods.

****End Of Report****

Please visit www.srlworld.com for related Test Information for this accession
TEST MARKED WITH '*' ARE OUTSIDE THE NABL ACCREDITED SCOPE OF THE LABORATORY.

Dr. Pranjali Vasisht
LAB HEAD

Dr. Nidhi Garg
Lab Consultant

DR. CHANDNI GARG
CONSULTANT PATHOLOGIST

CONDITIONS OF LABORATORY TESTING & REPORTING

1. It is presumed that the test sample belongs to the patient named or identified in the test requisition form.
2. All tests are performed and reported as per the turnaround time stated in the SRL Directory of Services.
3. Result delays could occur due to unforeseen circumstances such as non-availability of kits / equipment breakdown / natural calamities / technical downtime or any other unforeseen event.
4. A requested test might not be performed if:
 - i. Specimen received is insufficient or inappropriate
 - ii. Specimen quality is unsatisfactory
 - iii. Incorrect specimen type
 - iv. Discrepancy between identification on specimen container label and test requisition form
5. SRL confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity.
6. Laboratory results should not be interpreted in isolation; it must be correlated with clinical information and be interpreted by registered medical practitioners only to determine final diagnosis.
7. Test results may vary based on time of collection, physiological condition of the patient, current medication or nutritional and dietary changes. Please consult your doctor or call us for any clarification.
8. Test results cannot be used for Medico legal purposes.
9. In case of queries please call customer care (91115 91115) within 48 hours of the report.

SRL Limited

Fortis Hospital, Sector 62, Phase VIII,
Mohali 160062



Scan to View Details

Page 15 Of 15



Scan to View Report