SID No.

Collection On : 26/03/2022 7:57 AM

Report On : 27/03/2022 3:50 PM

 Age / Sex
 : 44 Year(s) / Female
 Report On
 : 27/03/2022 3:50 PM

 Type
 : OP
 Printed On
 : 01/04/2022 6:49 AM

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HAEMATOLOGY			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood'Spectrophotometry)	12.2	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood/Derived from Impedance)	37.2	%	37 - 47
RBC Count (EDTA Blood/Impedance Variation)	4.38	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (EDTA Blood/Derived from Impedance)	85.0	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood/Derived from Impedance)	27.9	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood/Derived from Impedance)	32.8	g/dL	32 - 36
RDW-CV (EDTA Blood/Derived from Impedance)	12.7	%	11.5 - 16.0
RDW-SD (EDTA Blood/Derived from Impedance)	37.78	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood/Impedance Variation)	6500	cells/cu.mm	4000 - 11000
Neutrophils (EDTA Blood/Impedance Variation & Flow Cytometry)	52.9	%	40 - 75
Lymphocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	37.1	%	20 - 45





 PID No.
 : MED111034201
 Register On
 : 26/03/2022 7:01 AM

 SID No.
 : 422024456
 Collection On
 : 26/03/2022 7:57 AM

Printed On

: 01/04/2022 6:49 AM

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Eosinophils (EDTA Blood/Impedance Variation & Flow Cytometry)	1.7	%	01 - 06
Monocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	7.6	%	01 - 10
Basophils (Blood/Impedance Variation & Flow Cytometry)	0.7	%	00 - 02
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	3.44	10^3 / µl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	2.41	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.11	10^3 / µl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.49	10^3 / µl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.05	10^3 / µl	< 0.2
Platelet Count (EDTA Blood/Impedance Variation)	289	10^3 / μl	150 - 450
MPV (EDTA Blood/Derived from Impedance)	8.2	fL	8.0 - 13.3
PCT (EDTA Blood/Automated Blood cell Counter)	0.24	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood/Modified Westergren)	21	mm/hr	< 20





 PID No.
 : MED111034201
 Register On
 : 26/03/2022 7:01 AM

 SID No.
 : 422024456
 Collection On
 : 26/03/2022 7:57 AM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
BIOCHEMISTRY			
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulfanilic Acid)	0.4	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.1	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.3	mg/dL	0.1 - 1.0
Total Protein (Serum/Biuret)	7.0	gm/dL	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.3	gm/dL	3.5 - 5.2
Globulin (Serum/Derived)	2.7	gm/dL	2.3 - 3.6
A: G Ratio (Serum/Derived)	1.6		1.1 - 2.2
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC Kinetic)	15	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC / Kinetic)	13	U/L	5 - 41
Alkaline Phosphatase (SAP) (Serum/IFCC Kinetic)	88	U/L	42 - 98
GGT(Gamma Glutamyl Transpeptidase) (Serum/SZASZ standarised IFCC)	18	U/L	< 38

: 01/04/2022 6:49 AM



VERIFIED BY



 PID No.
 : MED111034201
 Register On
 : 26/03/2022 7:01 AM

 SID No.
 : 422024456
 Collection On
 : 26/03/2022 7:57 AM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
<u>Lipid Profile</u>			
Cholesterol Total (Serum/Cholesterol oxidase/Peroxidase)	220	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol phosphate oxidase / peroxidase)	145	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

: 01/04/2022 6:49 AM

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the "usual" circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	55	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/Calculated)	136	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	29	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	165.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220



VERIFIED BY



PID No. : MED111034201 Register On : 26/03/2022 7:01 AM : 422024456 Collection On : 26/03/2022 7:57 AM SID No.

Age / Sex : 44 Year(s) / Female Report On : 27/03/2022 3:50 PM

Printed On

Type : OP

Ref. Dr

: MediWheel



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	Value		Reference Interval

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

: 01/04/2022 6:49 AM

Total Cholesterol/HDL Cholesterol Ratio	4	Optimal: < 3.3
(Serum/Calculated)		Low Risk: 3.4 - 4.4
		Average Risk: 4.5 - 7.1
		Moderate Risk: 7.2 - 11.0
		High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio	2.6	Optimal: < 2.5
(TG/HDL)		Mild to moderate risk: 2.5 - 5.0
(Serum/Calculated)		High Risk: > 5.0

(Serum/Calculated)

LDL/HDL Cholesterol Ratio 2.5 Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 (Serum/Calculated) High Risk: > 6.0



VERIFIED BY



PID No. : MED111034201 Register On : 26/03/2022 7:01 AM

Printed On

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/HPLC)	6.2	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

: 01/04/2022 6:49 AM

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 131.24 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.



VERIFIED BY



Printed On

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	Value		Reference Interval

: 01/04/2022 6:49 AM

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.32 ng/mL 0.7 - 2.04

(Serum/CMIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is

Metabolically active.

T4 (Thyroxine) - Total 9.44 μg/dL 4.2 - 12.0

(Serum/CMIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 3.10 µIU/mL 0.35 - 5.50

(Serum/Chemiluminescent Microparticle

Immunoassay(CMIA))

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values&lt,0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



PID No. : MED111034201 Register On : 26/03/2022 7:01 AM : 422024456 Collection On : 26/03/2022 7:57 AM SID No.

Report On : 27/03/2022 3:50 PM

Type : OP

Ref. Dr

Age / Sex : 44 Year(s) / Female

: MediWheel

Printed On : 01/04/2022 6:49 AM



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
CLINICAL PATHOLOGY			
PHYSICAL EXAMINATION			
Colour (Urine)	Pale Yellow		
Appearance (Urine)	Clear		Clear
Volume (Urine)	20	mL	
<u>CHEMICAL EXAMINATION(Automated-Urineanalyser)</u>			
pH (Urine/AUTOMATED URINANALYSER)	5.0		4.5 - 8.0
Specific Gravity (Urine)	1.020		1.002 - 1.035
Ketones (Urine)	Negative		Negative
Urobilinogen (Urine/AUTOMATED URINANALYSER)	0.2		0.2 - 1.0
Blood (Urine/AUTOMATED URINANALYSER)	Negative		Negative
Nitrite (Urine/AUTOMATED URINANALYSER)	Negative		Negative
Bilirubin (Urine/AUTOMATED URINANALYSER)	Negative		Negative

Negative



Protein (Urine)

VERIFIED BY



Negative

 PID No.
 : MED111034201
 Register On
 : 26/03/2022 7:01 AM

 SID No.
 : 422024456
 Collection On
 : 26/03/2022 7:57 AM

: 01/04/2022 6:49 AM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glucose (Urine)	Negative		Negative
Leukocytes (Urine)	Negative	leuco/uL	Negative
MICROSCOPY(URINE DEPOSITS)			
Pus Cells (Urine/Flow cytometry)	1-2	/hpf	3-5
Epithelial Cells (Urine)	1-2	/hpf	1-2
RBCs (Urine/Flow cytometry)	Nil	/hpf	NIL
Others (Urine)	Nil		Nil
Casts (Urine/Flow cytometry)	Nil	/hpf	0 - 1
Crystals (Urine)	Nil		NIL



VERIFIED BY



PID No. : MED111034201 Reg

SID No. : 422024456

Age / Sex : 44 Year(s) / Female

Type : OP

Ref. Dr : MediWheel

Register On : 26/03/2022 7:01 AM

Collection On : 26/03/2022 7:57 AM

Report On : 27/03/2022 3:50 PM

Printed On : 01/04/2022 6:49 AM

MEDALI

InvestigationObservedUnitBiologicalValueReference Interval

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING 'AB' 'Positive'

(EDTA Blood/Agglutination)





 PID No.
 : MED111034201
 Register On
 : 26/03/2022 7:01 AM

 SID No.
 : 422024456
 Collection On
 : 26/03/2022 7:57 AM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	17		6 - 22
Glucose Fasting (FBS) (Plasma - F/GOD - POD)	94	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

: 01/04/2022 6:49 AM

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose Fasting - Urine (Urine - F)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD - POD)	113	mg/dL	70 - 140

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Glucose Postprandial - Urine (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease-GLDH)	7	mg/dL	7.0 - 21
Creatinine (Serum/Jaffe Kinetic)	0.6	mg/dL	0.6 - 1.1

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine etc.

Uric Acid 5.0 mg/dL 2.6 - 6.0

(Serum/Uricase/Peroxidase)



VERIFIED BY



APPROVED BY

-- End of Report --

Name	Symala	ID	MED111034201
Age & Gender	44/FeMale		26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



2 D ECHOCARDIOGRAPHIC STUDY

M mode measurement:

AORTA : 2.3cms

LEFT ATRIUM : 2.9cms

AVS :----

LEFT VENTRICLE (DIASTOLE) : 4.8cms

(SYSTOLE) : 3.2cms

VENTRICULAR SEPTUM (DIASTOLE) : 0.9cms

(SYSTOLE) : 1.5cms

POSTERIOR WALL (DIASTOLE) : 0.9cms

(SYSTOLE) : 1.2cms

EDV : 107ml

ESV : 40ml

FRACTIONAL SHORTENING : 34%

EJECTION FRACTION : 63%

EPSS :---

RVID : 1.6cms

DOPPLER MEASUREMENTS:

MITRAL VALVE : E' 0.76 m/s A' 0.94 m/s NO MR

AORTIC VALVE : 1.24 m/s NO AR

TRICUSPID VALVE : E' 1.88 m/s A' - m/s NO TR

PULMONARY VALVE : 0.98 m/s NO PR

Name	Symala	ID	MED111034201
Age & Gender	44/FeMale	Visit Date	26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



2D ECHOCARDIOGRAPHY FINDINGS:

Left ventricle : Normal size, Normal systolic function.

No regional wall motion abnormalities.

Left Atrium : Normal.

Right Ventricle : Normal.

Right Atrium : Normal.

Mitral valve : Normal, No mitral valve prolapsed.

Aortic valve : Normal, Trileaflet.

Tricuspid valve : Normal.

Pulmonary valve : Normal.

IAS : Intact.

IVS : Intact.

Pericardium : No pericardial effusion.

IMPRESSION:

- > LV DIASTOLIC DYSFUNCTION.
- > NORMAL SIZED CARDIAC CHAMBERS.
- > NORMAL LV SYSTOLIC FUNCTION. EF:63 %.
- > NO REGIONAL WALL MOTION ABNORMALITIES.
- > NORMAL VALVES.
- > NO CLOTS / PERICARDIAL EFFUSION / VEGETATION.

DR. K.S. SUBRAMANI. MBBS, MD, DM (CARDIOLOGY) FESC SENIOR CONSULTANT INTERVENTIONAL CARDIOLOGIST

Name	Symala	ID	MED111034201
Age & Gender	44/FeMale	Visit Date	26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



Kss/da

- Note:

 * Report to be interpreted by qualified medical professional.

 * To be correlated with other clinical findings.

 * Parameters may be subjected to inter and intra observer variations.

Name	Symala	ID	MED111034201
Age & Gender	44/FeMale		26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER shows normal in shape, size and has uniform echopattern. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER shows normal shape and has clear contents. Gall bladder wall is of normal thickness. CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN shows normal shape, size and echopattern. Spleen measures 7.5cms in long axis and 3.3cms in short axis.

No demonstrable Para -aortic lymphadenopathy.

KIDNEYS move well with respiration and have normal shape, size and echopattern. Cortico- medullary differentiations are well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

·	Bipolar length (cms) Parenchymal thickness (cms)	
Right Kidney	11.1	1.3
Left Kidney	10.8	1.2

URINARY BLADDER shows normal shape and wall thickness. It has clear contents. No evidence of diverticula.

UTERUS is anteverted and mildly retroflexed and is slightly bulky. It has uniform myometrial echopattern.

No focal myometrial lesion is noted. Endometrial thickness measures 8mm and is triple line. Uterus measures as follows: LS: 9.2cms AP: 4.8cms TS: 5.5cms.

OVARIES are normal in size, shape and echotexture. No focal lesion seen.

Right ovary shows a large follicle measuring about 18 x 17mm.

Ovaries measure as follows: **Right ovary**: 3.4 x 1.8cms **Left ovary**: 2.6 x 1.4cms

POD & adnexae are free.

No evidence of ascites/pleural effusion.

IMPRESSION:

> NO SIGNIFICANT ABNORMALITY.

Name	Symala	ID	MED111034201
Age & Gender	44/FeMale		26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



DR. H.K. ANAND CONSULTANT RADIOLOGISTS MS/vp DR. MEERA S

Name	Symala	ID	MED111034201
Age & Gender	44/FeMale	Visit Date	26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



X-ray mammogram (mediolateral oblique and craniocaudal views) followed by Sonomammography was performed.

MAMMOGRAPHY OF BOTH BREASTS

Both breasts show symmetrical relatively dense fibro glandular fatty tissue.

No evidence of focal soft tissue lesion.

No evidence of cluster micro calcification.

Subcutaneous fat deposition is within normal limits.

SONOMAMMOGRAPHY OF BOTH BREASTS

Both breasts show diffuse fibrocystic changes.

No evidence of focal solid / cystic areas in either breast.

Retroareolar region is normal on either side.

Few lymphnodes with maintained fatty hilum are noted in both axillae.

IMPRESSION:

> BILATERAL FIBROCYSTIC CHANGES.

ASSESSMENT: BI-RADS CATEGORY -2

DR. H.K. ANAND CONSULTANT RADIOLOGISTS MS/vp DR. MEERA S

BI-RADS CLASSIFICATION

CATEGORY	RESULT
0	Assessment incomplete. Need additional imaging evaluation
1	Negative. Routine mammogram in 1 year recommended.
2	Benign finding. Routine mammogram in 1 year recommended.
3	Probably benign finding. Short interval follow-up suggested.
4	Suspicious. Biopsy should be considered.
5	Highly suggestive of malignancy. Appropriate action should be taken.

Name	Symala	ID	MED111034201
Age & Gender	44/FeMale	Visit Date	26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



Name	Symala	ID	MED111034201
Age & Gender	44/FeMale		26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

IMPRESSION:

No significant abnormality detected.