

Patient Name : Mr Satapathy Lalit Kumar

DOB/Age/Gender : 56 Y/Male Sample Collected : Apr 04, 2024, 10:00 PM

Patient ID / UHID : 1_7835348/RCL7008496 Report Date : Apr 04, 2024, 08:08 PM

Referred By : Self Barcode No : HY408330
Sample Type : Whole blood EDTA Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range



Hemogram (CBC + ESR)

Complete Blood Count (CBC)

RBC Parameters			
Hemoglobin	14.7	g/dL	13.0 - 17.0
colorimetric			
RBC Count	5.2	10^6/µl	4.5 - 5.5
Electrical impedance			
PCV Calculated	46	%	40 - 50
Calculated MCV	88.3	fl	83 - 101
Calculated	88.3	"	83 - 101
MCH	28.2	pg	27 - 32
Calculated	20.2	l P9	21 02
MCHC	31.9	g/dL	31.5 - 34.5
Calculated			
RDW (CV) *	13.1	%	11.6 - 14.0
Calculated			
RDW-SD *	46.2	fl	35.1 - 43.9
Calculated			
WBC Parameters		1010/	
TLC	5.6	10^3/µl	4 - 10
Electrical impedance and microscopy Differential Leucocyte Count			
<u>-</u>		0/	40.00
Neutrophils	56	%	40-80
Lymphocytes	36	%	20-40
Monocytes	5	%	2-10
Eosinophils	3	%	1-6
Basophils	0	%	<2
Absolute Leukocyte Counts Calculated			
Neutrophils.	3.14	10^3/µl	2 - 7
Lymphocytes.	2.02	10^3/µl	1 - 3
Monocytes.	0.28	10^3/µl	0.2 - 1.0
Eosinophils.	0.17	10^3/µl	0.02 - 0.5
Basophils.	0	10^3/µl	0.02 - 0.5
Platelet Parameters		10 0/μ.	0.02 0.0
Platelet Count	165	10^3/µl	150 - 410
Electrical impedance and microscopy		, s s, p	
Mean Platelet Volume (MPV) *	12.8	fL	9.3 - 12.1
Calculated			
PCT *	0.2	%	0.17 - 0.32
Calculated			

^(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.

Dr. Pallavi Rath MBBS, MD (Pathology) Consultant Pathologist

Pallari





Patient Name : Mr Satapathy Lalit Kumar

DOB/Age/Gender : 56 Y/Male Sample Collected : Apr 04, 2024, 10:00 PM

Patient ID / UHID : 1_7835348/RCL7008496 Report Date : Apr 04, 2024, 08:08 PM

Referred By : Self Barcode No : HY408330
Sample Type : Whole blood EDTA Report Status : Final Report

MABL M(EL)T LABS

Test Description	Value(s)	Unit(s)	Reference Range
PDW *	25.4	fL	8.3 - 25.0
Calculated			
P-LCR *	52.6	%	18 - 50
Calculated			
P-LCC *	87	%	44 - 140
Calculated			
Mentzer Index *	16.98	%	-
Calculated			

Interpretation:

CBC provides information about red cells, white cells and platelets. Results are useful in the diagnosis of anemia, infections, leukemias, clotting disorders and many other medical conditions.

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.

Dr. Pallavi Rath MBBS, MD (Pathology) Consultant Pathologist

Pallari





Patient Name : Mr Satapathy Lalit Kumar

DOB/Age/Gender : 56 Y/Male Sample Collected : Apr 04, 2024, 10:00 PM Patient ID / UHID : 1_7835348/RCL7008496 Report Date : Apr 04, 2024, 08:13 PM

Referred By : Self Barcode No : HY408330
Sample Type : Whole blood EDTA Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range

Erythrocyte Sedimentation Rate (ESR)

ESR - Erythrocyte Sedimentation Rate	8	mm/hr	0 - 12
MODIFIED WESTERGREN			

Interpretation:

ESR is also known as Erythrocyte Sedimentation Rate. An ESR test is used to assess inflammation in the body. Many conditions can cause an abnormal ESR, so an ESR test is typically used with other tests to diagnose and monitor different diseases. An elevated ESR may occur in inflammatory conditions including infection, rheumatoid arthritis, systemic vasculitis, anemia, multiple myeloma, etc. Low levels are typically seen in congestive heart failure, polycythemia, sickle cell anemia, hypo fibrinogenemia, etc.

AGE	MALE	FEMALE
1 DAY	0-2	0-2
2 - 7 DAYS	0-4	0-4
8 - 14 DAYS	0-17	0-17
15 DAYS - 17 YEARS	0-20	0-20
18 - 50 YEARS	0-10	0-12
51- 60 YEARS	0-12	0-19
61 - 70 YEARS	0-14	0-20
71 - 100 YEARS	0-30	0-35

Reference- Dacie and lewis practical hematology

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



Pallari





Patient Name : Mr Satapathy Lalit Kumar

DOB/Age/Gender : 56 Y/Male Sample Collected : Apr 04, 2024, 10:00 PM

Patient ID / UHID : 1_7835348/RCL7008496 Report Date : Apr 04, 2024, 08:00 PM

Referred By : Self Barcode No : HY408330
Sample Type : Whole blood EDTA Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range



HbA1C (Glycosylated Haemoglobin)

Glycosylated Hemoglobin (HbA1c) HPLC	5.5	%	<5.7
Estimated Average Glucose *	111.15	mg/dL	Refer Table Below

Interpretation:

Interpretation For HbA1c% As per American Diabetes Association (ADA)

Reference Group	HbA1c in %
Non diabetic adults >=18 years	<5.7
At risk (Prediabetes)	5.7 - 6.4
Diagnosing Diabetes	>= 6.5
Therapeutic goals for glycemic control	Age > 19 years Goal of therapy: < 7.0 Age < 19 years Goal of therapy: <7.5

Note:

- 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.
- 2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

Comments:

HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations ADA criteria for correlation between HbA1c & Mean plasma glucose levels.

HbA1c(%)	Mean Plasma Glucose (mg/dL)	HbA1c(%)	Mean Plasma Glucose (mg/dL)
6	126	12	298
8	183	14	355
10	240	16	413

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.

Dr. Pallavi Rath MBBS, MD (Pathology) Consultant Pathologist

Pallari





Patient Name : Mr Satapathy Lalit Kumar

 DOB/Age/Gender
 : 56 Y/Male
 Sample Collected
 : Apr 04, 2024, 10:00 PM

 Patient ID / UHID
 : 1_7835348/RCL7008496
 Report Date
 : Apr 04, 2024, 08:26 PM


Referred By : Self Barcode No : HY408330
Sample Type : Whole blood EDTA Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range

Blood Group ABO & Rh Typing

Blood Group	В	-	-
Rh Factor	Positive	-	-

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



Paulari





Patient Name : Mr Satapathy Lalit Kumar

DOB/Age/Gender : 56 Y/Male Sample Collected : Apr 04, 2024, 10:00 PM

Patient ID / UHID : 1_7835348/RCL7008496 Report Date : Apr 04, 2024, 07:46 PM

Referred By : Self Barcode No : ZC244583
Sample Type : FLUORIDE F Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range



Glucose Fasting (BSF)

Glucose Fasting	73	mg/dL	70 - 100
Hexokinase			

Interpretation:

Status	Fasting plasma glucose in mg/dL	
Normal	<100	
Impaired fasting glucose	100 - 125	
Diabetes	=>126	

Reference: American Diabetes Association

Comment:

Blood glucose determinations in commonly used as an aid in the diagnosis and treatment of diabetes. Elevated glucose levels (hyperglycemia) may also occur with pancreatic neoplasm, hyperthyroidism, and adrenal cortical hyper function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy insulinoma, or various liver diseases.

Note

- 1. The diagnosis of Diabetes requires a fasting plasma glucose of > or = 126 mg/dL or a random / 2 hour plasma glucose value of > or = 200 mg/dL with symptoms of diabetes mellitus.
- 2. Very high glucose levels (>450 mg/dL in adults) may result in Diabetic Ketoacidosis.

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.

Dr. Pallavi Rath MBBS, MD (Pathology) Consultant Pathologist

Pallari





Patient Name : Mr Satapathy Lalit Kumar

DOB/Age/Gender : 56 Y/Male Sample Collected : Apr 04, 2024, 10:00 PM

Patient ID / UHID : 1_7835348/RCL7008496 Report Date : Apr 04, 2024, 07:46 PM

Referred By : Self Barcode No : ZC244584
Sample Type : Serum Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range



Liver Function Test (LFT)

Bilirubin Total Photometric	0.6	mg/dL	0.2 - 1.2
Bilirubin Direct * Diazo Reaction	0.2	mg/dL	0.0 - 0.5
Bilirubin Indirect * Calculation (T Bil - D Bil)	0.4	mg/dL	0.1 - 1.0
SGOT/AST IFCC without P5P	25	U/L	5 - 34
SGPT/ALT IFCC without P5P	24	U/L	0 to 55
SGOT/SGPT Ratio *	1.04	-	-
Alkaline Phosphatase IFCC	55	U/L	40 - 150
Total Protein Biuret	8	g/dL	6.4 - 8.3
Albumin BCG	4.9	gm/dL	3.8 - 5.0
Globulin * Calculation (T.P - Albumin)	3.1	g/dL	2.3 - 3.5
Albumin :Globulin Ratio * Calculation (Albumin/Globulin)	1.58	-	1.0 - 2.1
Gamma Glutamyl Transferase (GGT) * Photometric	23	U/L	12 - 64

Interpretation:

The liver filters and processes blood as it circulates through the body. It metabolizes nutrients, detoxifies harmful substances, makes blood clotting proteins, and performs many other vital functions. The cells in the liver contain proteins called enzymes that drive these chemical reactions. When liver cells are damaged or destroyed, the enzymes in the cells leak out into the blood, where they can be measured by blood tests Liver tests check the blood for two main liver enzymes. Aspartate aminotransferase (AST),SGOT: The AST enzyme is also found in muscles and many other tissues besides the liver. Alanine aminotransferase (ALT), SGPT: ALT is almost exclusively found in the liver. If ALT and AST are found together in elevated amounts in the blood, liver damage is most likely present. Alkaline Phosphatase and GGT: Another of the liver's key functions is the production of bile, which helps digest fat. Bile flows through the liver in a system of small tubes (ducts), and is eventually stored in the gallbladder, under the liver. When bile flow is slow or blocked, blood levels of certain liver enzymes rise: Alkaline phosphatase Gamma-utamyl transpeptidase (GGT) Liver tests may check for any or all of these enzymes in the blood. Alkaline phosphatase is by far the most commonly tested of the three. If alkaline phosphatase and GGT are elevated, a problem with bile flow is most likely present. Bile flow problems can be due to a problem in the liver, the gallbladder, or the tubes connecting them. Proteins are important building blocks of all cells and tissues. Proteins are necessary for your body's growth, development, and health. Blood contains two classes of protein, albumin and globulin. Albumin proteins keep fluid from leaking out of blood vessels. Globulin proteins play an important role in your immune system. Low total protein may

Indicate:

- 1.Bleeding
- 2.Liver disorder
- 3.Malnutrition
- 4.Agammaglobulinemia High Protein levels 'Hyperproteinemia: May be seen in dehydration due to inadequate water intake or to excessive water loss (eg, severe vomiting, diarrhea, Addison's disease and diabetic acidosis) or as a result of increased production of proteins Low albumin levels may be

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.

Dr. Pallavi Rath MBBS, MD (Pathology) Consultant Pathologist

Paulari





NABL

M(EL)T

LABS

NABL-M(EL)T-00609

Patient Name : Mr Satapathy Lalit Kumar

DOB/Age/Gender : 56 Y/Male Sample Collected : Apr 04, 2024, 10:00 PM

Patient ID / UHID : 1_7835348/RCL7008496 Report Date : Apr 04, 2024, 07:46 PM

Referred By : Self Barcode No : ZC244584
Sample Type : Serum Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range

Caused by:

1.A poor diet (malnutrition).

2.Kidney disease.

3. Liver disease. High albumin levels may be caused by: Severe dehydration.

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.

Dr. Pallavi Rath MBBS, MD (Pathology) Consultant Pathologist

Paulari



Booking Centre :- Madyosis Diagnostics, Office No-406, 4th Floor, Bhakti Genesis, Wakad Rd, Shedge Vasti, Shankar Kalat Nagar, Wakad, Pimpri-Chinchwad, Maharashtra 411057

Processing Lab: Redcliffe Lifetech Pvt. Ltd., First Floor, B Wing. Aswani Chambers, S.No. 199+204+205 206/1, 209/1, Plot No. 45/B. Corresponding city/S No. 199 Village Lohgron Bune 411014, www.redcliffelabs.com



Patient Name : Mr Satapathy Lalit Kumar

DOB/Age/Gender : 56 Y/Male Sample Collected : Apr 04, 2024, 10:00 PM

Patient ID / UHID : 1_7835348/RCL7008496 Report Date : Apr 04, 2024, 07:46 PM

Referred By : Self Barcode No : ZC244584
Sample Type : Serum Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range



Kidney Function Test (KFT)

Blood Urea Urease	19	mg/dL	18 - 55
Creatinine Photometric	0.77	mg/dL	0.72 - 1.25
Bun * Urease	8.88	mg/dL	8.4 - 25.7
Bun/Creatinine Ratio *	11.53		
Urea / Creatinine Ratio *	24.68		
Uric Acid Uricase	4.9	mg/dL	3.5 - 7.2
Calcium Serum Arsenazo III	10.1	mg/dL	8.4 - 10.2
Phosphorus Photometric	3.9	mg/dL	2.3 - 4.7
Sodium Potentiometric	141	mmol/L	136 - 145
Potassium Potentiometric	4.2	mmol/L	3.5 - 5.1
Chloride Potentiometric	103	mmol/L	98 - 107

Interpretation:

Kidney function tests is a collective term for a variety of individual tests and proceduresthat can be done toevaluate how well the kidneys are functioning. Many conditions can affect the ability of the kidneys to carryout their vital functions. Somelead to a rapid (acute) decline in kidney functionothers lead to a gradual (chronic) declineinfunction. Both result in a buildup of toxic waste subst done on urine samples, as well as on blood samples. A number of symptoms may indicate a problem with your kidneys. These include: high blood pressure, blood in urine frequent urges to urinate, difficulty beginning urination, painful urination, swelling in the hands and feet due to a buildup of fluids in the body. A single symptom may not mean something serious. However, when occurring simultaneously, these symptoms suggest that your kidneys are not working properly. Kidney function tests can help determine the reason. Electrolytes (sodium, potassium, and chloride) are present in the human body and the balancing act of the electrolytes in our bodies is essential for normal function of our cells and organs. There has to be a balance. Ionized calcium this test if you have signs of kidney or parathyroid disease. The test may also be done to monitor progress and treatment of these diseases

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.

Dr. Pallavi Rath MBBS, MD (Pathology) Consultant Pathologist

Pallari





Patient Name : Mr Satapathy Lalit Kumar

DOB/Age/Gender : 56 Y/Male Sample Collected : Apr 04, 2024, 10:00 PM

Patient ID / UHID : 1_7835348/RCL7008496 Report Date : Apr 04, 2024, 07:46 PM

Referred By : Self Barcode No : ZC244584
Sample Type : Serum Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range



Lipid Profile

Total Cholesterol Enzymatic - Cholesterol Oxidase	211	mg/dL	<200
Triglycerides Colorimetric - Lip/Glycerol Kinase	227	mg/dL	<150
HDL Cholesterol Accelerator Selective Detergent	47	mg/dL	>40
Non HDL Cholesterol * Calculated	164	mg/dL	<130
LDL Cholesterol * Calculated	118.6	mg/dL	<100
V.L.D.L Cholesterol * Calculated	45.4	mg/dL	< 30
Chol/HDL Ratio * Calculated	4.49	Ratio	3.5 - 5.0
HDL/ LDL Ratio * Calculated	0.4	Ratio	0.5 - 3.0
LDL/HDL Ratio * Calculated	2.52	Ratio	-

Result Rechecked.

Kindly Correlate Clinically.

Adv. Close follow up for confirmation.

Interpretation:

Lipid level assessments must be made following 9 to 12 hours of fasting, otherwise assay results might lead to erroneous interpretation. NCEP recommends of 3 different samples to be drawn at intervals of 1 week for harmonizing biological variables that might be encountered in single assays.

National Lipid Association Recommendations (NLA-2014)	Total Cholesterol (mg/dL)	Triglyceride (mg/dL)	LDL Cholesterol (mg/dL)	Non HDL Cholesterol (mg/dL)
Optimal	<200	<150	<100	<130
Above Optimal			100-129	130 - 159
Borderline High	200-239	150-199	130-159	160 - 189
High	>=240	200-499	160-189	190 - 219
Very High	-	>=500	>=190	>=220

HDL Cholesterol			
Low	High		
<40	>=60		

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.

Dr. Pallavi Rath MBBS, MD (Pathology) Consultant Pathologist

Pallari





NABL

M(EL)T

LABS

NABL-M(EL)T-0060

Patient Name : Mr Satapathy Lalit Kumar

DOB/Age/Gender : 56 Y/Male Sample Collected : Apr 04, 2024, 10:00 PM

Patient ID / UHID : 1_7835348/RCL7008496 Report Date : Apr 04, 2024, 07:46 PM

Referred By : Self Barcode No : ZC244584
Sample Type : Serum Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range

Risk Stratification for ASCVD (Atherosclerotic Cardiovascular Disease) by Lipid Association of India.

Risk Category	A. CAD with > 1 feature of high risk group	
Extreme risk group	B. CAD with >1 feature of very high risk group of recurrent ACS (within 1 year) despite LDL-C <or 50="" =="" disease<="" dl="" mg="" or="" poly="" td="" vascular=""></or>	
Very High Risk	1.Established ASCVD 2.Diabetes with 2 major risk factors of evidence of end organ damage 3. Familial Homozygous Hypercholesterolemia	
High Risk	Three major ASCVD risk factors 2. Diabetes with 1 major risk factor or no evidence of end organ damage 3. CHD stage 3B or 4. 4 LDL >190 mg/dl 5. Extreme of a single risk factor 6. Coronary Artery Calcium - CAC > 300 AU 7. Lipoprotein a >/= 50 mg/dl 8. Non stenotic carotid plaque	
Moderate Risk	2 major ASCVD risk factors	
Low Risk	0-1 major ASCVD risk factors	
M	ajor ASCVD (Atherosclerotic cardiovascular disease) Risk Factors	
1. Age >/=45 years in Males & >/= 55 years in Females	Current Cigarette smoking or tobacco use	
2. Family history of premature ASCVD	4. High blood pressure	
5. Low HDL		

Newer treatment goals and statin initiation thresholds based on the risk categories proposed by Lipid Association of India in 2020.

Risk Group	Treatment Goals		Consider Drug Therapy	
	LDL-C (mg/dl)	Non-HDL (mg/dl)	LDL-C (mg/dl)	Non-HDL (mg/dl)
Extreme Risk Group Category A	<50 (Optional goal <or 30)<="" =="" td=""><td><80 (Optional goal <or 60)<="" =="" td=""><td>>OR = 50</td><td>>OR = 80</td></or></td></or>	<80 (Optional goal <or 60)<="" =="" td=""><td>>OR = 50</td><td>>OR = 80</td></or>	>OR = 50	>OR = 80
Extreme Risk Group Category B	>OR = 30	>OR = 60	> 30	> 60
Very High Risk	<50	<80	>OR = 50	>OR = 80
High Risk	<70	<100	>OR = 70	>OR = 100
Moderate Risk	<100	<130	>OR = 100	>OR = 130
Low Risk	<100	<130	>OR = 130*	>OR = 160

^{*} After an adequate non-pharmacological intervention for at least 3 months.

References: Management of Dyslipidaemia for the Prevention of Stroke: Clinical practice Recommendations from the Lipid Association of India. Current Vascular Pharmacology,2022,20,134-155.

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.

Dr. Pallavi Rath MBBS, MD (Pathology) Consultant Pathologist

Pallari





Patient Name : Mr Satapathy Lalit Kumar

DOB/Age/Gender : 56 Y/Male Sample Collected : Apr 04, 2024, 10:00 PM Patient ID / UHID : 1_7835348/RCL7008496 Report Date : Apr 04, 2024, 08:25 PM

Referred By : Self Barcode No : ZC244584
Sample Type : Serum Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range

Thyroid Profile Total

Triiodothyronine (T3) CMIA	108.3	ng/dL	35 - 193
TotalnThyroxine (T4) CMIA	7.3	μg/dL	4.87 - 11.2
Thyroid Stimulating Hormone (Ultrasensitive) CMIA	1.87	μIU/mL	0.35 - 4.94

Interpretation:

Pregnancy	Reference ranges TSH
1 st Trimester	0.1 - 2.5
2 ed Trimester	0.2 - 3.0
3 rd Trimester	0.3 - 3.0

Primary malfunction of the thyroid gland may result in excessive (hyper) or below normal (hypo) release of T3 or T4. In addition as TSH directly affects thyroid function, malfunction of the pituitary or the hypo - thalamus influences the thyroid gland activity. Disease in any portion of the thyroid-pitutary-hypothala- mus system may influence the levels of T3 and T4 in the blood. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels may be low. In addition, in the Euthyroid Sick Syndrome, multiple alterations in serum thyroid function test findings have been recognized in patients with a wide variety of non-thyroidal illnesses (NTI) without evidence of preexisting thyroid or hypothalami c-pitutary diseases. Thyroid Binding Globulin (TBG) concentrations remain relatively constant in healthy individuals. However, pregnancy, excess estrogen's, androgen's, antibiotic steroids and glucocorticoids are known to alter TBG levels and may cause false thyroid values for Total T3 and T4 tests.

TSH	T4	T3	INTERPRETATION
High	Normal	Normal	Mild (subclinical) hypothyroidism
High	Low	Low or normal	Hypothyroidism
Low	Normal	Normal	Mild (subclinical) hyperthyroidism
Low	High or normal	High or normal	Hyperthyroidism
Low	Low or normal	Low or normal	Nonthyroidal illness; pituitary (secondary) hypothyroidism
Normal	High	High	Thyroid hormone resistance syndrome (a mutation in the thyroid hormone receptor decreases thyroid hormone function)

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



Pallari





Patient Name : Mr Satapathy Lalit Kumar

DOB/Age/Gender : 56 Y/Male Sample Collected : Apr 04, 2024, 10:00 PM Patient ID / UHID : 1_7835348/RCL7008496 Report Date : Apr 04, 2024, 08:25 PM

: ZC244584 Referred By : Self Barcode No Sample Type : Serum Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range

Prostate Specific Antigen (PSA) Total

>=70

Prostate Specific Antigen-Total (PSA-Total) CMIA	0.2 ng/mL		<4	.0	
Interpretation:					
Age (years)	Ranges				
< 40	<1.4				
40 - 49	<2.0				
50 - 59	<3.1				
60 - 69	<4.1				

Prostate Specific Antigen (PSA) is a single-chain glycoprotein normally found in the cytoplasm of the epithelial cells lining the acini and ducts of the prostate gland. PSA is detected in the serum of males with normal, benign hyperplastic and malignant prostate tissue and in patients with prostatetiss. PSA is not detected (or detected at very low levels) in the serum of males without prostate tissue (because of radical prostatectomy or cytoprostatectomy) or in the serum of most females. The fact that PSA is unique to prostate tissue makes it a suitable marker for monitoring men with cancer of the prostate. PSA is also useful for determining possible recurrence after therapy when used in conjunction with other diagnostic indices. PSA levels increase in men with cancer of the prostate. After radical prostatectomy PSA levels routinely fall to a very low level, which may not be seen in patients undergoing radiation therapy. Monitoring PSA levels appears to be useful in detecting residual disease and early recurrence of tumor. Therefore, serial PSA levels can help determine the success of prostatectomy and the need for further treatment, such as radiation, endocrine of

chemotherapy and in the monitoring of the effectiveness of therapy.

PSA levels should not be interpreted as absolute evidence of presence or the absence of malignant disease. Before treatment, patients with confirmed prostate carcinoma frequently have levels of PSA within the range observed in healthy individuals. Elevated levels of PSA can be observed in the patients with nonmalignant disease. Measurement of PSA should always be used in conjunction with other diagnostic procedures, including information from the patients and

The concentration of total PSA in a given specimen determined with assays from different manufacturers can vary due to differences in assay methods, calibration, and reagent specificity.

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.

Dr. Pallavi Rath MBBS, MD (Pathology) **Consultant Pathologist**

Pallari





Patient Name : Mr Satapathy Lalit Kumar

DOB/Age/Gender : 56 Y/Male Sample Collected : Apr 04, 2024, 10:00 PM

Patient ID / UHID : 1_7835348/RCL7008496 Report Date : Apr 04, 2024, 07:40 PM

Referred By : Self Barcode No : YA485024
Sample Type : Spot Urine Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range

NABL M(EL)T LABS

Urine Routine and Microscopic Examination

Physical Examination *			
Volume *	20	ml	-
Colour *	Pale yellow	-	Pale yellow
Transparency *	Clear	-	Clear
Deposit *	Absent	-	Absent
Chemical Examination *			
Reaction (pH) Double Indicator	5	-	4.5 - 8.0
Specific Gravity Ion Exchange	1.02	-	1.010 - 1.030
Urine Glucose (sugar) Oxidase / Peroxidase	Negative	-	Negative
Urine Protein (Albumin) Acid / Base Colour Excahnge	Negative	-	Negative
Urine Ketones (Acetone) Legals Test	Negative	-	Negative
Blood Peroxidase Hemoglobin	Positive(+)	-	Negative
Leucocyte esterase Enzymatic Reaction	Negative	-	Negative
Bilirubin Urine Coupling Reaction	Negative	-	Negative
Nitrite Griless Test	Negative	-	Negative
Urobilinogen Ehrlichs Test	Normal	-	Normal
Microscopic Examination *			
Pus Cells (WBCs) *	1-2	/hpf	0 - 5
Epithelial Cells *	1-2	/hpf	0 - 4
Red blood Cells *	1-2	/hpf	Absent
Crystals *	Absent	-	Absent
Cast *	Absent	-	Absent
Yeast Cells *	Absent	-	Absent
Amorphous deposits *	Absent	-	Absent
Bacteria *	Absent	-	Absent
Protozoa *	Absent	-	Absent

Dr. Pallavi Rath MBBS, MD (Pathology) Consultant Pathologist

Pallari



^(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



Patient Name : Mr Satapathy Lalit Kumar

 DOB/Age/Gender
 : 56 Y/Male
 Sample Collected
 : Apr 04, 2024, 10:00 PM

 Patient ID / UHID
 : 1_7835348/RCL7008496
 Report Date
 : Apr 04, 2024, 07:41 PM

Referred By : Self Barcode No : YA485025
Sample Type : URINE F Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range

Urine Glucose Fasting

Urine Glucose (sugar)	Negative	-	Negative
Oxidase / Peroxidase			

*** End Of Report ***

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



Paulari



Booking Centre :- Madyosis Diagnostics, Office No-406, 4th Floor, Bhakti Genesis, Wakad Rd, Shedge Vasti, Shankar Kalat Nagar, Wakad, Pimpri-Chinchwad, Maharashtra 411057

Processing Lab: Redcliffe Lifetech Pvt. Ltd., First Floor, B Wing. Aswani Chambers, S.No. 199+204+205 206/1, 209/1,

Terms and Conditions of Reporting

- 1. The presented findings in the Reports are intended solely for informational and interpretational purposes by the referring physician or other qualified medical professionals possessing a comprehensive understanding of reporting units, reference ranges, and technological limitations. The laboratory shall not be held liable for any interpretation or misinterpretation of the results, nor for any consequential or incidental damages arising from such interpretation.
- 2. It is to be presumed that the tests performed pertain to the specimen/sample attributed to the Customer's name or identification. It is presumed that the verification particulars have been cleared out by the customer or his/her representation at the point of generation of said specimen / sample. It is hereby clarified that the reports furnished are restricted solely to the given specimen only.
- 3. It is to be noted that variations in results may occur between different laboratories and over time, even for the same parameter for the same Customer. The assays are performed and conducted in accordance with standard procedures, and the reported outcomes are contingent on the specific individual assay methods and equipment(s) used, as well as the quality of the received specimen.
- 4. This report shall not be deemed valid or admissible for any medico-legal purposes.
- 5. The Customers assume full responsibility for apprising the Company of any factors that may impact the test finding. These factors, among others, includes dietary intake, alcohol, or medication / drug(s) consumption, or fasting. This list of factors is only representative and not exhaustive.



Patient Name : Mr Satapathy Lalit Kumar

DOB/Age/Gender : 56 Y/Male Sample Collected : Apr 04, 2024, 10:00 PM

Patient ID / UHID : 7835458/RCL7008496 Report Date : Apr 04, 2024, 07:46 PM

Referred By : Self Barcode No : ZC244579
Sample Type : FLUORIDE PP Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range



Glucose Post Prandial (BSPP)

Glucose Post Prandial	98	mg/dL	70 - 140
(Fluoride Plasma-P,Hexokinase)			

Interpretation:

Status	PP plasma glucose in mg/dL	
Normal	<140	
Impaired glucose tolerance	140 - 199	
Diabetes	=>200	

Reference: American Diabetes Association

Comment

Blood glucose determinations in commonly used as an aid in the diagnosis and treatment of diabetes. Elevated glucose levels (hyperglycemia) may also occur with pancreatic neoplasm, hyperthyroidism, and adrenal cortical hyper function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy insulinoma, or various liver diseases.

Note

1. The diagnosis of Diabetes requires a fasting plasma glucose of > or = 126 mg/dL or a random / 2 hour plasma glucose value of > or = 200 mg/dL with symptoms of diabetes mellitus.

2. Very high glucose levels (>450 mg/dL in adults) may result in Diabetic Ketoacidosis.

*** End Of Report ***





Terms and Conditions of Reporting

- 1. The presented findings in the Reports are intended solely for informational and interpretational purposes by the referring physician or other qualified medical professionals possessing a comprehensive understanding of reporting units, reference ranges, and technological limitations. The laboratory shall not be held liable for any interpretation or misinterpretation of the results, nor for any consequential or incidental damages arising from such interpretation.
- 2. It is to be presumed that the tests performed pertain to the specimen/sample attributed to the Customer's name or identification. It is presumed that the verification particulars have been cleared out by the customer or his/her representation at the point of generation of said specimen / sample. It is hereby clarified that the reports furnished are restricted solely to the given specimen only.
- 3. It is to be noted that variations in results may occur between different laboratories and over time, even for the same parameter for the same Customer. The assays are performed and conducted in accordance with standard procedures, and the reported outcomes are contingent on the specific individual assay methods and equipment(s) used, as well as the quality of the received specimen.
- 4. This report shall not be deemed valid or admissible for any medico-legal purposes.
- 5. The Customers assume full responsibility for apprising the Company of any factors that may impact the test finding. These factors, among others, includes dietary intake, alcohol, or medication / drug(s) consumption, or fasting. This list of factors is only representative and not exhaustive.



Patient Name : Mr Satapathy Lalit Kumar

 DOB/Age/Gender
 : 56 Y/Male
 Sample Collected
 : Apr 04, 2024, 10:00 PM

 Patient ID / UHID
 : 1_7835921/RCL7008496
 Report Date
 : Apr 05, 2024, 12:16 PM


Referred By : Self Barcode No : YA485952
Sample Type : URINE PP Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range

Urine Glucose PP

Urine Glucose (sugar)	Negative	-	Negative
-----------------------	----------	---	----------

*** End Of Report ***



Pallari



Terms and Conditions of Reporting

- 1. The presented findings in the Reports are intended solely for informational and interpretational purposes by the referring physician or other qualified medical professionals possessing a comprehensive understanding of reporting units, reference ranges, and technological limitations. The laboratory shall not be held liable for any interpretation or misinterpretation of the results, nor for any consequential or incidental damages arising from such interpretation.
- 2. It is to be presumed that the tests performed pertain to the specimen/sample attributed to the Customer's name or identification. It is presumed that the verification particulars have been cleared out by the customer or his/her representation at the point of generation of said specimen / sample. It is hereby clarified that the reports furnished are restricted solely to the given specimen only.
- 3. It is to be noted that variations in results may occur between different laboratories and over time, even for the same parameter for the same Customer. The assays are performed and conducted in accordance with standard procedures, and the reported outcomes are contingent on the specific individual assay methods and equipment(s) used, as well as the quality of the received specimen.
- 4. This report shall not be deemed valid or admissible for any medico-legal purposes.
- 5. The Customers assume full responsibility for apprising the Company of any factors that may impact the test finding. These factors, among others, includes dietary intake, alcohol, or medication / drug(s) consumption, or fasting. This list of factors is only representative and not exhaustive.






Mr. Lalir Kumar Satapathy

Wagholi Shivranjani Ahmedabad Gujarat India

Gendr/DOB (Age) **Medico ID** : 24040402609116 : Male/04-Apr-1968(56Y 0M) Referred By Date : 04-Apr-2024 / 11:22 AM

History



			3333 3333 3333 3333 3333				
VITALS	:	TEMP	: - (F)	PULSE RATE	: - /MIN	RBS	: - mg/dL
		HR	: 70 /MIN	BP	: 0 / 0 mmHg	SPO2	: 0.0 %
MEASUREMENTS*	:	PR	: 139.35 ms	QT	: 385.42 ms	Р	: 52.1 deg
(ECG Parameters)		ST	: 0.45 ms	QTc	: 415.0 ms	QRs	: 65.64 deg
		R-R	: 862.5 ms	QRS	: 65.64 ms	Т	: 21.09 deg

FINDINGS : NORMAL SINUS RHYTHM. NO SIGNIFICANT ST CHANGES NOTED


: THIS ECG IS FOUND TO BE WITHIN NORMAL LIMITS. **IMPRESSION**

: CLINICAL CORRELATION RECOMMENDATION

This is electronically authenticated report; hence doesn't require signature.

* Software calculated values; to be verified manually.

Printed By: M4 Diagnostics Center On 05-Apr-2024 / 12:57 PM (Rs. 300.00/- Received for this ECG)



Reported By **Express Diagnostics HQ**



Name : MR. LALIT KUMAR SATAPATHY Age/Sex : 56 YEARS/M

Ref By : Dr. MADYOASIS MEDICAL SERVICES -- Date : 04 Apr 2024

2D ECHOCARDIOGRAPHY & COLOUR DOPPLER STUDY

Friday, April 05, 2024

Left Ventricle:

The left ventricle is normal in size. No e/o RWMA.

The left ventricular ejection fraction is normal .

Left Atrium:

The left atrium is normal size. No clot.

Right Ventricle:

The right ventricular is normal size. There is normal right Ventricular wall thickness.

Aorta:

The aortic root is normal.

Pulmonary Artery:

The Pulmonary artery is normal.

Pericardium:

There is no pericardial effusion. No calcification.

Aortic Valve:

The aortic valve is tri-leaflet with thin, pliable leaflets that move normally. There is no aortic Stenosis. No aortic regurgitation is present.

Mitral Valve:

The mitral valve leaflets are thin. Normal mitral gradients. There is no evidence of stenosis, prolapse. Diastolic flows are altered . No mitral regurgitation noted.

Tricuspid Valve:

The tricuspid valve leaflets are thin and pliable and the valve motion is normal. No tricuspid Regurgitation is noted.

Pulmonary Valve:

The pulmonary valve leaflets are thin and pliable and the valve motion is normal. No pulmonary Valvular regurgitation is noted.

Proximal Coronaries:

Not visualized.

IAS and IVS are intact.

M-MODE/2D PARAMETERS

AO	29	(23-37mm)
LA	28	(19-40mm)
RVD		(7-23mm)
LVD	41	(35-55mm)
LVS	29	(24-42mm)
IVS	10.2	(6-11mm)
LVPW	10.2	(6-11mm)
EF	55-60%	(50-70%)

Parameters in brackets indicate normal adult Values.

IMPRESSION:

- No e/o RWMA
- Normal EF.
- RA / RV not dilated.
- No e/o pulmonary hypertension
- Normal valves and velocities.
- No clot, vegetations or effusions.



Dr Ganesh Sanap MBBS DMRD DNB





Patient Name : MR. LALIT KUMAR SATAPATHY Date : 04 Apr 2024

Referred By : Dr. MADYOASIS MEDICAL SERVICES - Age : 56 YEARS Sex : M

-

USG ABDOMEN AND PELVIS

Liver:

The liver is normal in size and shows increased echotexture. No focal lesion is seen. The intrahepatic biliary radicles are normal. The common bile duct and the portal vein appear normal.

Gall Bladder

The gall bladder is well distended. No e/o calculus seen_ . The wall thickness is normal.

Pancreas

The pancreas is normal in size and shape. No focal lesion or calcifications are seen within it. The pancreatic duct is normal.

Spleen

The spleen measures 10.1cm in size and is normal in echotexture. No focal lesion is seen.

Kidneys

The right kidney measures 9.6 x 4.1cm. The left kidney measures 9.8 x 5.4 cm. Both kidneys show normal parenchymal echo texture. The cortico-medullary differentiation is maintained bilaterally. The pelvicalyceal system is normal in both the kidneys.

Aorta/IVC

The aorta and IVC appear grossly normal. No ascites or lymphadenopathy is seen.

Urinary bladder

The bladder is well distended. The wall thickness is normal. No vesical calculus is seen.

Prostate

The prostate corresponding to a weight of about 21 gms. No focal lesion or calcification is seen.

Impression

- Diffuse fatty infiltration of liver- Grade I.
- No other abnormality seen.



Dr Ganesh Sanap MBBS DMRD DNB







Patient Name: MR. LALIT KUMAR SATAPATHY Date: 04 Apr 2024
Ref. By: Dr. MADYOASIS MEDICAL SERVICES -- Age/sex :56 YEARS/M

X RAY CHEST PA VIEW

Both the lung fields are clear.

Both diaphragmatic domes have normal contours and positions.

Cardio-aortic silhouette has a normal appearance.

There is no evidence of any pleural effusion.

Bony thorax appears normal

IMPRESSION:

No obvious abnormality seen at present study.



Dr Ganesh Sanap MBBS DMRD DNB



🕜 Shop/ofc. No. 2, Ground floor, Building A, City Vista Downtown Kolte Patil , opposite Victorius School, Kharadi Pune 411014

