

#### CERTIFICATE OF MEDICAL FITNESS

NAME: Varanahalaxmi	
AGE/GENDER: 534 Fernale	
HEIGHT: 153 cm	WEIGHT: 56.3 kg
IDENTIFICATION MARK:	
BLOOD PRESSURE: 110/70 mm49	
PULSE: 94 b/m	
CVS: Gromal RS:P	
ANY OTHER DISEASE DIAGNOSED IN THE PAST: ( )	
ALLERGIES, IF ANY:	
LIST OF PRESCRIBED MEDICINES:	
ANY OTHER REMARKS:	
of Ms Sheshappa who has signed in midisease and is fit for employment.	
S. Varanaha Jak Shmi	Dr. BINDURAJ, R Internal BBS, MD
Signature of candidate	Signature of Medical Officer
Place: Sportoum Diagnostick theats	th core
Date: <u>@G   D4   2 </u>	
	=1

Disclaimer: The patient has not been checked for COVID. This certificate does not relate to the covid status of the patient examined





NAME	AGE	GENDER
MM. Valornohelakelmi	5377	fende

### DENTAL EXAMINATION REPORT:

8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8
8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8

C: CAVITY -> DOC ON - FOR INVESS KET & Clovn.

M: MISSING

O: OTHERS -> Incomplete orthodalae beauty in

gays bloo feets.

ADVISED:

CLEANING / SCALING / ROOTS PLANNING / FLOSSING & POLISHING / OTHERS

**REMARKS:** 

SIGNATURE OF THE DENTAL SURGEON

SEAL

DATE

Dr. SACHDEV NAGARKAR B.D.S., F.A.G.E., F.P.F.A. (USA) Reg. No: 2247/A







Dr. Ashok S Bsc., MBBS., D.O.M.S Consultant Opthalmologist KMC No: 31827 DATE: 06 04.24.

#### EYE EXAMINATION

NAME: MS. Volamaha	laksani AGE: 53 y	GENDER: F/M
	RIGHT EYE	LEFT EYE
Vision	6137:NO	6/3/10
Vision With glass	616g-N6	Olep role
Color Vision	Normal	Normal
Anterior segment examination	Normal	Normal
Fundus Examination	Normal	Normal
Any other abnormality	Nill	Nill
Diagnosis/ impression	Normal	Normal
	Dr. ASHOI  Consultant (6)	K SARODHE M.B.B.S., D.O.M.S.  pthalmologist) 31827

SCAN FOR LOCATION



ID: 240031	06-04-2024 10·50·34 Fo	or Rel
MRS VARAMAHALAXMI	<i>R</i>	Diagnosis Information:
Female 53Years	PR : 149 ms	Sinus Rhythm Abnormal q and Q Wave(III,VI)
	S : 81 QTc : 348/4 RS/T : 54/23 5/SV1 : 0.901	Flat I Wave(V5,V6)
	. 0.701/0.017	Report Confirmed by:
		VI
		\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\
aVF		



NAME : MRS. VARAMAHALAXMI	DATE : 05/04/2024
NAME : MRS. VARAMAHALAXMI AGE/SEX : 53 YEARS/FEMALE	REG NO: 0504240031
REF BY : APOLO CLINIC	

# CHEST PA VIEW

- Visualised lungs are clear .
- Bilateral hila appears normal.
- Cardia is normal in size
- No pleural effusion

IMPRESSION: No significant abnormality .

Traveors

DR PRAVEEN B, DMRD, DNB Consultant Radiologist





### **SPECTRUM DIAGNOSTICS**

Bangalore


Patient ID: 0304

Name: VARAMAHALAKSHMI

CR Number : 20240406103404 Registration Date : 06-Apr-2024 Age: 53

Gender: Female

Operator: spectrum diagnostics



	125 Hz	250 Hz	500 Hz	750 Hz	1000 H	1500 H	2000 H	3000 H	4000 H	6000 H	8000 H
X - Air Left	25	(20)	20	25	25	20	20	(15)	20	20	20
O - Air Right	25	25	15	20	25	25	25	15	20	20	15
> - Bone Left											
< - Bone Right						75-					

	Average	High	Mid	Low
AIR Left	20.91 dB	18.75 dB	21.67 dB	22.50 dB
AIR Right	20.91 dB	17.50 dB	25.00 dB	21.25 dB

#### **Clinical Notes:**

Not Found





PATIENT NAME	MRS VARAMAHALAXMI	ID NO	230031
AGE	53 YEARS	SEX	MALE
REF BY	DRAPOLLO CLINIC	DATE	06.04.2024

## 2D ECHO CARDIOGRAHIC STUDY

### M-MODE

	1111000
AORTA	27mm
LEFT ATRIUM	35mm
RIGHT VENTRICLE	18mm
LEFT VENTRICLE (DIASTOLE )	44mm
LEFT VENTRICLE(SYSTOLE)	30mm
VENTRICULAR SEPTUM (DIASTOLE)	11mm
VENTRICULAR SEPTUM (SYSTOLE)	12mm
POSTERIOR WALL (DIASTOLE)	10mm
POSTERIOR WALL (SYSTOLE)	11mm
FRACTIONAL SHORTENING	30%
EJECTION FRACTION	60%

## DOPPLER /COLOUR FLOW

MITRAL VALVE	E-0.85 m/sec	A-0.62m/sec	TRIVIAL MR
AORTIC VALVE	1.00 m/sec		NO AR
PULMONARY VALVE			NO PR
TO BE A DOUBLE OF THE	1.20 m/sec		
TRISCUSPID VALVE	128 228 527 122		
	THE RESERVE OF THE	20mmHg	TRIVIAL TR







PATIENT NAME	MRS VARAMAHALAXMI	ID NO	230031
AGE	53 YEARS	SEX	MALE
REF BY	DRAPOLLO CLINIC	DATE	06.04.2024

### 2D ECHO CARDIOGRAHIC STUDY

LEFT VENTRICLE SIZE& THICKNESS		NORMAL		
CONTRACTILITY	REGIONAL GLOBAL	NO RWMA		

RIGHT VENTRICLE : NORMAL LEFT ATRIUM : NORMAL RIGHT ATRIUM: NORMAL MITRAL VALVE : NORMAL AORTIC VALVE : NORMAL PULMONARY VALVE: NORMAL TRICUSPID VALVE: NORMAL INTER ATRIAL SEPTUM: INTACT INTER VENTRICULAR SEPTUM: INTACT PERICARDIUM: NORMAL **OTHERS** 

#### IMPRESSION

- NO RWMA OF LV AT REST
- NORMAL LV FUNCTION, LVEF-60%
- NO LV DIASTOLIC DYSFUNCTION
- NO CLOT / VEGITATION / PERICARDIAL EFFUSION

MS LIKITHA M ECHO TECHNICIAN

The science of radiology is based upon interpretation of shadows of normal and abnormal tissue. This is neither complete nor accurate; hence, findings should always be interpreted in to the light of clinico-pathological correction.







NAME AND LAB NO	MRS VARAMAHALAXMI	REG -40031
AGE & SEX	53 YRS	FEMALE
DATE AND AREA OF INTEREST	06.04.2024	ABDOMEN & PELVIS
REF BY	C/O APOLO CLINIC	

USG ABDOMEN AND PELVIS

LIVER: Normal in size and shows diffuse increased echogenicity

> No e/o IHBR dilatation. No evidence of focal lesion Portal vein appears normal. CBD appears normal.

**GALL BLADDER:** Partially distended .No obvious calculus in the visualised luminal portion

SPLEEN: Normal in size and echotexture. No focal lesion

PANCREAS: Head and body appears normal. Tail obscured by bowel gas shadows

RETROPERITONEUM: Suboptimal visualised due to bowel gas.

RIGHT KIDNEY: Right kidney is normal in size & echotexture

No evidence of calculus/ hydronephrosis.

LEFT KIDNEY: Left kidney is normal in size & echotexture

No evidence of calculus/ hydronephrosis.

**URINARY BLADDER:** Well distended. No wall thickening/ calculi.

**UTERUS & OVARIES:** Atrophic

No obvious adnexal mass lesions.

No evidence of ascites/pleural effusion.

#### IMPRESSION:

Grade I fatty liver.

Suggested clinical / lab correlation.

DR PRAVEEN B, DMRD, DNB CONSULTANT RADIOLOGIST









Age / Gender : 53 years / Female

**Ref. By Dr.** : Dr. APOLO CLINIC **Reg. No.** : 0604240031

C/o : Apollo Clinic

Bill Date : 06-Apr-2024 08:57 AM
Sample Col. Date : 06-Apr-2024 08:57 AM
Result Date : 06-Apr-2024 11:39 AM

0604240031 Report Status : Final

Test Name	Result	Unit	Reference Value	Method	
Glycosylated Haemoglobin (HbA1c)-Whole Blood EDTA					***************************************
Glycosylated Haemoglobin	5.40	%	Non diabetic adults :<5.7	HPLC	
IbA1c)			At risk (Prediabetes): 5.7 - 6.4	4	
			Diagnosing Diabetes :>= 6.5		
			Diabetes		
			Excellent Control: 6-7		
			Fair to good Control: 7-8 Unsatisfactory Control:8-10		
Sstimated Average	108.28	m a/JT	Poor Control :>10		
Glucose(eAG)	100.20	mg/dL		Calculated	

Note: 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not

Comments: HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.



Printed By

: spectrum

Printed On : 0

: 06 Apr, 2024 05:02 pm

N. Chum

Dr. Nithun Reddy C,MD,Consultant Pathologist







Age / Gender : 53 years / Female Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 0604240031 C/o : Apollo Clinic **Bill Date** 

: 06-Apr-2024 08:57 AM

Sample Col. Date: 06-Apr-2024 08:57 AM **Result Date** : 06-Apr-2024 01:35 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Lipid Profile-Serum				
Cholesterol Total-Serum	257.00	mg/dL	Female: 0.0 - 200	Cholesterol
Triglycerides-Serum	314.00	mg/dL	Female: 0.0 - 150	Oxidase/Peroxidase Lipase/Glycerol
High-density lipoprotein (HDL) Cholesterol-Serum	61.00	mg/dL	Female: 40.0 - 60.0	Dehydrogenase Accelerator/Selective
Non-HDL cholesterol-Serum	196	mg/dL	Female: 0.0 - 130	Detergent Calculated
Low-density lipoprotein (LDL) Cholesterol-Serum	172.0	mg/dL	Female: 0.0 - 100.0	Cholesterol esterase and cholesterol
Very-low-density lipoprotein (VLDL) cholesterol-Serum	63	mg/dL	Female: 0.0 - 40	oxidase Calculated
Cholesterol/HDL Ratio-Serum	4.21	Ratio	Female: 0.0 - 5.0	Calculated

: 0604240031

0604240031

UHID

#### Interpretation:

Parameter	Desirable	Borderline High		
Total Cholesterol		8	High	Very High
	<200	200-239	>240	
Triglycerides	<150	150-199	200-499	>500
Non-HDL cholesterol	<130	160-189		
Low-density lipoprotein (LDL) Cholesterol		100-189	190-219	>220
dem density apoptotem (LDL) Cholesterol	<100	100-129	160-189	>190

Comments: As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atherosclerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus, sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.



Printed By

: spectrum

Printed On

: 06 Apr, 2024 05:02 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 2 of 12 Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010









Age / Gender : 53 years / Female Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 0604240031

C/o : Apollo Clinic **Bill Date** 

: 06-Apr-2024 08:57 AM

Sample Col. Date: 06-Apr-2024 08:57 AM

**Result Date** Report Status : Final

: 06-Apr-2024 01:35 PM

Test Name	Result	Unit	Reference Value	Method
LFT-Liver Function Test -Seru	m			
Bilirubin Total-Serum	0.48	mg/dL	0.2-1.0	Caffeine
Bilirubin Direct-Serum	0.08	mg/dL	0.0-0.2	Benzoate Diazotised Sulphanilic
Bilirubin Indirect-Serum Aspartate Aminotransferase AST/SGOT)-Serum	0.40 20.00	mg/dL U/L	Female: 0.0 - 1.10 Female: 15.0 - 37.0	Acid Direct Measure UV with
Alanine Aminotransferase ALT/SGPT)-Serum	18.00	U/L	Female: 14.0 - 59.0	Pyridoxal - 5 - Phosphate UV with Pyridoxal - 5 -
lkaline Phosphatase (ALP)- erum	90.00	U/L	Female: 45.0 - 117.0	Phosphate PNPP,AMP- Buffer
rotein, Total-Serum	7.05	g/dL	6.40-8.20	Biuret/Endpoint-
lbumin-Serum	4.48	g/dL	Female: 3.40 - 5.50	With Blank Bromocresol
lobulin-Serum bumin/Globulin Ratio-Serum	2.57 1.74	g/dL Ratio	2.0-3.50 0.80-2.0	Purple Calculated Calculated

UHID

: 0604240031

0604240031



Printed By

: spectrum

Printed On : 06 Apr, 2024 05:02 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

**SCAN FOR LOCATION** 

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010







Age / Gender : 53 years / Female Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 0604240031

C/o : Apollo Clinic **Bill Date** 

: 06-Apr-2024 08:57 AM

Sample Col. Date: 06-Apr-2024 08:57 AM **Result Date** 

**Report Status** : Final

: 06-Apr-2024 01:35 PM

Test Name	Result	Unit	Reference Value	Method
Calcium, Total- Serum	9.10	mg/dL	8.50-10.10	Spectrophotometry (O- Cresolphthalein
Gamma-Glutamyl Transferas (GGT)-Serum	15.00	U/L	Male: 15.0-85.0	complexone) Other g-Glut-3-
			Female: 5.0-55.0	carboxy-4 nitro

0604240031

: 0604240031

UHID

Comments: Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system, and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of serum GGT levels in detecting alcohol-induced liver disease. Elevated serum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.

Fasting Urine Glucose-Urine	Negative		Negative	Dipstick/Benedicts
Fasting Blood Sugar (FBS)- Plasma	89	mg/dL	60.0-110.0	(Manual) Hexo Kinase









Name

: MRS. VARAMAHALAXMI

Age / Gender Ref. By Dr.

: 53 years / Female : Dr. APOLO CLINIC

Reg. No. C/o

: 0604240031

: Apollo Clinic

**Bill Date** 

**Result Date** 

: 06-Apr-2024 08:57 AM

Sample Col. Date: 06-Apr-2024 08:57 AM

: 06-Apr-2024 01:35 PM

Report Status : Final

**Test Name** 

Result

Unit

UHID

Reference Value

: 0604240031

Method

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

0604240031

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total



Printed By

: spectrum

Printed On

: 06 Apr, 2024 05:02 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 +91 77604 97644 | 080 2337 1555

linfo@spectrumdiagnostics.org







Age / Gender : 53 years / Female

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 0604240031

C/o : Apollo Clinic **Bill Date** : 06-Apr-2024 08:57 AM

Sample Col. Date: 06-Apr-2024 08:57 AM **Result Date** : 06-Apr-2024 01:35 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Thyroid function tests (TFT Serum	)-			
Tri-Iodo Thyronine (T3)-Sei	rum 1.08	ng/mL	Female: 0.60 - 1.81	Chemiluminescence Immunoassay
Гhyroxine (Т4)-Serum	6.90	μg/dL	Female: 5.50 - 12.10	(CLIA) Chemiluminescence Immunoassay
Thyroid Stimulating Hormon TSH)-Serum	ne 3.40	μIU/mL	Female: 0.35 - 5.50	(CLIA) Chemiluminescence Immunoassay (CLIA)

0604240031

: 0604240031

Comments: Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population without

Reference range: Cord: (37 Weeks): 0.5-1.41, Children:1-3 Days: 1.0-7.40,1-11 Months: 1.05-2.45,1-5 Years: 1.05-2.69,6-10 Years: 0.94-2.41,11-15

Reference range: Adults: 20-50 Years: 0.70-2.04, 50-90 Years: 0.40-1.81,

Reference range in Pregnancy: First Trimester: 0.81-1.90, Second Trimester: 1.0-2.60

Increased Levels: Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG). Decreased Levels: Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

Comments: Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4). It also helps to monitor treatment of Hyperthyroidism with

Reference Range: Males: 4.6-10.5, Females: 5.5-11.0, 60 Years: 5.0-10.70, Cord: 7.40-13.10, Children: 1-3 Days: 11.80-22.60, 1-2 Weeks: 9.90-16.60,1-4 Months: 7.20-14.40,1-5 Years: 7.30-15.0,5-10 Years: 6.4-13.3

1-15 Years: 5.60-11.70, Newborn Screen: 1-5 Days: >7.5,6 Days : >6.5

Increased Levels: Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia, Increased transthyretin, estrogen therapy, pregnancy. Decreased Levels: Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.

Comments: TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, caloric intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after  $\sim 3$  weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality. Reference range in Pregnancy: I- trimester:0.1-2.5; II -trimester:0.2-3.0; III- trimester:0.3-3.0

Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks:1.7-9.1

Increased Levels: Primary hypothyroidism, Subclinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance. els: Graves disease, Autonomous thyroid hormone secretion, TSH defic

Printed By : spectrum

Printed On : 06 Apr, 2024 05:02 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010

+91 77604 97644 | 080 2337 1555

info@spectrumdiagnostics.org





Age / Gender Ref. By Dr.

: 53 years / Female

Reg. No.

: Dr. APOLO CLINIC : 0604240031

C/o : Apollo Clinic **UHID** : 0604240031

0604240031

**Bill Date** 

: 06-Apr-2024 08:57 AM

Sample Col. Date: 06-Apr-2024 08:57 AM **Result Date** : 06-Apr-2024 01:35 PM

**Report Status** : Final

Test Name	Result	Unit	Reference Value	Method
Complete Haemogram-Whole	Blood EDTA			
Haemoglobin (HB)	11.60	g/dL	Male: 14.0-17.0 Female:12.0-15.0	Spectrophotmete
Red Blood Cell (RBC)	3.80	million/cu	Newborn:16.50 - 19.50 mm3.50 - 5.50	Volumetric
Packed Cell Volume (PCV)	32.70	%	Male: 42.0-51.0 Female: 36.0-45.0	Impedance Electronic Pulse
Mean corpuscular volume (MCV)	86.10	fL	78.0- 94.0	Calculated
Mean corpuscular hemoglobin (MCH)		pg	27.50-32.20	Calculated
Mean corpuscular hemoglobin concentration (MCHC)	35.50	%	33.00-35.50	Calculated
Red Blood Cell Distribution Width SD (RDW-SD)	42.00	fL	40.0-55.0	Volumetric
Red Blood Cell Distribution CV (RDW-CV)	15.40	%	Male: 11.80-14.50 Female: 12.20-16.10	Impedance Volumetric
Mean Platelet Volume (MPV)	9.90	fL	8.0-15.0	Impedance Volumetric
Platelet	2.99	lakh/cumm	1.50-4.50	Impedance Volumetric
latelet Distribution Width PDW)	10.60	%	8.30 - 56.60	Impedance Volumetric
White Blood cell Count (WBC)	6640.00	cells/cumm	Male: 4000-11000 Female 4000-11000 Children: 6000-17500 Infants: 9000-30000	Impedance Volumetric Impedance
eutrophils	58.70	%	40.0-75.0	Light
mphocytes	36.50	%	20.0-40.0	scattering/Manual Light
sinophils	1.80	%	0.0-8.0	scattering/Manual Light scattering/Manual









Age / Gender : 53 years / Female Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 0604240031 C/o : Apollo Clinic

**Bill Date** 

: 06-Apr-2024 08:57 AM

Sample Col. Date: 06-Apr-2024 08:57 AM **Result Date** 

: 06-Apr-2024 01:35 PM Report Status

: Final

Test Name	Result	Unit	Reference Value	Method
Monocytes	3.00	%	0.0-10.0	Light
Basophils	0.00	%	0.0-1.0	scattering/Manual Light
Absolute Neutrophil Count Absolute Lymphocyte Count Absolute Monocyte Count Absolute Eosinophil Count Absolute Basophil Count Erythrocyte Sedimentation Rate (ESR)	3.90 2.42 0.20 120.00 0.00 42	10^3/uL 10^3/uL 10^3/uL cells/cumm 10^3/uL mm/hr	2.0- 7.0 1.0-3.0 0.20-1.00 40-440 0.0-0.10 Female: 0.0-20.0 Male: 0.0-10.0	scattering/Manual Calculated Calculated Calculated Calculated Calculated Calculated Westergren

0604240031

: 0604240031

UHID

# Peripheral Smear Examination-Whole Blood EDTA

Method: (Microscopy-Manual)

RBC'S : Normocytic Normochromic.

WBC'S : Are normal in total number, morphology and distribution. Platelets

: Adequate in number and normal in morphology. No abnormal cells or hemoparasites are present.

Impression: Mild degree of Normocytic Normochromic Anaemia.




Printed By

: spectrum

Printed On : 06 Apr, 2024 05:02 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

SCAN FOR LOCATION







Name

: MRS. VARAMAHALAXMI

Age / Gender Ref. By Dr.

: 53 years / Female

Reg. No.

: Dr. APOLO CLINIC

: 0604240031 C/o : Apollo Clinic UHID

: 0604240031

0604240031

**Bill Date** 

: 06-Apr-2024 08:57 AM Sample Col. Date: 06-Apr-2024 08:57 AM

**Result Date** 

: 06-Apr-2024 01:35 PM

Report Status : Final

**Test Name** 

Result

Unit

Reference Value

Method

Post Prandial Urine Sugar

Negative

Negative

Dipstick/Benedicts(Man



Printed By Printed On

: spectrum

: 06 Apr, 2024 05:02 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist



SCAN FOR LOCATION

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 +91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org





Age / Gender : 53 years / Female Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 0604240031

C/o : Apollo Clinic **Bill Date** 

: 0604240031

0604240031

: 06-Apr-2024 08:57 AM Sample Col. Date: 06-Apr-2024 08:57 AM

**Result Date** 

: 06-Apr-2024 01:35 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Urine Routine Examination- Physical Examination	<u>Urine</u>			
Colour Appearance Reaction (pH) Specific Gravity Biochemical Examination	Pale Yellow Clear 5.50 1.025		Pale Yellow Clear 5.0-7.5 1.000-1.030	Visual Visual Dipstick Dipstick
Albumin Glucose Bilirubin Ketone Bodies Urobilinogen Nitrite Microscopic Examination	Negative Negative Negative Negative Normal Negative		Negative Negative Negative Negative Normal Negative	Dipstick/Precipitation Dipstick/Benedicts Dipstick/Fouchets Dipstick/Rotheras Dipstick/Ehrlichs Dipstick
Pus Cells Epithelial Cells RBCs Easts Erystals Others	2-4 1-2 Absent Absent Absent	hpf hpf hpf	0.0-5.0 0.0-10.0 Absent Absent Absent Absent	Microscopy Microscopy Microscopy Microscopy Microscopy Microscopy

UHID

Comments: The kidneys help infiltration of the blood by eliminating waste out of the body through urine. They also regulate water in the body by conserving electrolytes, proteins, and other compounds. But due to some conditions and abnormalities in kidney function, the urine may encompass some abnormal constituents, which are not normally present. A complete urine examination helps in detecting such abnormal constituents in urine. Several disorders can be detected by identifying and measuring the levels of such substances. Blood cells, bilirubin, bacteria, pus cells, epithelial cells may be present in urine due to kidney disease or infection. Routine urine examination helps to diagnose kidney diseases, urinary tract infections,



Printed By

: spectrum

Printed On : 06 Apr, 2024 05:02 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 +91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostics.org





Name

: MRS. VARAMAHALAXMI

Age / Gender Ref. By Dr.

: 53 years / Female

Reg. No.

**Test Name** 

: Dr. APOLO CLINIC : 0604240031

C/o

: Apollo Clinic

UHID : 0604240031

> 0604240031

**Bill Date** 

: 06-Apr-2024 08:57 AM

Sample Col. Date: 06-Apr-2024 08:57 AM

**Result Date** Report Status

: 06-Apr-2024 02:04 PM : Final

Result

Unit

Reference Value

Method

Post prandial Blood Glucose (PPBS)-Plasma

117

mg/dL

70-140

Hexo Kinase

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula  $C_6H_{12}O_6$ . It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total .

Blood Group & Rh Typing-Whole Blood EDTA

**Blood Group** 

Rh Type

**Positive** 

Slide/Tube agglutination

Slide/Tube

agglutination

Note: Confirm by tube or gel method.

Comments: ABO blood group system, the classification of human blood based on the inherited properties of red blood cells (erythrocytes) asdetermined by the presence or absence of the antigens A and B, which are carried on the surface of the red cells. Persons may thus have type A, type



Printed By

: spectrum


Printed On


: 06 Apr, 2024 05:02 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 +91 77604 97644 | 080 2337 1555









Age / Gender : 53 years / Female Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 0604240031 C/o : Apollo Clinic UHID : 0604240031

0604240031

**Bill Date** : 06-Apr-2024 08:57 AM

Sample Col. Date: 06-Apr-2024 08:57 AM **Result Date** : 06-Apr-2024 02:41 PM

**Report Status** : Final

Test Name	Result	Unit	Reference Value	Method
KFT ( Kidney Function Tes Blood Urea Nitrogen (BUN) Serum		mg/dL	7.0-18.0	GLDH,Kinetic Assay
Creatinine-Serum	0.55	mg/dL	Male: 0.70-1.30	Modified
Uric Acid-Serum	3.44	mg/dL	Female: 0.55-1.02 Male: 3.50-7.20	kinetic Jaffe Uricase PAP
Sodium (Na+)-Serum	141.6	mmol/L	Female: 2.60-6.00 135.0-145.0	Ion-Selective
otassium (K+)-Serum	4.16	mmol/L	3.5 to 5.5	Electrodes (ISE) Ion-Selective Electrodes
hloride(Cl-)-Serum	106.00	mmol/L	96.0-108.0	(ISE) Ion-Selective Electrodes

Comments: Renal Function Test (RFT), also called kidney function tests, are a group of tests performed to evaluate the functions of the kidneys. The kidneys play a vital role in removing waste, toxins, and extra water from the body. They are responsible for maintaining a healthy balance of water, salts, and minerals such as calcium, sodium, potassium, and phosphorus. They are also essential for blood pressure control, maintenance of the body's pH balance, making red blood cell production hormones, and promoting bone health. Hence, keeping your kidneys healthy is essential for maintaining overall health. It helps diagnose inflammation, infection or damage in the kidneys. The test measures Uric Acid, Creatinine, BUN and electrolytes in the blood to determine the health of the kidneys. Risk factors for kidney dysfunction such as hypertension, diabetes, cardiovascular disease, obesity, elevated cholesterol or a family history of kidney disease. It may also be when has signs and symptoms of kidney disease, though in early stage often no noticeable symptoms are observed. Kidney panel is useful for general health screening; screening patients at risk of developing kidney disease; management of patients with known kidney disease. Estimated GFR is especially important in CKD patients CKD for monitoring, it helps to identify disease at early stage in those with risk factors for CKD (diabetes, hypertension, cardiovascular disease, and family history of kidney disease). Early recognition and intervention are important in slowing the progression of CKD and preventing its complications.



Printed By

: spectrum

Printed On

: 06 Apr, 2024 05:02 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 12 of 12





