

CERTIFICATE OF MEDICAL FITNESS

· ·
NAME: Mr Erachmappa. K.S
AGE/GENDER: H291M
HEIGHT: 156CM WEIGHT: 65Kg
IDENTIFICATION MARK:
BLOOD PRESSURE: 130 Go my 1 Hg
PULSE: FO me
CVS: RS:P Normal
ANY OTHER DISEASE DIAGNOSED IN THE PAST:
ALLERGIES, IF ANY:
LIST OF PRESCRIBED MEDICINES:
ANY OTHER REMARKS:
I Certify that I have carefully examined Mr/Mrs. <u>Evac Lineapparks</u> son/daughter of MI · K · N · Shivaling phino has signed in my presence. He/ she has no physical disease and is fit for employment.
Signature of Medical Officer
Place: Spectrum diagnostic & health Caus
Date: 93 03 24
Disclaimer: The patient has not been checked for COVID. This certificate does not relate to the

covid status of the patient examined





Dr. Ashok S Bsc., MBBS., D.O.M.S **Consultant Opthalmologist** KMC No: 31827

DATE: 23.38.18.

EYE EXAMINATION

NAME: Mg. E Ja China	12 AGE: 427	GENDER: F/M
	RIGHT EYE	LEFT EYE
Vision	Olep-2010	Elly NO
Vision With glass	asin	enim
Color Vision	Normal	Normal
Anterior segment examination	Normal	Normal
Fundus Examination	Normal	Normal
Any other abnormality	Nill	Nill
Diagnosis/ impression	Normal	Normal









NAME	AGE	GENDER
Mo Educhinnapla k-5	427×	Mole.

DENTAL EXAMINATION REPORT:

8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8
8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8

C: CAVITY -> NOTE.

M: MISSING - Nove.

O: OTHERS - Genobulilis

ADVISED:

CLEANING / SCALING / ROOTS PLANNING / FLOSSING & POLISHING / OTHERS

REMARKS:


SIGNATURE OF THE DENTAL SURGE

SEAL

DATE







RMS

SPECTRUM DIAGNOSTICS


Bangalore

Patient ID: 0263

Name: ERACHINNAPPA K S CR Number: 20240323143858 Registration Date: 23-Mar-2024 Age: 42

Gender : Male

Operator: spectrum diagnostics



X - Air Left	35	30	45	45	40	40	40	40	35	35	20
O - Air Right	40	35	40	40	35	35	30	25	30	30	25
> - Bone Left											
< - Bone Right											

	Average	High	Mid	Low
AIR Left	36.82 dB	32.50 dB	40.00 dB	38.75 dB
AIR Right	33.18 dB	27.50 dB	33.33 dB	38.75 dB

Clinical Notes:

Not Found





NAME	: MR.ERACHINNAPPA K S	DATE : 23/03/2024
AGE/SEX	: 42YEARS/MALE	REG NO: 2303240070
REF BY	: APOLO CLINIC	

CHEST PA VIEW

- Visualised lungs are clear.
- Bilateral hila appears normal .
- Cardia is normal in size
- No pleural effusion

IMPRESSION: No significant abnormality .

Transort

DR PRAVEEN B, DMRD, DNB **Consultant Radiologist**







PATIENT NAME	MR ERACHINNAPPA K S	ID NO	2303240070
AGE	42YEARS	SEX	MALE
REF BY	DR.APOLO CLINIC	DATE	23.03.2024

2D ECHO CARDIOGRAHIC STUDY

M-MODE

1 7 1	IVIODE
AORTA	22mm
LEFT ATRIUM	29mm
RIGHT VENTRICLE	20mm
LEFT VENTRICLE (DIASTOLE)	31mm
LEFT VENTRICLE(SYSTOLE)	27mm
VENTRICULAR SEPTUM (DIASTOLE)	10mm
VENTRICULAR SEPTUM (SYSTOLE)	11mm
POSTERIOR WALL (DIASTOLE)	09mm
POSTERIOR WALL (SYSTOLE)	11mm
FRACTIONAL SHORTENING	30%
EJECTION FRACTION	58%

DOPPLER / COLOUR FLOW

Mitral Valve Velocity: MVE- 0.94m/s MVA - 0.63m/s E/A-0.64

Tissue Doppler : e' (Septal) -10cm/s E/e'(Septal) -9

Velocity/ Gradient across the Pulmonic valve : 0.83m/s 3mmHg

Max. Velocity / Gradient across the Aortic valve: 1.19m/s 4mmHg

Velocity / Gradient across the Tricuspid valve : 1.87 m/s 19mmHg







PATIENT NAME	MR ERACHINNAPPA K S	ID NO	2303240070
AGE	42YEARS	SEX	MALE
REF BY	DR.APOLO CLINIC	DATE	23.03.2024

2D ECHO CARDIOGRAHIC STUDY

LEFT VENTRICLE	SIZE& THICKNESS	NORMAL
CONTRACTILITY	REGIONAL GLOBAL	NO RWMA

RIGHT VENTRICLE	:	NORMAL	
LEFT ATRIUM	:	NORMAL	
RIGHT ATRIUM	:	NORMAL	
MITRAL VALVE		NORMAL	
AORTIC VALVE	:	NORMAL	
PULMONARY VALVE	:	NORMAL	
TRICUSPID VALVE	;	NORMAL	
INTER ATRIAL SEPTUM	;	INTACT	
INTER VENTRICULAR SEPT	UM:	INTACT	
PERICARDIUM	:	NORMAL	
OTHERS	: -	· NIL	

IMPRESSION

- NO REGIONAL WALL MOTION ABNORMALITY PRESENT
- NORMAL VALVES AND DIMENSIONS
- NORMAL LV FUNCTION, LVEF- 58%
- TRIVIAL MR / TRIVIAL TR
- NO CLOT / VEGETATION / EFFUSION

DURGA V ECHO TECHNICIAN

The science of radiology is based upon interpretation of shadows of normal and abnormal tissue. This is neither complete nor accurate; hence, findings should always be interpreted in to the light of clinico-pathological correction.







NAME AND LAB NO MR ERACHINAPPA K		REG-40070
AGE & SEX	42 YRS	MALE
DATE AND AREA OF INTEREST	23.03.2024	ABDOMEN & PELVIS
REF BY	C/O APOLO CLINIC	

USG ABDOMEN AND PELVIS

LIVER:

Normal in size and shows diffuse increased echogenicity

No e/o IHBR dilatation. No evidence of focal lesion. Portal vein appears normal. CBD appears normal.

GALL BLADDER:

Partially distended .No obvious calculus in the visualised luminal portion.

SPLEEN:

Normal in size and echotexture. No e/o focal lesion.

PANCREAS:

Head and body appears normal . Tail obscured by bowel gas shadows .

RETROPERITONEUM:

Suboptimal visualised due to bowel gas

RIGHT KIDNEY:

Right kidney, is normal in size & echotexture.

No evidence of calculus/ hydronephrosis.

No solid lesions.

LEFT KIDNEY:

Left kidney, is normal in size & echotexture.

No evidence of calculus/ hydronephrosis.

No solid lesions.

URINARY BLADDER:

Well distended. No wall thickening/calculi.

PROSTATE:

Normal in size and echotexture.

No evidence of ascites/pleural effusion.

IMPRESSION:

> Grade I fatty liver.

Suggested clinical / lab correlation.

EEN B, DMRD, DNB CONSULTANT RADIOLOGIST









Age / Gender : 42 years / Male

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 2303240070

C/o : Apollo Clinic **Bill Date** : 23-Mar-2024 09:59 AM

Sample Col. Date: 23-Mar-2024 09:59 AM **Result Date** : 23-Mar-2024 03:14 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
LFT-Liver Function Test -Seru	ım			
Bilirubin Total-Serum	0.60	mg/dL	0.2-1.0	Caffeine
Bilirubin Direct-Serum	0.15	mg/dL	0.0-0.2	Benzoate Diazotised Sulphanilic
Bilirubin Indirect-Serum Aspartate Aminotransferase AST/SGOT)-Serum	0.45 28.00	mg/dL U/L	Male: 0.0 - 1.10 Male: 15.0 - 37.0	Acid Direct Measure UV with Pyridoxal - 5 -
Alanine Aminotransferase ALT/SGPT)-Serum	53.00	U/L	Male: 16.0 - 63.0	Phosphate UV with Pyridoxal - 5 -
alkaline Phosphatase (ALP)- erum	104.00	U/L	Male: 45.0 - 117.0	Phosphate PNPP,AMP- Buffer
rotein, Total-Serum	6.40	g/dL	6.40-8.20	Biuret/Endpoint-
lbumin-Serum	3.80	g/dL	Male: 3.40 - 5.50	With Blank Bromocresol
lobulin-Serum lbumin/Globulin Ratio-Serum	2.60 1.46	g/dL Ratio	2.0-3.50 0.80-2.0	Purple Calculated Calculated

UHID

: 2303240070

2303240070



Printed By

: spectrum


Printed On : 23 Mar, 2024 07:44 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengal կելել - 15-ի Գգի



info@spectrumdiagnostics.org









Age / Gender : 42 years / Male

Ref. By Dr. : Dr. APOLO CLINIC Reg. No.

C/o : Apollo Clinic

: 2303240070

Bill Date

: 23-Mar-2024 09:59 AM

Sample Col. Date: 23-Mar-2024 09:59 AM

Result Date

: 23-Mar-2024 03:14 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Lipid Profile-Serum				
Cholesterol Total-Serum	192.00	mg/dL	Male: 0.0 - 200	Cholesterol
Triglycerides-Serum	163.00	mg/dL	Male: 0.0 - 150	Oxidase/Peroxidase Lipase/Glycerol
High-density lipoprotein (HDL) Cholesterol-Serum	43.00	mg/dL	Male: 40.0 - 60.0	Dehydrogenase Accelerator/Selective
Non-HDL cholesterol-Serum Low-density lipoprotein (LDL) Cholesterol-Serum	149 136.0	mg/dL mg/dL	Male: 0.0 - 130 Male: 0.0 - 100.0	Detergent Calculated Cholesterol esterase and cholesterol
Very-low-density lipoprotein (VLDL) cholesterol-Serum	33	mg/dL	Male: 0.0 - 40	oxidase Calculated
Cholesterol/HDL Ratio-Serum	4.47	Ratio	Male: 0.0 - 5.0	Calculated

: 2303240070

2303240070

UHID

Interpretation:

Parameter	Desirable	In. d. W. Str.		
Total Cholesterol		Borderline High	High	Very High
	<200	200-239	>240	
Triglycerides	<150	150-199	200 400	
Non-HDL cholesterol	<130		200-499	>500
	<130	160-189	190-219	>220
Low-density lipoprotein (LDL) Cholesterol	<100	100-129	160-189	>190
			100-109	130

Comments: As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atherosclerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus, sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.



Printed By

: spectrum

Printed On


: 23 Mar, 2024 07:44 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalւթւկեր 2600գ10













Age / Gender : 42 years / Male

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 2303240070

C/o

: Apollo Clinic

Bill Date

: 23-Mar-2024 09:59 AM

Sample Col. Date: 23-Mar-2024 09:59 AM

Result Date : 23-Mar-2024 03:14 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method	
Fasting Blood Sugar (FBS)- Plasma	79	mg/dL	60.0-110.0	Hexo Kinase	**************************************

: 2303240070

UHID

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula C₆H₁₂O₆. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total

Glycosylated Haemoglobin (HbA1c)-Whole Blood EDTA

Glycosylated Haemoglobin (HbA1c)

5.60

%

Non diabetic adults:<5.7

HPLC

At risk (Prediabetes): 5.7 - 6.4

Diagnosing Diabetes :>= 6.5

Diabetes

Excellent Control: 6-7

Fair to good Control: 7-8 Unsatisfactory Control:8-10


Poor Control:>10

Estimated Average Glucose(eAG)

114.01

mg/dL

Calculated









Age / Gender : 42 years / Male

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. C/o : Apollo Clinic

: 2303240070

Bill Date

Result Date

: 23-Mar-2024 09:59 AM

Sample Col. Date: 23-Mar-2024 09:59 AM

Report Status

: 23-Mar-2024 03:14 PM : Final

Test Name

Result

Unit

UHID

Reference Value

: 2303240070

Method

Note: 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

Comments: HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.



Printed By

: spectrum

Printed On

: 23 Mar, 2024 07:44 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalալել - 45/բգր10

🧠 +91 77604 97644 | 080 2337 1555

info@spectrumdiagnostics.org







Age / Gender : 42 years / Male

Ref. By Dr. : Dr. APOLO CLINIC Reg. No.

: 2303240070

C/o : Apollo Clinic **Bill Date**

: 23-Mar-2024 09:59 AM

Sample Col. Date: 23-Mar-2024 09:59 AM

Result Date : 23-Mar-2024 03:14 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Fasting Urine Glucose-Urine	Negative		Negative	Dipstick/Benedicts (Manual)
Gamma-Glutamyl Transferase (GGT)-Serum	48.00	U/L	Male: 15.0-85.0	Other g-Glut-3-
			Female: 5.0-55.0	carboxy-4 nitro

: 2303240070

UHID

Comments: Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system, and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of serum GGT levels in detecting alcohol-induced liver disease. Elevated serum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.



Printed By

: spectrum

Printed On

: 23 Mar, 2024 07:44 pm

Dr. Nithun Reddy C, MD, Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalurage 56/94/2

+91 77604 97644 | 080 2337 1555

info@spectrumdiagnostics.org







Age / Gender : 42 years / Male

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 2303240070

C/o : Apollo Clinic **Bill Date**

: 23-Mar-2024 09:59 AM

Sample Col. Date: 23-Mar-2024 09:59 AM

Result Date

: 23-Mar-2024 03:14 PM

Report Status : Final

Test Name

Result

Positive

Unit

UHID

Reference Value

: 2303240070

2303240070

Method

Blood Group & Rh Typing-Whole Blood EDTA

Blood Group

Rh Type

Slide/Tube

agglutination

Slide/Tube

agglutination

Note: Confirm by tube or gel method.

Comments: ABO blood group system, the classification of human blood based on the inherited properties of red blood cells (erythrocytes) as determined by the presence or absence of the antigens A and B, which are carried on the surface of the red cells. Persons may thus have type A, type



Printed By

: spectrum

Printed On

: 23 Mar, 2024 07:44 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalւթթեց 5,600գ10



info@spectrumdiagnostics.org







Age / Gender : 42 years / Male Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 2303240070

C/o : Apollo Clinic **Bill Date**

: 23-Mar-2024 09:59 AM

Sample Col. Date: 23-Mar-2024 09:59 AM

Result Date

: 23-Mar-2024 03:14 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Kidney Function Test (KFT)-I Kidney Function Test (KFT) Serum	BUN,CREA,Ur -	ic Acid,Na,K,(Cl-Serum	
Blood Urea Nitrogen (BUN)	10.00	mg/dL	7.0-18.0	GLDH,Kinetic
Creatinine-Serum	0.71	mg/dL	Male: 0.70-1.30 Female: 0.55-1.02	Assay Modified
Uric Acid-Serum	3.80	mg/dL	Male: 3.50-7.20 Female: 2.60-6.0	kinetic Jaffe
Electrolytes				
Sodium (Na+)-Serum	140.8	mmol/L	135.0-145.0	ISE-Direct
Potassium (K+)-Serum Chloride (Cl-)-Serum	3.67 96.80	mmol/L mmol/L	3.50-5.50 96.0-108.0	ISE-Direct ISE-Direct

2303240070

: 2303240070

Comments: Renal Function Test (RFT), also called kidney function tests, are a group of tests performed to evaluate the functions of the kidneys. The kidneys play a vital role in removing waste, toxins, and extra water from the body. They are responsible for maintaining a healthy balance of water, salts, and minerals such as calcium, sodium, potassium, and phosphorus. They are also essential for blood pressure control, maintenance of the body's pH balance, making red blood cell production hormones, and promoting bone health. Hence, keeping your kidneys healthy is essential for maintaining overall health. It helps diagnose inflammation, infection or damage in the kidneys. The test measures Uric Acid, Creatinine, BUN and electrolytes in the blood to determine the health of the kidneys. Risk factors for kidney dysfunction such as hypertension, diabetes, cardiovascular disease, obesity, elevated cholesterol or a family history of kidney disease. It may also be when has signs and symptoms of kidney disease, though in early stage often no noticeable symptoms are observed. Kidney panel is useful for general health screening; screening patients at risk of developing kidney disease; management of patients with known kidney disease. Estimated GFR is especially important in CKD patients CKD for monitoring, it helps to identify disease at early stage in those with risk factors for CKD (diabetes, hypertension, cardiovascular disease, and family history of kidney disease). Early recognition and intervention are important in slowing the progression of CKD and preventing its complications.



Printed By

: spectrum

Printed On

: 23 Mar, 2024 07:44 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalயுழ் நடுவு 10



info@spectrumdiagnostics.org









Age / Gender : 42 years / Male

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 2303240070

C/o : Apollo Clinic **Bill Date**

: 23-Mar-2024 09:59 AM

Sample Col. Date: 23-Mar-2024 09:59 AM

Result Date : 23-Mar-2024 03:14 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method	
Prostate-Specific Antige Serum	en(PSA)-0.81	ng/mL	0.0-4.0	CLIA	

: 2303240070

Note: 1. This is a recommended test for detection of prostate cancer along with Digital Rectal Examination (DRE) in males above 50 years of age.

2. False negative / positive results are observed in patients receiving mouse monoclonal antibodies for diagnosis or therapy.

UHID

3. PSA levels may appear consistently elevated / depressed due to the interference by heterophilic antibodies & nonspecific protein binding.

4. Immediate PSA testing following digital rectal examination, ejaculation, prostatic massage, indwelling catheterization, ultrasonography and needle biopsy of prostate is not recommended as they falsely elevate levels

5. PSA values regardless of levels should not be interpreted as absolute evidence of the presence or absence of disease. All values should be

clinical findings and results of other investigations

6. Sites of Non-prostatic PSA production are breast epithelium, salivary glands, periurethral & anal glands, cells of male urethra & breast milk

7. Physiological decrease in PSA level by 18% has been observed in hospitalized /sedentary patients either due to supine position or suspended sexual activity.

Recommended Testing Intervals: Pre-operatively (Baseline), 2-4 days post-operatively, Prior to discharge from hospital, Monthly followup if levels are

Clinical Use: -An aid in the early detection of Prostate cancer when used in conjunction with Digital rectal examination in males more than 50 years of age and in those with two or more affected first degree relatives.

-Followup and management of Prostate cancer patients

-Detect metastatic or persistent disease in patients following surgical or medical treatment of Prostate cancer.

Increased Levels: Prostate cancer, Benign Prostatic Hyperplasia, Prostatitis, Genitourinary infections.



Printed By

: spectrum

Printed On

: 23 Mar, 2024 07:44 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalupկခွင့် နှင့်ဝှာညှှု၊၀



info@spectrumdiagnostics.org









Age / Gender : 42 years / Male

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 2303240070

C/o

: Apollo Clinic

Bill Date : 23-Mar-2024 09:59 AM

Sample Col. Date: 23-Mar-2024 09:59 AM **Result Date** : 23-Mar-2024 03:14 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Thyroid function tests (TF Serum	T)-			
Tri-Iodo Thyronine (T3)-So	erum 0.77	ng/mL	Male: 0.60 - 1.81	Chemiluminescence Immunoassay
Thyroxine (T4)-Serum	5.56	μg/dL	Male: 5.50 - 12.10	(CLIA) Chemiluminescence
Thyroid Stimulating Hormo TSH)-Serum	one 3.16	μIU/mL	Male: 0.35 - 5.50	Immunoassay (CLIA) Chemiluminescence Immunoassay (CLIA)

2303240070

: 2303240070

Comments: Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population without

Reference range: Cord: (37 Weeks): 0.5-1.41, Children:1-3 Days: 1.0-7.40,1-11 Months: 1.05-2.45,1-5 Years: 1.05-2.69,6-10 Years: 0.94-2.41,11-15

Reference range: Adults: 20-50 Years; 0.70-2.04, 50-90 Years: 0.40-1.81,

Reference range in Pregnancy: First Trimester: 0.81-1.90, Second Trimester: 1.0-2.60

Increased Levels: Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG). Decreased Levels: Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

Comments: Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4). It also helps to monitor treatment of Hyperthyroidism with

Reference Range: Males: 4.6-10.5, Females: 5.5-11.0, 60 Years: 5.0-10.70, Cord: 7.40-13.10, Children: 1-3 Days: 11.80-22.60, 1-2 Weeks: 9.90-16.60,1-4 Months: 7.20-14.40,1-5 Years: 7.30-15.0,5-10 Years: 6.4-13.3

1-15 Years: 5.60-11.70, Newborn Screen: 1-5 Days: >7.5,6 Days :>6.5

Increased Levels: Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia, Increased transthyretin, estrogen therapy, pregnancy. Decreased Levels: Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.

Comments: TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, caloric intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after ~ 3 weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality.

Reference range in Pregnancy: I- trimester:0.1-2.5; II -trimester:0.2-3.0; III- trimester:0.3-3.0

Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks:1.7-9.1

Increased Levels: Primary hypothyroidism, Subclinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance. Decreased Levels: Graves disease, Autonomous thyroid hormone secretion, TSH deficiency.

Printed By : spectrum

Printed On : 23 Mar, 2024 07:44 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalurակը 5 6 የ በ













Age / Gender : 42 years / Male

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 2303240070

C/o : Apollo Clinic **Bill Date** : 23-Mar-2024 09:59 AM

Sample Col. Date: 23-Mar-2024 09:59 AM **Result Date** : 23-Mar-2024 03:14 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Urine Routine Examination	-Urine			
Physical Examination				
Colour	Pale Yellow		Pale Yellow	
Appearance	Clear		Clear	Visual
Reaction (pH)	6.00		5.0-7.5	Visual
Specific Gravity	1.025			Dipstick
Biochemical Examination			1.000-1.030	Dipstick
Albumin	Negative		Manatin	
Glucose	Negative		Negative	Dipstick/Precipitation
Bilirubin	Negative		Negative	Dipstick/Benedicts
Ketone Bodies	Negative		Negative	Dipstick/Fouchets
Urobilinogen	Normal		Negative	Dipstick/Rotheras
Vitrite	Negative		Normal	Dipstick/Ehrlichs
Aicroscopic Examination	rogative		Negative	Dipstick
Pus Cells	2-4	ha f	0.0.7.0	
Cpithelial Cells	1-2	hpf	0.0-5.0	Microscopy
BCs	Absent	hpf	0.0-10.0	Microscopy
Casts	Absent	hpf	Absent	Microscopy
rystals	Absent		Absent	Microscopy
others			Absent	Microscopy
	Absent		Absent	Microscopy

UHID

: 2303240070

2303240070

Comments: The kidneys help infiltration of the blood by eliminating waste out of the body through urine. They also regulate water in the body by conserving electrolytes, proteins, and other compounds. But due to some conditions and abnormalities in kidney function, the urine may encompass some abnormal constituents, which are not normally present. A complete urine examination helps in detecting such abnormal constituents in urine. Several disorders can be detected by identifying and measuring the levels of such substances. Blood cells, bilirubin, bacteria, pus cells, epithelial cells may be present in urine due to kidney disease or infection. Routine urine examination helps to diagnose kidney diseases, urinary tract infections,



Printed By

: spectrum

Printed On


: 23 Mar, 2024 07:44 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalម្ពុផ្តេច 156pអ្ន0

+91 77604 97644 | 080 2337 1555











Age / Gender : 42 years / Male

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 2303240070

C/o : Apollo Clinic : 2303240070

Result Date

2303240070

Bill Date : 23-Mar-2024 09:59 AM

Sample Col. Date: 23-Mar-2024 09:59 AM : 23-Mar-2024 03:47 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Calcium, Total-Serum	8.80	mg/dL	8.50-10.10	Spectrophotometry (O-
				Cresolphthalein complexone)

UHID



Printed By

: spectrum

Printed On : 23 Mar, 2024 07:44 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalարդը 156թին +91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostics.org









Age / Gender : 42 years / Male

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 2303240070

C/o

: Apollo Clinic

Bill Date

: 23-Mar-2024 09:59 AM

Sample Col. Date: 23-Mar-2024 09:59 AM

Result Date : 23-Mar-2024 05:21 PM

Report Status

: Final

Test Name Result Unit Reference Value Method Postprandial Urine glucose-Negative Negative Dipstick/Benedicts Urine (Manual)

2303240070

: 2303240070

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

UHID

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total

Post prandial Blood Glucose (PPBS)-Plasma

130

mg/dL

70-140

Hexo Kinase

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula $C_6H_{12}O_6$. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total



Printed By

: spectrum

Printed On

: 23 Mar, 2024 07:44 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalម្ពុជ្យ-156pណ្ដ



info@spectrumdiagnostics.org









Age / Gender : 42 years / Male

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 2303240070

C/o : Apollo Clinic

Bill Date : 23-Mar-2024 09:59 AM Sample Col. Date: 23-Mar-2024 09:59 AM : 2303240070

Result Date : 23-Mar-2024 05:21 PM 2303240070

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Complete Haemogram-Whole	Blood EDTA			
Haemoglobin (HB)	12.80	g/dL	Male: 14.0-17.0 Female:12.0-15.0 Newborn:16.50 - 19.50	Spectrophotmeter
Red Blood Cell (RBC)	3.96	million/cu	mm3.50 - 5.50	Volumetric Impedance
Packed Cell Volume (PCV)	36.30	%	Male: 42.0-51.0 Female: 36.0-45.0	Electronic Pulse
Mean corpuscular volume (MCV)	91.50	fL	78.0- 94.0	Calculated
Mean corpuscular hemoglobin (MCH)		pg	27.50-32.20	Calculated
Mean corpuscular hemoglobin concentration (MCHC)	35.20	%	33.00-35.50	Calculated
Red Blood Cell Distribution Width SD (RDW-SD)	39.20	fL	40.0-55.0	Volumetric
Red Blood Cell Distribution CV (RDW-CV)	14.00	%	Male: 11.80-14.50 Female: 12.20-16.10	Impedance Volumetric
Mean Platelet Volume (MPV)	9.70	fL	8.0-15.0	Impedance Volumetric
Platelet	2.25	lakh/cumm	1.50-4.50	Impedance Volumetric
latelet Distribution Width PDW)	10.50	%	8.30 - 56.60	Impedance Volumetric
White Blood cell Count (WBC)		cells/cumm	Male: 4000-11000 Female 4000-11000 Children: 6000-17500 Infants: 9000-30000	Impedance Volumetric Impedance
eutrophils	51.20	%	40.0-75.0	Light
ymphocytes	41.20	%	20.0-40.0	scattering/Manual Light
osinophils	3.90	%	0.8-0.0	scattering/Manual Light scattering/Manual

UHID









Age / Gender : 42 years / Male

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 2303240070

C/o : Apollo Clinic **Bill Date**

: 23-Mar-2024 09:59 AM

Sample Col. Date: 23-Mar-2024 09:59 AM

Result Date

: 23-Mar-2024 05:21 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Monocytes	3.70	%	0.0-10.0	Light scattering/Manual
Basophils	0.00	%	0.0-1.0	Light scattering/Manual
Absolute Neutrophil Count	3.10	10^3/uL	2.0-7.0	Calculated
Absolute Lymphocyte Count	2.49	10^3/uL	1.0-3.0	Calculated
Absolute Monocyte Count	0.22	10^3/uL	0.20-1.00	Calculated
Absolute Eosinophil Count	240.00	cells/cumm	40-440	Calculated
Absolute Basophil Count	0.00	10^3/uL	0.0-0.10	Calculated
Erythrocyte Sedimentation Rate (ESR)	03	mm/hr	Female: 0.0-20.0 Male: 0.0-10.0	Westergren

2303240070

: 2303240070

UHID

Peripheral Smear Examination-Whole Blood EDTA

Method: (Microscopy-Manual)

RBC'S : Normocytic Normochromic.

WBC'S : Are normal in total number, morphology and distribution.

: Adequate in number and normal in morphology. Platelets

No abnormal cells or hemoparasites are present.

Impression: Normocytic Normochromic Blood picture.



Printed By

: spectrum

Printed On

: 23 Mar, 2024 07:44 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalur🌬 🛱 🖓 🗛



info@spectrumdiagnostics.org

