

PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON W

ACCESSION NO : 0002XD020787

PATIENT ID : AMITM1910872

CLIENT PATIENT ID:

ABHA NO :

AGE/SEX : 37 Years Male

DRAWN : 13/04/2024 08:48:36

RECEIVED : 13/04/2024 08:50:04

REPORTED : 17/04/2024 17:27:55

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE**XRAY-CHEST**

IMPRESSION NO ABNORMALITY DETECTED

ECG

ECG WITHIN NORMAL LIMITS

ANTHROPOMETRIC DATA & BMI

HEIGHT IN METERS	1.71	mts
WEIGHT IN KGS.	86.8	Kgs
BMI	30	kg/sqmts

BMI & Weight Status as follows:
 Below 18.5: Underweight
 18.5 - 24.9: Normal
 25.0 - 29.9: Overweight
 30.0 and Above: Obese

GENERAL EXAMINATION

PULSE 72/MIN REGULAR, ALL PERIPHERAL PULSES WELL FELT, NO CAROTID BRUIT

CARDIOVASCULAR SYSTEM


BP 120/70MM HG (SUPINE) mm/Hg

BASIC EYE EXAMINATION

DISTANT VISION RIGHT EYE WITHOUT GLASSES WITHIN NORMAL LIMIT (6/6)



Dr. Swati Karmarkar,
MD,DNB,DMRD
Consultant Radiologist



Dr. J N Shukla ,MBBS, AFIH
Consultant Physician

Page 1 Of 21



View Details



View Report

PERFORMED AT :

Agilus Diagnostics Ltd
 Prime Square Building,Plot No 1,Gaiwadi Industrial Estate,S.V. Road,Goregaon (W)
 Mumbai, 400062
 Maharashtra, India
 Tel : 9111591115, Fax :
 CIN - U74899PB1995PLC045956



ULR No.2000012351057-0002

PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON WACCESSION NO : **0002XD020787**

PATIENT ID : AMITM1910872

CLIENT PATIENT ID:

ABHA NO :

AGE/SEX : 37 Years Male

DRAWN : 13/04/2024 08:48:36

RECEIVED : 13/04/2024 08:50:04

REPORTED : 17/04/2024 17:27:55

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------


DISTANT VISION LEFT EYE WITHOUT GLASSES	WITHIN NORMAL LIMIT (6/6)
NEAR VISION RIGHT EYE WITHOUT GLASSES	WITHIN NORMAL LIMIT (N6)
NEAR VISION LEFT EYE WITHOUT GLASSES	WITHIN NORMAL LIMIT (N6)
COLOUR VISION	NORMAL (17/17)

SUMMARY

RELEVANT HISTORY	NOT SIGNIFICANT
RELEVANT GP EXAMINATION FINDINGS	OVERWEIGHT
RELEVANT LAB INVESTIGATIONS	RAISED BILIRUBIN DIRECT (0.60)
RELEVANT NON PATHOLOGY DIAGNOSTICS	USG : GRADE I FATTY LIVER
REMARKS / RECOMMENDATIONS	LOW CALORIC DIET AND REGULAR PHYSICAL EXERCISE. ADV. TO FOLLOW UP WITH PHYSICIAN FOR 1. RAISED BILIRUBIN DIRECT 2. FATTY LIVER.



Dr. Swati Karmarkar,
MD,DNB,DMRD
Consultant Radiologist



Dr. J N Shukla ,MBBS, AFIH
Consultant Physician

Page 2 Of 21



View Details



View Report

PERFORMED AT :

Agilus Diagnostics Ltd
Prime Square Building,Plot No 1,Gaiwadi Industrial Estate,S.V. Road,Goregaon (W)
Mumbai, 400062
Maharashtra, India
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956



ULR No.2000012351057-0002

PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON WACCESSION NO : **0002XD020787**

PATIENT ID : AMITM1910872

CLIENT PATIENT ID:

ABHA NO :

AGE/SEX : 37 Years Male

DRAWN : 13/04/2024 08:48:36

RECEIVED : 13/04/2024 08:50:04

REPORTED : 17/04/2024 17:27:55

Test Report Status **Final**

Results

Units

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE**ULTRASOUND ABDOMEN****ULTRASOUND ABDOMEN**

GRADE I FATTY LIVER.


TMT OR ECHO**CLINICAL PROFILE**

2 DECHO DONE IMPRESSION

- Good LV systolic function at rest. No RWMA.
- LVEF 55-60%.
- All valves structurally normal.
- No evidence of PE/clot/vegetation.



Dr. Swati Karmarkar,
MD,DNB,DMRD
Consultant Radiologist



Dr. J N Shukla ,MBBS, AFIH
Consultant Physician

Page 3 Of 21



View Details



View Report

PERFORMED AT :**Agilus Diagnostics Ltd**

Prime Square Building,Plot No 1,Gaiwadi Industrial Estate,S.V. Road,Goregaon (W)

Mumbai, 400062

Maharashtra, India

Tel : 9111591115, Fax :

CIN - U74899PB1995PLC045956



ULR No.2000012351057-0002

PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON W

ACCESSION NO : 0002XD020787

PATIENT ID : AMITM1910872

CLIENT PATIENT ID:

ABHA NO :

AGE/SEX : 37 Years Male

DRAWN : 13/04/2024 08:48:36

RECEIVED : 13/04/2024 08:50:04

REPORTED : 17/04/2024 17:27:55

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

HAEMATOLOGY - CBC

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

BLOOD COUNTS, EDTA WHOLE BLOOD

HEMOGLOBIN (HB)	14.3	13.0 - 17.0	g/dL
METHOD : CYANIDE FREE DETERMINATION			
RED BLOOD CELL (RBC) COUNT	5.38	4.5 - 5.5	mil/ μ L
METHOD : FLUORESCENCE FLOW CYTOMETRY			
WHITE BLOOD CELL (WBC) COUNT	7.46	4.0 - 10.0	thou/ μ L
METHOD : ELECTRICAL IMPEDANCE			
PLATELET COUNT	307	150 - 410	thou/ μ L
METHOD : ELECTRONIC IMPEDENCE & MICROSCOPY			

RBC AND PLATELET INDICES

HEMATOCRIT (PCV)	45.6	40 - 50	%
METHOD : CALCULATED PARAMETER			
MEAN CORPUSCULAR VOLUME (MCV)	84.7	83.0 - 101.0	fL
METHOD : DERIVED PARAMETER FROM RBC HISTOGRAM			
MEAN CORPUSCULAR HEMOGLOBIN (MCH)	26.6 Low	27.0 - 32.0	pg
METHOD : CALCULATED PARAMETER			
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION (MCHC)	31.4 Low	31.5 - 34.5	g/dL
METHOD : CALCULATED PARAMETER			
RED CELL DISTRIBUTION WIDTH (RDW)	14.3 High	11.6 - 14.0	%
METHOD : DERIVED PARAMETER FROM RBC HISTOGRAM			
MENTZER INDEX	15.7		
MEAN PLATELET VOLUME (MPV)	9.2	6.8 - 10.9	fL
METHOD : DERIVED PARAMETER FROM PLATELET HISTOGRAM			

WBC DIFFERENTIAL COUNT

NEUTROPHILS	64	40 - 80	%
METHOD : FLUORESCENCE FLOW CYTOMETRY			
LYMPHOCYTES	29	20 - 40	%
METHOD : FLUORESCENCE FLOW CYTOMETRY			
MONOCYTES	5	2 - 10	%


Dr. Sushant Chikane
Consultant Pathologist

Dr. Akshaya Mandloi, MD, PDF
Consultant Hematopathologist
and Head,
Dept. of Hematopathology and
Flowcytometry

Page 4 Of 21



View Details



View Report

PERFORMED AT :

Agilus Diagnostics Ltd
Prime Square Building, Plot No 1, Gaiwadi Industrial Estate, S.V. Road, Goregaon (W)
Mumbai, 400062
Maharashtra, India
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956



ULR No. 2000012351057-0002



PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON W

ACCESSION NO : **0002XD020787**
 PATIENT ID : AMITM1910872
 CLIENT PATIENT ID:
 ABHA NO :

AGE/SEX : 37 Years Male
 DRAWN : 13/04/2024 08:48:36
 RECEIVED : 13/04/2024 08:50:04
 REPORTED : 17/04/2024 17:27:55

Test Report Status	Final	Results	Biological Reference Interval	Units
METHOD : FLUORESCENCE FLOW CYTOMETRY				
EOSINOPHILS		2	1 - 6	%
METHOD : FLUORESCENCE FLOW CYTOMETRY				
BASOPHILS		0	0 - 1	%
METHOD : FLUORESCENCE FLOW CYTOMETRY				
ABSOLUTE NEUTROPHIL COUNT		4.77	2.0 - 7.0	thou/ μ L
METHOD : CALCULATED PARAMETER				
ABSOLUTE LYMPHOCYTE COUNT		2.16	1.0 - 3.0	thou/ μ L
METHOD : CALCULATED PARAMETER				
ABSOLUTE MONOCYTE COUNT		0.37	0.2 - 1.0	thou/ μ L
METHOD : CALCULATED PARAMETER				
ABSOLUTE EOSINOPHIL COUNT		0.15	0.02 - 0.50	thou/ μ L
METHOD : CALCULATED PARAMETER				
ABSOLUTE BASOPHIL COUNT		0.00 Low	0.02 - 0.10	thou/ μ L
METHOD : CALCULATED PARAMETER				
NEUTROPHIL LYMPHOCYTE RATIO (NLR)		2.2		
METHOD : CALCULATED				

Interpretation(s)

BLOOD COUNTS,EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.
 RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait (<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.
 WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.
 (Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients ; A.-P. Yang, et al.; International Immunopharmacology 84 (2020) 106504
 This ratio element is a calculated parameter and out of NABL scope.

Dr. Sushant Chikane
 Consultant Pathologist

Dr. Akshaya Mandloi,MD,PDF
 Consultant Hematopathologist
 and Head,
 Dept.of Hematopathology and
 Flowcytometry



View Details



View Report

PERFORMED AT :

Agilus Diagnostics Ltd
 Prime Square Building,Plot No 1,Gaiwadi Industrial Estate,S.V. Road,Goregaon (W)
 Mumbai, 400062
 Maharashtra, India
 Tel : 9111591115, Fax :
 CIN - U74899PB1995PLC045956



PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON W

ACCESSION NO : 0002XD020787

PATIENT ID : AMITM1910872

CLIENT PATIENT ID:

ABHA NO :

AGE/SEX : 37 Years Male

DRAWN : 13/04/2024 08:48:36

RECEIVED : 13/04/2024 08:50:04

REPORTED : 17/04/2024 17:27:55

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

HAEMATOLOGY

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

ERYTHROCYTE SEDIMENTATION RATE (ESR), EDTA BLOOD

E.S.R 8 = or < 10 mm at 1 hr

METHOD : MODIFIED WESTERGREN METHOD BY AUTOMATED ANALYSER

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD

HBA1C	5.6	Non-diabetic Adult < 5.7 % Pre-diabetes 5.7 - 6.4 Diabetes diagnosis: > or = 6.5 Therapeutic goals: < 7.0 Action suggested : > 8.0 (ADA Guideline 2021)
ESTIMATED AVERAGE GLUCOSE(EAG)	114.0	< 116 mg/dL

METHOD : ION- EXCHANGE HPLC

Interpretation(s)

ERYTHROCYTE SEDIMENTATION RATE (ESR), EDTA BLOOD-TEST DESCRIPTION :-

Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic; it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition. CRP is superior to ESR because it is more sensitive and reflects a more rapid change.

TEST INTERPRETATION

Increase in: Infections, Vasculitides, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy, Estrogen medication, Aging.

Finding a very accelerated ESR (>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias, Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis).

In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum.

Decreased in: Polycythemia vera, Sickle cell anemia

LIMITATIONS

False elevated ESR : Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia

False Decreased : Poikilocytosis, (Sickle Cells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs(Quinine, salicylates)

REFERENCE :

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition; 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin; 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th edition.

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-Used For:



Dr. Akshaya Mandloi, MD, PDF
Consultant Hematopathologist
and Head,
Dept. of Hematopathology and
Flowcytometry



Dr. Sushant Chikane
Consultant Pathologist

Page 6 Of 21



View Details



View Report

PERFORMED AT :

Agilus Diagnostics Ltd

Prime Square Building, Plot No 1, Gaiwadi Industrial Estate, S.V. Road, Goregaon (W)
Mumbai, 400062

Maharashtra, India

Tel : 9111591115, Fax :

CIN - U74899PB1995PLC045956



ULN No. 2000012351057-0002



PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON W

ACCESSION NO : **0002XD020787**
PATIENT ID : AMITM1910872
CLIENT PATIENT ID:
ABHA NO :

AGE/SEX : 37 Years Male
DRAWN : 13/04/2024 08:48:36
RECEIVED : 13/04/2024 08:50:04
REPORTED : 17/04/2024 17:27:55

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

1. Evaluating the long-term control of blood glucose concentrations in diabetic patients.
2. Diagnosing diabetes.
3. Identifying patients at increased risk for diabetes (prediabetes).

The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for well-controlled type 2 diabetic patients) to determine whether a patients metabolic control has remained continuously within the target range.

1. eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.
2. eAG gives an evaluation of blood glucose levels for the last couple of months.
3. eAG is calculated as $eAG (mg/dl) = 28.7 * HbA1c - 46.7$

HbA1c Estimation can get affected due to :

1. Shortened Erythrocyte survival : Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss,hemolytic anemia) will falsely lower HbA1c test results.Fructosamine is recommended in these patients which indicates diabetes control over 15 days.
- 2.Vitamin C & E are reported to falsely lower test results.(possibly by inhibiting glycation of hemoglobin.
3. Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia,uremia, hyperbilirubinemia, chronic alcoholism,chronic ingestion of salicylates & opiates addition are reported to interfere with some assay methods,falsely increasing results.
4. Interference of hemoglobinopathies in HbA1c estimation is seen in

- a) Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c.
- b) Heterozygous state detected (D10 is corrected for HbS & HbC trait.)
- c) HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c.Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy

Dr. Akshaya Mandloi,MD,PDF
Consultant Hematopathologist
and Head,
Dept.of Hematopathology and
Flowcytometry

Dr. Sushant Chikane
Consultant Pathologist



View Details



View Report

PERFORMED AT :

Agilus Diagnostics Ltd
Prime Square Building,Plot No 1,Gaiwadi Industrial Estate,S.V. Road,Goregaon (W)
Mumbai, 400062
Maharashtra, India
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956



ULR No.2000012351057-0002

PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON WACCESSION NO : **0002XD020787**

PATIENT ID : AMITM1910872

CLIENT PATIENT ID:

ABHA NO :

AGE/SEX : 37 Years Male

DRAWN : 13/04/2024 08:48:36

RECEIVED : 13/04/2024 08:50:04

REPORTED : 17/04/2024 17:27:55

Test Report Status **Final**

Results

Biological Reference Interval Units

IMMUNOHAEMATOLOGY

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE**ABO GROUP & RH TYPE, EDTA WHOLE BLOOD**

ABO GROUP

A

METHOD : HAEMAGGLUTINATION (AUTOMATED)

RH TYPE

POSITIVE

METHOD : HAEMAGGLUTINATION (AUTOMATED)

Interpretation(s)

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same."

The test is performed by both forward as well as reverse grouping methods.



Dr. Sushant Chikane
Consultant Pathologist



Dr. Akshaya Mandloi, MD, PDF
Consultant Hematopathologist
and Head,
Dept. of Hematopathology and
Flowcytometry

Page 8 Of 21



View Details



View Report

PERFORMED AT :**Agilus Diagnostics Ltd**

Prime Square Building, Plot No 1, Gaiwadi Industrial Estate, S.V. Road, Goregaon (W)

Mumbai, 400062

Maharashtra, India

Tel : 9111591115, Fax :

CIN - U74899PB1995PLC045956



ULR No.2000012351057-0002



PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON W

ACCESSION NO : **0002XD020787**
PATIENT ID : AMITM1910872
CLIENT PATIENT ID:
ABHA NO :

AGE/SEX : 37 Years Male
DRAWN : 13/04/2024 08:48:36
RECEIVED : 13/04/2024 08:50:04
REPORTED : 17/04/2024 17:27:55

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

BIOCHEMISTRY

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

GLUCOSE FASTING, FLUORIDE PLASMA

FBS (FASTING BLOOD SUGAR)	88	Normal <100 mg/dL Impaired fasting glucose: 100 to 125 Diabetes mellitus: > = 126 (on more than 1 occasion) (ADA guidelines 2021)
---------------------------	----	--

METHOD : SPECTROPHOTOMETRY HEXOKINASE

GLUCOSE, POST-PRANDIAL, PLASMA

PPBS (POST PRANDIAL BLOOD SUGAR)	100	Normal <140 mg/dL Impaired glucose tolerance: 140 to 199 Diabetes mellitus : > = 200 (on more than 1 occasion) ADA guideline 2021
----------------------------------	-----	--

METHOD : SPECTROPHOTOMETRY HEXOKINASE

LIPID PROFILE WITH CALCULATED LDL, SERUM

CHOLESTEROL, TOTAL	168	Desirable : < 200 mg/dL Borderline : 200 - 239 High : > / = 240
--------------------	-----	---

METHOD : SPECTROPHOTOMETRY, ENZYMATIC COLORIMETRIC - CHOLESTEROL OXIDASE, ESTERASE, PEROXIDASE

TRIGLYCERIDES	104	Normal: < 150 mg/dL Borderline high: 150 - 199 High: 200 - 499 Very High: > / = 500
---------------	-----	--

METHOD : SPECTROPHOTOMETRY, ENZYMATIC ENDPOINT WITH GLYCEROL BLANK

HDL CHOLESTEROL	40	At Risk: < 40 mg/dL Desirable: > or = 60
-----------------	----	---

METHOD : SPECTROPHOTOMETRY, HOMOGENEOUS DIRECT ENZYMATIC COLORIMETRIC

Dr. Apeksha Sharma
D.P.B., DNB (PATH)
(Reg.no.MMC2008/06/2561)
Consultant Pathologist

Dr. Deepak Sanghavi, M.D (Path)
(Reg.no.MMC2004/03/1530)
Chief of Mumbai Reference Lab.



View Details



View Report

PERFORMED AT :

Agilus Diagnostics Ltd
Prime Square Building, Plot No 1, Gaiwadi Industrial Estate, S.V. Road, Goregaon (W)
Mumbai, 400062
Maharashtra, India
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956





PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON W

ACCESSION NO : **0002XD020787**
PATIENT ID : AMITM1910872
CLIENT PATIENT ID:
ABHA NO :

AGE/SEX : 37 Years Male
DRAWN : 13/04/2024 08:48:36
RECEIVED : 13/04/2024 08:50:04
REPORTED : 17/04/2024 17:27:55

Test Report Status	Final	Results	Biological Reference Interval	Units
CHOLESTEROL LDL		107 High	Optimal : < 100 Near optimal/above optimal : 100-129 Borderline high : 130-159 High : 160-189 Very high : = 190	mg/dL
METHOD : CALCULATED PARAMETER				
NON HDL CHOLESTEROL		128	Desirable : < 130 Above Desirable : 130 -159 Borderline High : 160 - 189 High : 190 - 219 Very high : > / = 220	mg/dL
METHOD : CALCULATED PARAMETER				
VERY LOW DENSITY LIPOPROTEIN		20.8	< or = 30.0	mg/dL
METHOD : CALCULATED PARAMETER				
CHOL/HDL RATIO		4.2	Low Risk : 3.3 - 4.4 Average Risk : 4.5 - 7.0 Moderate Risk : 7.1 - 11.0 High Risk : > 11.0	
METHOD : CALCULATED PARAMETER				
LDL/HDL RATIO		2.7	Desirable/Low Risk : 0.5 - 3.0 Borderline/Moderate Risk : 3.1 - 6.0 High Risk : > 6.0	
METHOD : CALCULATED PARAMETER				

Interpretation(s)

Serum lipid profile is measured for cardiovascular risk prediction. Lipid Association of India recommends LDL-C as primary target and Non HDL-C as co-primary treatment target.

Risk Stratification for ASCVD (Atherosclerotic cardiovascular disease) by Lipid Association of India

Risk Category	
Extreme risk group	A.CAD with > 1 feature of high risk group B. CAD with > 1 feature of Very high risk group or recurrent ACS (within 1 year) despite LDL-C < or = 50 mg/dl or polyvascular disease
Very High Risk	1. Established ASCVD 2. Diabetes with 2 major risk factors or evidence of end organ damage 3. Familial Homozygous Hypercholesterolemia
High Risk	1. Three major ASCVD risk factors. 2. Diabetes with 1 major risk factor or no evidence of end organ damage. 3. CKD stage 3B or 4. 4. LDL >190 mg/dl 5. Extreme of a single risk factor. 6. Coronary Artery Calcium - CAC >300 AU. 7. Lipoprotein a >= 50mg/dl 8. Non stenotic carotid plaque

Dr. Apeksha Sharma
D.P.B.,DNB (PATH)
(Reg.no.MMC2008/06/2561)
Consultant Pathologist

Dr. Deepak Sanghavi, M.D(Path)
(Reg.no.MMC2004/03/1530)
Chief of Mumbai Reference Lab.



View Details



View Report

PERFORMED AT :

Agilus Diagnostics Ltd
Prime Square Building, Plot No 1, Gaiwadi Industrial Estate, S.V. Road, Goregaon (W)
Mumbai, 400062
Maharashtra, India
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956





PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON W

ACCESSION NO : **0002XD020787**
PATIENT ID : AMITM1910872
CLIENT PATIENT ID :
ABHA NO :

AGE/SEX : 37 Years Male
DRAWN : 13/04/2024 08:48:36
RECEIVED : 13/04/2024 08:50:04
REPORTED : 17/04/2024 17:27:55

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

Moderate Risk	2 major ASCVD risk factors
Low Risk	0-1 major ASCVD risk factors
Major ASCVD (Atherosclerotic cardiovascular disease) Risk Factors	
1. Age > or = 45 years in males and > or = 55 years in females	3. Current Cigarette smoking or tobacco use
2. Family history of premature ASCVD	4. High blood pressure
5. Low HDL	

Newer treatment goals and statin initiation thresholds based on the risk categories proposed by LAI in 2020.

Risk Group	Treatment Goals		Consider Drug Therapy	
	LDL-C (mg/dl)	Non-HDL (mg/dl)	LDL-C (mg/dl)	Non-HDL (mg/dl)
Extreme Risk Group Category A	<50 (Optional goal < OR = 30)	< 80 (Optional goal <OR = 60)	>OR = 50	>OR = 80
Extreme Risk Group Category B	<OR = 30	<OR = 60	> 30	>60
Very High Risk	<50	<80	>OR= 50	>OR= 80
High Risk	<70	<100	>OR= 70	>OR= 100
Moderate Risk	<100	<130	>OR= 100	>OR= 130
Low Risk	<100	<130	>OR= 130*	>OR= 160

*After an adequate non-pharmacological intervention for at least 3 months.

References: Management of Dyslipidaemia for the Prevention of Stroke: Clinical Practice Recommendations from the Lipid Association of India. Current Vascular Pharmacology, 2022, 20, 134-155.

LIVER FUNCTION PROFILE, SERUM

BILIRUBIN, TOTAL	1.00	Upto 1.2	mg/dL
METHOD : SPECTROPHOTOMETRY, COLORIMETRIC -DIAZO METHOD			
BILIRUBIN, DIRECT	0.60 High	< or = 0.3	mg/dL
METHOD : SPECTROPHOTOMETRY, JENDRASSIK & GROFF - DIAZOTIZATION			
BILIRUBIN, INDIRECT	0.4	0.0 - 0.9	mg/dL
METHOD : CALCULATED PARAMETER			
TOTAL PROTEIN	7.0	6.0 - 8.0	g/dL
METHOD : SPECTROPHOTOMETRY, COLORIMETRIC -BIURET, REAGENT BLANK, SERUM BLANK			
ALBUMIN	4.4	3.97 - 4.94	g/dL
METHOD : SPECTROPHOTOMETRY, BROMOCRESOL GREEN(BCG) - DYE BINDING			
GLOBULIN	2.6	2.0 - 3.5	g/dL
METHOD : CALCULATED PARAMETER			
ALBUMIN/GLOBULIN RATIO	1.7	1.0 - 2.1	RATIO
METHOD : CALCULATED PARAMETER			
ASPARTATE AMINOTRANSFERASE(AST/SGOT)	25	Upto 40	U/L
METHOD : SPECTROPHOTOMETRY, WITHOUT PYRIDOXAL PHOSPHATE ACTIVATION(P5P) - IFCC			
ALANINE AMINOTRANSFERASE (ALT/SGPT)	35	Upto 41	U/L
METHOD : SPECTROPHOTOMETRY, WITHOUT PYRIDOXAL PHOSPHATE ACTIVATION(P5P) - IFCC			
ALKALINE PHOSPHATASE	85	40 - 129	U/L
METHOD : SPECTROPHOTOMETRY, PNPP, AMP BUFFER - IFCC			
GAMMA GLUTAMYL TRANSFERASE (GGT)	33	< 60	U/L

Dr. Apeksha Sharma
D.P.B.,DNB (PATH)
(Reg.no.MMC2008/06/2561)
Consultant Pathologist

Dr. Deepak Sanghavi, M.D(Path)
(Reg.no.MMC2004/03/1530)
Chief of Mumbai Reference Lab.



View Details



View Report

PERFORMED AT :

Agilus Diagnostics Ltd
Prime Square Building, Plot No 1, Gaiwadi Industrial Estate, S.V. Road, Goregaon (W)
Mumbai, 400062
Maharashtra, India
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956





PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON W

ACCESSION NO : **0002XD020787**
 PATIENT ID : AMITM1910872
 CLIENT PATIENT ID:
 ABHA NO :

AGE/SEX : 37 Years Male
 DRAWN : 13/04/2024 08:48:36
 RECEIVED : 13/04/2024 08:50:04
 REPORTED : 17/04/2024 17:27:55

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

METHOD : SPECTROPHOTOMETRY, ENZYMATIC COLORIMETRIC - G-GLUTAMYL-CARBOXY-NITROANILIDE - IFCC
LACTATE DEHYDROGENASE 228 < 232 U/L
 METHOD : SPECTROPHOTOMETRY, LACTATE TO PYRUVATE - UV-IFCC

BLOOD UREA NITROGEN (BUN), SERUM

BLOOD UREA NITROGEN 10 6 - 20 mg/dL
 METHOD : SPECTROPHOTOMETRY, UREASE -COLORIMETRIC

CREATININE, SERUM

CREATININE 0.79 Low 0.90 - 1.30 mg/dL
 METHOD : SPECTROPHOTOMETRY, JAFFE'S ALKALINE PICRATE KINETIC - RATE BLANKED - IFCC-IDMS STANDARDIZED

BUN/CREAT RATIO

BUN/CREAT RATIO 12.66 8 - 15
 METHOD : CALCULATED PARAMETER

URIC ACID, SERUM


URIC ACID 6.7 3.4 - 7.0 mg/dL
 METHOD : SPECTROPHOTOMETRY, ENZYMATIC COLORIMETRIC- URICASE


TOTAL PROTEIN, SERUM

TOTAL PROTEIN 7.0 6.0 - 8.0 g/dL
 METHOD : SPECTROPHOTOMETRY, COLORIMETRIC -BIURET, REAGENT BLANK, SERUM BLANK

ALBUMIN, SERUM

ALBUMIN 4.4 3.97 - 4.94 g/dL
 METHOD : SPECTROPHOTOMETRY, BROMOCRESOL GREEN(BCG) - DYE BINDING


Dr. Apeksha Sharma
D.P.B.,DNB (PATH)
(Reg.no.MMC2008/06/2561)
Consultant Pathologist


Dr. Deepak Sanghavi,M.D(Path)
(Reg.no.MMC2004/03/1530)
Chief of Mumbai Reference Lab.



View Details



View Report

PERFORMED AT :
Agilus Diagnostics Ltd
 Prime Square Building,Plot No 1,Gaiwadi Industrial Estate,S.V. Road,Goregaon (W)
 Mumbai, 400062
 Maharashtra, India
 Tel : 9111591115, Fax :
 CIN - U74899PB1995PLC045956





PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON W

ACCESSION NO : **0002XD020787**
 PATIENT ID : AMITM1910872
 CLIENT PATIENT ID :
 ABHA NO :

AGE/SEX : 37 Years Male
 DRAWN : 13/04/2024 08:48:36
 RECEIVED : 13/04/2024 08:50:04
 REPORTED : 17/04/2024 17:27:55

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

GLOBULIN

GLOBULIN	2.6	2.0 - 3.5	g/dL
METHOD : CALCULATED PARAMETER			

ELECTROLYTES (NA/K/CL), SERUM

SODIUM, SERUM	138	136 - 145	mmol/L
METHOD : ISE INDIRECT			
POTASSIUM, SERUM	4.90	3.5 - 5.1	mmol/L
METHOD : ISE INDIRECT			
CHLORIDE, SERUM	101	98 - 106	mmol/L
METHOD : ISE INDIRECT			

Interpretation(s)

Sodium	Potassium	Chloride
Decreased in: CCF, cirrhosis, vomiting, diarrhea, excessive sweating, salt-losing nephropathy, adrenal insufficiency, nephrotic syndrome, water intoxication, SIADH. Drugs: thiazides, diuretics, ACE inhibitors, chlorpropamide, carbamazepine, anti depressants (SSRI), antipsychotics.	Decreased in: Low potassium intake, prolonged vomiting or diarrhea, RTA types I and II, hyperaldosteronism, Cushing's syndrome, osmotic diuresis (e.g., hyperglycemia), alkalosis, familial periodic paralysis, trauma (transient). Drugs: Adrenergic agents, diuretics.	Decreased in: Vomiting, diarrhea, renal failure combined with salt deprivation, over-treatment with diuretics, chronic respiratory acidosis, diabetic ketoacidosis, excessive sweating, SIADH, salt-losing nephropathy, porphyria, expansion of extracellular fluid volume, adrenal insufficiency, hyperaldosteronism, metabolic alkalosis. Drugs: chronic laxative, corticosteroids, diuretics.
Increased in: Dehydration (excessive sweating, severe vomiting or diarrhea), diabetes mellitus, diabetes insipidus, hyperaldosteronism, inadequate water intake. Drugs: steroids, licorice, oral contraceptives.	Increased in: Massive hemolysis, severe tissue damage, rhabdomyolysis, acidosis, dehydration, renal failure, Addison's disease, RTA type IV, hyperkalemic familial periodic paralysis. Drugs: potassium salts, potassium-sparing diuretics, NSAIDs, beta-blockers, ACE inhibitors, high-dose trimethoprim-sulfamethoxazole.	Increased in: Renal failure, nephrotic syndrome, RTA, dehydration, overtreatment with saline, hyperparathyroidism, diabetes insipidus, metabolic acidosis from diarrhea (Loss of HCO3-), respiratory alkalosis, hyperadrenocorticism. Drugs: acetazolamide, androgens, hydrochlorothiazide, salicylates.

Dr. Apeksha Sharma
D.P.B., DNB (PATH)
(Reg.no.MMC2008/06/2561)
Consultant Pathologist

Dr. Deepak Sanghavi, M.D(Path)
(Reg.no.MMC2004/03/1530)
Chief of Mumbai Reference Lab.



View Details



View Report

PERFORMED AT :

Agilus Diagnostics Ltd
 Prime Square Building, Plot No 1, Gaiwadi Industrial Estate, S.V. Road, Goregaon (W)
 Mumbai, 400062
 Maharashtra, India
 Tel : 9111591115, Fax :
 CIN - U74899PB1995PLC045956





PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON W

ACCESSION NO : **0002XD020787**

PATIENT ID : AMITM1910872

CLIENT PATIENT ID:

ABHA NO :

AGE/SEX : 37 Years Male

DRAWN : 13/04/2024 08:48:36

RECEIVED : 13/04/2024 08:50:04

REPORTED : 17/04/2024 17:27:55

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

Interferences: Severe lipemia or hyperproteinemi, if sodium analysis involves a dilution step can cause spurious results. The serum sodium falls about 1.6 mEq/L for each 100 mg/dL increase in blood glucose.

Interferences: Hemolysis of sample, delayed separation of serum, prolonged fist clenching during blood drawing, and prolonged tourniquet placement. Very high WBC/PLT counts may cause spurious. Plasma potassium levels are normal.

Interferences: Test is helpful in assessing normal and increased anion gap metabolic acidosis and in distinguishing hypercalcemia due to hyperparathyroidism (high serum chloride) from that due to malignancy (Normal serum chloride)

Interpretation(s)

GLUCOSE FASTING, FLUORIDE PLASMA-TEST DESCRIPTION

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and so that no glucose is excreted in the urine.

Increased in: Diabetes mellitus, Cushing's syndrome (10 – 15%), chronic pancreatitis (30%). Drugs: corticosteroids, phenytoin, estrogen, thiazides.

Decreased in : Pancreatic islet cell disease with increased insulin, insulinoma, adrenocortical insufficiency, hypopituitarism, diffuse liver disease, malignancy (adrenocortical, stomach, fibrosarcoma), infant of a diabetic mother, enzyme deficiency diseases (e.g. galactosemia), Drugs- insulin, ethanol, propranolol; sulfonylureas, tolbutamide, and other oral hypoglycemic agents.

NOTE: While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values), there is wide fluctuation within individuals. Thus, glycosylated hemoglobin (HbA1c) levels are favored to monitor glycemic control.

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glycosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.

GLUCOSE, POST-PRANDIAL, PLASMA- High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glycosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc. Additional test HbA1c LIVER FUNCTION PROFILE, SERUM-

Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. **Elevated levels** results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors & Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health. AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis.

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Pagets disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilsons disease.

GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc.

Total Protein also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease. Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

Albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc

BLOOD UREA NITROGEN (BUN), SERUM- Causes of Increased levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)

Causes of decreased level include Liver disease, SIADH.

CREATININE, SERUM- Higher than normal level may be due to:

• Blockage in the urinary tract, Kidney problems, such as kidney damage or failure, infection, or reduced blood flow, Loss of body fluid (dehydration), Muscle problems, such as breakdown of muscle fibers, Problems during pregnancy, such as seizures (eclampsia), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to: • Myasthenia Gravis, Muscuophy

URIC ACID, SERUM- Causes of Increased levels: -Dietary (High Protein Intake, Prolonged Fasting, Rapid weight loss), Gout, Lesch nyhan syndrome, Type 2 DM, Metabolic syndrome **Causes of decreased levels:** -Low Zinc intake, OCP, Multiple Sclerosis

TOTAL PROTEIN, SERUM- is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin.

Dr. Apeksha Sharma
D.P.B., DNB (PATH)
(Reg.no.MMC2008/06/2561)
Consultant Pathologist

Dr. Deepak Sanghavi, M.D (Path)
(Reg.no.MMC2004/03/1530)
Chief of Mumbai Reference Lab.



View Details



View Report

PERFORMED AT :

Agilus Diagnostics Ltd
Prime Square Building, Plot No 1, Gaiwadi Industrial Estate, S.V. Road, Goregaon (W)
Mumbai, 400062
Maharashtra, India
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956





PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON W

ACCESSION NO : **0002XD020787**
 PATIENT ID : AMITM1910872
 CLIENT PATIENT ID:
 ABHA NO :

AGE/SEX : 37 Years Male
 DRAWN : 13/04/2024 08:48:36
 RECEIVED : 13/04/2024 08:50:04
 REPORTED : 17/04/2024 17:27:55

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease.
Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.
ALBUMIN, SERUM-Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. **Low blood albumin levels (hypoalbuminemia) can be caused by:** Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.

Dr. Apeksha Sharma
D.P.B., DNB (PATH)
(Reg.no.MMC2008/06/2561)
Consultant Pathologist

Dr. Deepak Sanghavi, M.D(Path)
(Reg.no.MMC2004/03/1530)
Chief of Mumbai Reference Lab.



View Details



View Report

PERFORMED AT :
Agilus Diagnostics Ltd
 Prime Square Building, Plot No 1, Gaiwadi Industrial Estate, S.V. Road, Goregaon (W)
 Mumbai, 400062
 Maharashtra, India
 Tel : 9111591115, Fax :
 CIN - U74899PB1995PLC045956



ULR No.2000012351057-0002



PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON W

ACCESSION NO : **0002XD020787**
 PATIENT ID : AMITM1910872
 CLIENT PATIENT ID:
 ABHA NO :

AGE/SEX : 37 Years Male
 DRAWN : 13/04/2024 08:48:36
 RECEIVED : 13/04/2024 08:50:04
 REPORTED : 17/04/2024 17:27:55

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

CLINICAL PATH - URINALYSIS

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

PHYSICAL EXAMINATION, URINE

COLOR PALE YELLOW
 APPEARANCE CLEAR

CHEMICAL EXAMINATION, URINE

PH	6.5	4.6 - 8.0
SPECIFIC GRAVITY	1.000 Low	1.003 - 1.035
PROTEIN	NOT DETECTED	NOT DETECTED
GLUCOSE	NOT DETECTED	NOT DETECTED
KETONES	NOT DETECTED	NOT DETECTED
BLOOD	NOT DETECTED	NOT DETECTED
BILIRUBIN	NOT DETECTED	NOT DETECTED
UROBILINOGEN	NOT DETECTED	NOT DETECTED
NITRITE	NOT DETECTED	NOT DETECTED
LEUKOCYTE ESTERASE	NOT DETECTED	NOT DETECTED

MICROSCOPIC EXAMINATION, URINE

RED BLOOD CELLS	NOT DETECTED	NOT DETECTED	/HPF
PUS CELL (WBC'S)	0-1	0-5	/HPF
EPITHELIAL CELLS	0-1	0-5	/HPF
CASTS	NOT DETECTED		
CRYSTALS	NOT DETECTED		
BACTERIA	NOT DETECTED	NOT DETECTED	
YEAST	NOT DETECTED	NOT DETECTED	

METHOD : URINE ROUTINE & MICROSCOPY EXAMINATION BY INTEGRATED AUTOMATED SYSTEM. (PH-DOUBLE INDICATOR,SP. GRAVITY-IONIC CONCEN, GLUCOSE-GOD/POD,PROTEIN- ERROR OF INDICATORS,KETONE-LEGAL'S,BLOOD- PEROXIDASE ACTIVITY-HB,BILIRUBIN-DIAZOTIZATION,UROBILINOGEN-DIAZOTIZATION,NITRITE-GRIESS,LEUKOCYTES- ESTERASES ACTIVITY)

Dr. Deepak Sanghavi, M.D(Path)
 (Reg.no.MMC2004/03/1530)
 Chief of Mumbai Reference Lab.

Dr. Apeksha Sharma
 D.P.B., DNB (PATH)
 (Reg.no.MMC2008/06/2561)
 Consultant Pathologist



View Details



View Report

PERFORMED AT :

Agilus Diagnostics Ltd
 Prime Square Building, Plot No 1, Gaiwadi Industrial Estate, S.V. Road, Goregaon (W)
 Mumbai, 400062
 Maharashtra, India
 Tel : 9111591115, Fax :
 CIN - U74899PB1995PLC045956



ULR No.2000012351057-0002



PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON W

ACCESSION NO : **0002XD020787**
 PATIENT ID : AMITM1910872
 CLIENT PATIENT ID:
 ABHA NO :

AGE/SEX : 37 Years Male
 DRAWN : 13/04/2024 08:48:36
 RECEIVED : 13/04/2024 08:50:04
 REPORTED : 17/04/2024 17:27:55

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

Interpretation(s)

The following table describes the probable conditions, in which the analytes are present in urine

Presence of	Conditions
Proteins	Inflammation or immune illnesses
Pus (White Blood Cells)	Urinary tract infection, urinary tract or kidney stone, tumors or any kind of kidney impairment
Glucose	Diabetes or kidney disease
Ketones	Diabetic ketoacidosis (DKA), starvation or thirst
Urobilinogen	Liver disease such as hepatitis or cirrhosis
Blood	Renal or genital disorders/trauma
Bilirubin	Liver disease
Erythrocytes	Urological diseases (e.g. kidney and bladder cancer, urolithiasis), urinary tract infection and glomerular diseases
Leukocytes	Urinary tract infection, glomerulonephritis, interstitial nephritis either acute or chronic, polycystic kidney disease, urolithiasis, contamination by genital secretions
Epithelial cells	Urolithiasis, bladder carcinoma or hydronephrosis, ureteric stents or bladder catheters for prolonged periods of time
Granular Casts	Low intratubular pH, high urine osmolality and sodium concentration, interaction with Bence-Jones protein
Hyaline casts	Physical stress, fever, dehydration, acute congestive heart failure, renal diseases
Calcium oxalate	Metabolic stone disease, primary or secondary hyperoxaluria, intravenous infusion of large doses of vitamin C, the use of vasodilator naftidrofuryl oxalate or the gastrointestinal lipase inhibitor orlistat, ingestion of ethylene glycol or of star fruit (Averrhoa carambola) or its juice
Uric acid	arthritis
Bacteria	Urinary infection when present in significant numbers & with pus cells.
Trichomonas vaginalis	Vaginitis, cervicitis or salpingitis

Dr. Deepak Sanghavi, M.D(Path)
 (Reg.no.MMC2004/03/1530)
 Chief of Mumbai Reference Lab.

Dr. Apeksha Sharma
 D.P.B., DNB (PATH)
 (Reg.no.MMC2008/06/2561)
 Consultant Pathologist



View Details



View Report

PERFORMED AT :

Agilus Diagnostics Ltd
 Prime Square Building, Plot No 1, Gaiwadi Industrial Estate, S.V. Road, Goregaon (W)
 Mumbai, 400062
 Maharashtra, India
 Tel : 9111591115, Fax :
 CIN - U74899PB1995PLC045956



PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON WACCESSION NO : **0002XD020787**

PATIENT ID : AMITM1910872

CLIENT PATIENT ID:

ABHA NO :

AGE/SEX : 37 Years Male

DRAWN : 13/04/2024 08:48:36

RECEIVED : 13/04/2024 08:50:04

REPORTED : 17/04/2024 17:27:55

Test Report Status **Final**

Results

Biological Reference Interval Units

CLINICAL PATH - STOOL ANALYSIS

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE**MICROSCOPIC EXAMINATION,STOOL**

REMARK

TEST CANCELLED AS SPECIMEN NOT RECEIVED

Interpretation(s)

Stool routine analysis is only a screening test for disorders of gastrointestinal tract like infection, malabsorption, etc. The following table describes the probable conditions, in which the analytes are present in stool.

PRESENCE OF	CONDITION
Pus cells	Pus in the stool is an indication of infection
Red Blood cells	Parasitic or bacterial infection or an inflammatory bowel condition such as ulcerative colitis
Parasites	Infection of the digestive system. Stool examination for ova and parasite detects presence of parasitic infestation of gastrointestinal tract. Various forms of parasite that can be detected include cyst, trophozoite and larvae. One negative result does not rule out the possibility of parasitic infestation. Intermittent shedding of parasites warrants examinations of multiple specimens tested on consecutive days. Stool specimens for parasitic examination should be collected before initiation of antidiarrheal therapy or antiparasitic therapy. This test does not detect presence of opportunistic parasites like Cyclospora, Cryptosporidia and Isospora species. Examination of Ova and Parasite has been carried out by direct and concentration techniques.
Mucus	Mucus is a protective layer that lubricates, protects & reduces damage due to bacteria or viruses.
Charcot-Leyden crystal	Parasitic diseases.
Ova & cyst	Ova & cyst indicate parasitic infestation of intestine.
Frank blood	Bleeding in the rectum or colon.
Occult blood	Occult blood indicates upper GI bleeding.
Macrophages	Macrophages in stool are an indication of infection as they are protective cells.
Epithelial cells	Epithelial cells that normally line the body surface and internal organs show up in stool when there is inflammation or infection.
Fat	Increased fat in stool maybe seen in conditions like diarrhoea or malabsorption.
pH	Normal stool pH is slightly acidic to neutral. Breast-fed babies generally have an acidic stool.

ADDITIONAL STOOL TESTS :


Dr. Ekta Patil, MD
(Reg.No. MMC2008/04/1142)
Senior Microbiologist

Page 18 Of 21



View Details



View Report

PERFORMED AT :

Agilus Diagnostics Ltd
Prime Square Building, Plot No 1, Gaiwadi Industrial Estate, S.V. Road, Goregaon (W)
Mumbai, 400062
Maharashtra, India
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956



ULR No. 2000012351057-0002



PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON W

ACCESSION NO : **0002XD020787**
 PATIENT ID : AMITM1910872
 CLIENT PATIENT ID:
 ABHA NO :

AGE/SEX : 37 Years Male
 DRAWN : 13/04/2024 08:48:36
 RECEIVED : 13/04/2024 08:50:04
 REPORTED : 17/04/2024 17:27:55

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

- Stool Culture**:- This test is done to find cause of GI infection, make decision about best treatment for GI infection & to find out if treatment for GI infection worked.
- Fecal Calprotectin**: It is a marker of intestinal inflammation. This test is done to differentiate Inflammatory Bowel Disease (IBD) from Irritable Bowel Syndrome (IBS).
- Fecal Occult Blood Test(FOBT)**: This test is done to screen for colon cancer & to evaluate possible cause of unexplained anaemia.
- Clostridium Difficile Toxin Assay**: This test is strongly recommended in healthcare associated bloody or waterydiarrhoea, due to overuse of broad spectrum antibiotics which alter the normal GI flora.
- Biofire (Film Array) GI PANEL**: In patients of Diarrhoea, Dysentery, Rice watery Stool, FDA approved, Biofire Film Array Test,(Real Time Multiplex PCR) is strongly recommended as it identifies organisms, bacteria,fungi,virus ,parasite and other opportunistic pathogens, Vibrio cholera infections only in 3 hours. Sensitivity 96% & Specificity 99%.
- Rota Virus Immunoassay**: This test is recommended in severe gastroenteritis in infants & children associated with watery diarrhoea, vomiting& abdominal cramps. Adults are also affected. It is highly contagious in nature.

Dr. Ekta Patil,MD
 (Reg.No. MMC2008/04/1142)
 Senior Microbiologist



View Details



View Report

PERFORMED AT :

Agilus Diagnostics Ltd
 Prime Square Building,Plot No 1,Gaiwadi Industrial Estate,S.V. Road,Goregaon (W)
 Mumbai, 400062
 Maharashtra, India
 Tel : 9111591115, Fax :
 CIN - U74899PB1995PLC045956



ULR No.2000012351057-0002



PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON W

ACCESSION NO : **0002XD020787**
PATIENT ID : AMITM1910872
CLIENT PATIENT ID:
ABHA NO :

AGE/SEX : 37 Years Male
DRAWN : 13/04/2024 08:48:36
RECEIVED : 13/04/2024 08:50:04
REPORTED : 17/04/2024 17:27:55

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

SPECIALISED CHEMISTRY - HORMONE

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

THYROID PANEL, SERUM

T3	136.0	80.0 - 200.0	ng/dL
METHOD : COMPETITIVE ELECTROCHEMILUMINESCENCE IMMUNOASSAY			
T4	8.47	5.10 - 14.10	µg/dL
METHOD : COMPETITIVE ELECTROCHEMILUMINESCENCE IMMUNOASSAY			
TSH (ULTRASENSITIVE)	1.620	0.270 - 4.200	µIU/mL
METHOD : SANDWICH ELECTROCHEMILUMINESCENCE IMMUNOASSAY			

Interpretation(s)

Triiodothyronine T3, **Thyroxine T4**, and **Thyroid Stimulating Hormone TSH** are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate. Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH. Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hyperthyroidism, TSH levels are low. Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3. Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Sr. No.	TSH	Total T4	FT4	Total T3	Possible Conditions
1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3) Post Thyroidectomy (4) Post Radio-Iodine treatment
2	High	Normal	Normal	Normal	(1) Subclinical Hypothyroidism (2) Patient with insufficient thyroid hormone replacement therapy (3) In cases of Autoimmune/Hashimoto thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical inflammation, drugs like amphetamines, Iodine containing drug and dopamine antagonist e.g. domperidone and other physiological reasons.
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre (3) Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4 replacement therapy (7) First trimester of Pregnancy
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism

Dr. Deepak Sanghavi, M.D(Path)
(Reg.no.MMC2004/03/1530)
Chief of Mumbai Reference Lab.

Dr. Apeksha Sharma
D.P.B., DNB (PATH)
(Reg.no.MMC2008/06/2561)
Consultant Pathologist



View Details



View Report

PERFORMED AT :

Agilus Diagnostics Ltd
Prime Square Building, Plot No 1, Gaiwadi Industrial Estate, S.V. Road, Goregaon (W)
Mumbai, 400062
Maharashtra, India
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956



PATIENT NAME : AMIT JAMBHULKAR

REF. DOCTOR : SELF

NEW SURISE TOWER CHSL UNNAT NAGAR
GOREGAON W

ACCESSION NO : 0002XD020787

PATIENT ID : AMITM1910872

CLIENT PATIENT ID:

ABHA NO :

AGE/SEX : 37 Years Male

DRAWN : 13/04/2024 08:48:36

RECEIVED : 13/04/2024 08:50:04

REPORTED : 17/04/2024 17:27:55

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent treatment for Hyperthyroidism
8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies

REF: 1. TIETZ Fundamentals of Clinical chemistry 2.Guidlines of the American Thyroid association during pregnancy and Postpartum, 2011.

TSH in pregnancy

There's reduction in both the lower and the upper limit of maternal TSH relative to the non-pregnant TSH reference range. This is because of elevated levels of serum hCG that directly stimulates the TSH receptor, thereby increasing thyroid hormone production. The largest decrease in serum TSH is observed during the first trimester. Thereafter, serum TSH and its reference range gradually increases in the second and third trimesters, but nonetheless remains lower than in non-pregnant women.

NOTE: It is advisable to detect Free T3,FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

****End Of Report****Please visit www.agilusdiagnostics.com for related Test Information for this accession**CONDITIONS OF LABORATORY TESTING & REPORTING**

1. It is presumed that the test sample belongs to the patient named or identified in the test requisition form.
2. All tests are performed and reported as per the turnaround time stated in the AGILUS Directory of Services.
3. Result delays could occur due to unforeseen circumstances such as non-availability of kits / equipment breakdown / natural calamities / technical downtime or any other unforeseen event.
4. A requested test might not be performed if:
 - i. Specimen received is insufficient or inappropriate
 - ii. Specimen quality is unsatisfactory
 - iii. Incorrect specimen type
 - iv. Discrepancy between identification on specimen container label and test requisition form
5. AGILUS Diagnostics confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity.
6. Laboratory results should not be interpreted in isolation; it must be correlated with clinical information and be interpreted by registered medical practitioners only to determine final diagnosis.
7. Test results may vary based on time of collection, physiological condition of the patient, current medication or nutritional and dietary changes. Please consult your doctor or call us for any clarification.
8. Test results cannot be used for Medico legal purposes.
9. In case of queries please call customer care (91115 91115) within 48 hours of the report.

Agilus Diagnostics LimitedFortis Hospital, Sector 62, Phase VIII,
Mohali 160062


Dr. Deepak Sanghavi, M.D(Path)
(Reg.no.MMC2004/03/1530)
Chief of Mumbai Reference Lab.



Dr. Apeksha Sharma
D.P.B., DNB (PATH)
(Reg.no.MMC2008/06/2561)
Consultant Pathologist

Page 21 Of 21



View Details



View Report

PERFORMED AT :

Agilus Diagnostics Ltd
Prime Square Building, Plot No 1, Gaiwadi Industrial Estate, S.V. Road, Goregaon (W)
Mumbai, 400062
Maharashtra, India
Tel : 9111591115, Fax :
CIN - U74899PB1995PLC045956



ULR No.2000012351057-0002