





भारतीय विशिष्ट पहचान प्राधिकरण UNIQUE IDENTIFICATION AUTHORITY OF INDIA

पताः

W/O दिनेश कुमार शर्मा, बार्ड न १२ कसेरा भवन माहल्ला लोकनाथ का, श्री माधोपुर, सीकर, राजस्थान - 332715

Address:

W/O Dinesh Kumar Sharma, ward n 12 kasera bhawan mohalla loknath ka, Sri Madhopur, Srimadhopur, Sikar, Rajasthan - 332715

8807




help@uldal.gov.in www.uldal.gov.in P.O. Box No. 1947, Bengaluru-560 001



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

 B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023





General Physical Examination

Age: 42 DOB: 01/01/1922 Sex:fample
* v
77
Wt: <u>80</u> (Kg)
Abdomen Circumference: 17 % (cm)
min RR: 18 / min Temp: Afallace
A
. 416 ACB
ntally fit: Yes / No
- Name of Examinee: SPROS.DEV)
Name Medical Examiner - PINUSH COOMS Ologist)

P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axix Bank
 Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⑥ +91 141 4824885 ⑤ maxcarediagnostics1@gmail.com





NAME :- Mrs. SAROJ DEVI

Age:- 47 Yrs 2 Mon 25 Days

Sex :- Female

Patient ID :-12234987

Date :- 26/03/2024

10:30:52

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :-

Mr.MEDIWHEEL

Final Authentication: 26/03/2024 17:37:27

HAEMOGARAM

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval	=
FULL BODY HEALTH CHECKUP ABOVE 401	FEMALE			_
HAEMOGLOBIN (Hb)	11.8 └	g/dL	12.0 - 15.0	
TOTAL LEUCOCYTE COUNT	6.30	/cumm	4.00 - 10.00	
DIFFERENTIAL LEUCOCYTE COUNT				
NEUTROPHIL	51.0	%	40.0 - 80.0	
LYMPHOCYTE	43.0 H	%	20.0 - 40.0	
EOSINOPHIL	2.0	%	1.0 - 6.0	
MONOCYTE	4.0	%	2.0 - 10.0	
BASOPHIL	0.0	%	0.0 - 2.0	
TOTAL RED BLOOD CELL COUNT (RBC)	4.08	x10^6/uL	3.80 - 4.80	
HEMATOCRIT (HCT)	37.50	%	36.00 - 46.00	
MEAN CORP VOLUME (MCV)	92.0	T.	83.0 - 101.0	
MEAN CORP HB (MCH)	28.9	pg	27.0 - 32.0	
MEAN CORP HB CONC (MCHC)	31.5	g/dL	31.5 - 34.5	
PLATELET COUNT	271	x10^3/uL	150 - 410	
RDW-CV	15.0 H	%	11.6 - 14.0	

Technologist MGR Page No: 1 of 15



O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

③ +91 141 4824885 ☑ maxcarediagnostics1@gmail.com





NAME :- Mrs. SAROJ DEVI

Age :-47 Yrs 2 Mon 25 Days

Female Sex :-

Patient ID: -12234987

Date :- 26/03/2024

10:30:52

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :-

Mr.MEDIWHEEL

Final Authentication: 26/03/2024 17:37:27

HAEMATOLOGY

Erythrocyte Sedimentation Rate (ESR)

mm in 1st hr

00 - 20

The erythrocyte sedimentation rate (ESR or sed rate) is a relatively simple, inexpensive, non-specific test that has been used for many years to help detect inflammation associated with conditions such as infections, cancers, and autoimmune diseases.ESR is said to be a non-specific test because an elevated result often indicates the presence of inflammation but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other tests, such as C-reactive protein. ESR is used to help diagnose certain specific inflammatory diseases, including temporal arteritis, systemic vasculitis and polymyalgia rheumatica. (For more on these, read the article on Vasculitis.) A significantly elevated ESR is one of the main test results used to support the diagnosis. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as



Technologist Page No: 2 of 15



O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023





NAME :- Mrs. SAROJ DEVI

Age :-47 Yrs 2 Mon 25 Days

Sex :-Female Patient ID :-12234987

Date :- 26/03/2024

10:30:52

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company:-

Mr.MEDIWHEEL

(CBC): Methodology: TLC,DLC Fluorescent Flow cytometry, HB SLS method,TRBC,PCV,PLT Hydrodynamically focused Impedance. and MCH,MCV,MCHC,MENTZER INDEX are calculated. InstrumentName: Sysmex 6 part fully automatic analyzer XN-L,Japan



MGR Page No: 3 of 15



B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

(8) +91 141 4824885 (2) maxcarediagnostics 1@gmail.com





NAME :- Mrs. SAROJ DEVI

47 Yrs 2 Mon 25 Days

Female

Age :-

Sex :-

Patient ID :-12234987

Date :- 26/03/2024

10:30:52

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :-

Mr.MEDIWHEEL

Final Authentication: 26/03/2024 17:37:27

BIOCHEMISTRY

DIOCHEMISTRI				
Test Name	Value	Unit	Biological Ref Interval	
FASTING BLOOD SUGAR (Plasma) Methord:- GOD POD	77.1	mg/dl	70.0 - 115.0	
Impaired glucose tolerance (IGT)	111 - 125 mg/dL			
Diabetes Mellitus (DM)	> 126 mg/dL			

Instrument Name: HORIBA CA60 Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm,

hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin

therapy or various liver diseases.

BLOOD SUGAR PP (Plasma)

Methord:- GOD PAP

111.0

mg/dl

70.0 - 140.0

Instrument Name: HORIBA Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels(hypoglycemia) may result from excessive insulin therapy or various liver diseases.

Technologist MGR Page No: 4 of 15 DR.TANU RUNGTA MD (Pathology) RMC No. 17226

This Report Is Not Valid For Medico Legal Purpose



O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

③ +91 141 4824885 ۞ maxcarediagnostics1@gmail.com





NAME: - Mrs. SAROJ DEVI

Age :-

47 Yrs 2 Mon 25 Days

Sex :-Female Patient ID: -12234987

Date :- 26/03/2024

10:30:52

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :-Mr.MEDIWHEEL

Final Authentication: 26/03/2024 17:37:27

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
GLYCOSYLATED HEMOGLOBIN (Hb.A. Methord:- CAPILLARY with EDTA	MIC) 5.4	%	Non-diabetic: < 5.7 Pre-diabetics: 5.7-6.4 Diabetics: = 6.5 or higher ADA Target: 7.0 Action suggested: > 6.5
MEAN PLASMA GLUCOSE Methord:- Calculated Parameter	106	mg/dL	68 - 125

INTERPRETATION

AS PER AMERICAN DIABETES ASSOCIATION (ADA) Reference Group HbA1c in % Non diabetic adults >=18 years < 5.7 At risk (Prediabetes) 5.7 - 6.4 Diagnosing Diabetes >= 6.5

CLINICAL NOTES

In vitro quantitative determination of HbA1c in whole blood is utilized in long term monitoring of glycemia. The HbA1c level correlates with the mean glucose concentration prevailing in the course of the patient's recent history (approx - 6-8 weeks) and therefore provides much more reliable information for glycemia monitoring than do determinations of blood glucose or urinary glucose. It is recommended that the determination of HbA1c be performed at intervals of 4-6 weeks during Diabetes Mellitus therapy. Results of HbA1c should be assessed in conjunction with the patient's medical history, clinical examinations and other findings. Some of the factors that influence HbA1c and its measurement [Adapted from Gallagher et al]


- 1. Erythropoiesis
- Increased HbA1c: iron, vitamin B12 deficiency, decreased erythropolesis
- Decreased HbA1c: administration of erythropoletin, iron, vitamin B12, reticulocytosis, chronic liver disease,
 Altered Haemoglobin-Genetic or chemical alterations in hemoglobin: hemoglobin-behaptines, HbF, methemoglobin, may increase or decrease HbA1c.
- 3. Glycation
- Increased HbA1c: alcoholism, chronic renal failure, decreased intraerythrocytic pH.
- Decreased HbA1c: certain hemoglobinopathies, increased intra-erythrocyte pH
- 4. Erythrocyte destruction
- Increased HbA1c: increased erythrocyte life span: Splenectomy
- Decreased A1c: decreased RBC life span: hemoglobinopathies, splencmegaly, rheumatoid arthritis or drugs such as antiretrovirals, ribavirin & dapsone
- 5. Others
- Increased HbA1c: hyperbilirubinemia, carbamylated hemoglobin, alcoholism, large doses of aspirin, chronic opiate use,chronic renal failure

- Decreased HbA1c: hypertriglyceridemia, reticulocytosis, chronic liver disease, aspirin, vitamin C and E, splenomegaly, rheumatoid arthritis or drugs

Technologist MGR Page No: 5 of 15



O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023





NAME :- Mrs. SAROJ DEVI

Age :-

47 Yrs 2 Mon 25 Days

Sex :-Female Patient ID :-12234987

Date :- 26/03/2024

10:30:52

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :-

Mr.MEDIWHEEL

Final Authentication: 26/03/2024 17:37:27

HAEMATOLOGY

BLOOD GROUP ABO Methord:- Haemagglutination reaction "O" POSITIVE



Technologist MGR Page No: 6 of 15



B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023





NAME :- Mrs. SAROJ DEVI

Patient ID: -12234987

Date :- 26/03/2024

10:30:52

Age :-

47 Yrs 2 Mon 25 Days

Ref. By Doctor:-BANK OF BARODA

Sex :-Female Lab/Hosp :-Company:-

Mr.MEDIWHEEL

Final Authentication: 26/03/2024 17:37:27

BIOCHEMISTRY			
Test Name	Value	Unit	Biological Ref Interval
LIPID PROFILE			8
TOTAL CHOLESTEROL Methord:- CHOD-PAP methodology	171.00	mg/dl	Desirable <200 Borderline 200-239 High> 240
InstrumentName:MISPA PLUS Interpreta disorders.	tion: Cholesterol measurements	are used in the diagnosis	and treatments of lipid lipoprotein metabolism
TRIGLYCERIDES Methord:- GPO-PAP	105.60	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500
InstrumentNama: Randov Ry Imola Intern	retation : Triplyceride measure	ments are used in the diag	mosis and treatment of diseases involving linid

InstrumentName: Randox Rx Imola Interpretation: Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction.

DIRECT HDL CHOLESTEROL Methord:- Direct clearance Method

43.20

mg/dl

MALE- 30-70 **FEMALE - 30-85**

Instrument Name: Rx Daytona plus Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.

LDL CHOLESTEROL Methord:- Calculated Method	110.20	mg/dl	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190
VLDL CHOLESTEROL Methord: - Calculated	21.12	mg/dl	0.00 - 80.00
T.CHOLESTEROL/HDL CHOLESTEROL RATIO Methord:- Calculated	3.96		0.00 - 4.90
LDL / HDL CHOLESTEROL RATIO Methord:- Calculated	2.55		0.00 - 3.50
TOTAL LIPID Methord: CALCULATED	511.21	mg/dl	400.00 - 1000.00

¹ Measurements in the same patient can show physiological& analytical variations. Three serialsamples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL& LDL Cholesterol

Technologist MGR Page No: 7 of 15

^{2.} As per NCEP guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is



② B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊚ +91 141 4824885 © maxcarediagnostics1@gmail.com





NAME :- Mrs. SAROJ DEVI

Age :-47 Yrs 2 Mon 25 Days

Sex :-Female Patient ID: -12234987

Date :- 26/03/2024

10:30:52

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Mr.MEDIWHEEL Company:-

Final Authentication: 26/03/2024 17:37:27

BIOCHEMISTRY

recommended

3. Low HDL levels are associated with Coronary Heart Disease due to insufficient HDL being available to participate in reverse cholesterol transport, the process by which cholesterol is eliminated fromperipheral tissues.



Technologist MGR Page No: 8 of 15



O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885 ⊕ maxcarediagnostics1@gmail.com





NAME :- Mrs. SAROJ DEVI

47 Yrs 2 Mon 25 Days

Sex :-Female

Age :-

Patient ID: -12234987

Date :- 26/03/2024

10:30:52

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp:-

Company:-

Mr.MEDIWHEEL

Final Authentication: 26/03/2024 17:37:27

BIOCHEMISTRY

LIVER PROFILE WITH GGT			
SERUM BILIRUBIN (TOTAL) Methord:- DMSO/Diazo	0.70	mg/dL	Infants: 0.2-8.0 mg/dL Adult - Up to - 1.2 mg/dL
SERUM BILIRUBIN (DIRECT) Methord:- DMSO/Diazo	0.20	mg/dL	Up to 0.40 mg/dL
SERUM BILIRUBIN (INDIRECT) Methord:- Calculated	0.50	mg/dl	0.30-0.70
SGOT Methord:- IFCC	219	U/L	0.0 - 40.0
SGPT Methord:- IFCC	26.8	U/L	0.0 - 35.0
SERUM ALKALINE PHOSPHATASE Methord:- DGKC - SCE	101.20	U/L	64.00 - 306.00

InstrumentName: MISPA PLUS Interpretation: Measurements of alkaline phosphatase are of use in the diagnosis, treatment and investigation of hepatobilary disease and in bone disease associated with increased osteoblastic activity. Alkaline phosphatase is also used in the diagnosis of parathyroid and intestinal disease.

SERUM GAMMA GT

Methord:- Szasz methodology Instrument Name Randox Rx Imola

22.30

U/L

5.00 - 32.00

Interpretation. Elevations in GGT levels are seen earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and

metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or post-

hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal) are observed with infectious hepatitis.

SERUM TOTAL PROTEIN Methord:- Direct Biuret Reagent	6.58	g/dl	6.00 - 8.40
SERUM ALBUMIN Methord:- Bromocresol Green	4.21	g/dl	3.50 - 5.50
SERUM GLOBULIN Methord:- CALCULATION	2.37	gm/dl	2.20 - 3.50
A/G RATIO	1.78		1.30 - 2.50

Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

Note: These are group of tests that can be used to detect the presence of liver disease, distinguish among different types of liver disorders, gauge the extent of known liver damage, and monitor the response to treatment. Most liver diseases cause only mild symptoms initially, but these diseases must be detected early. Some tests are associated with functionality (e.g., albumin), some with cellular integrity (e.g., transaminase), and some with conditions linked to the biliary tract (gamma-glutamyl transferase and alkaline phosphatase). Conditions with elevated levels of ALT and AST include hepatitis A,B,C, paracetamol toxicity etc. Several biochemical tests are useful in the evaluation and management of patients with hepatic dysfunction. Some or all of these measurements are also carried out (usually about twice a year for routine cases) on those individuals taking certain medications, such as

Technologist age No: 9 of 15



O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885 maxcarediagnostics1@gmail.com





NAME :- Mrs. SAROJ DEVI

Patient ID: -12234987

Date :- 26/03/2024

10:30:52

Age :-

47 Yrs 2 Mon 25 Days

Ref. By Doctor:-BANK OF BARODA

Sex :-Female

Lab/Hosp:-Company :-

Mr.MEDIWHEEL

Final Authentication: 26/03/2024 17:37:27

BIOCHEMISTRY

RFT / KFT WITH ELECTROLYTES

SERUM UREA Methord:- Urease/GLDH 33.20

mg/dl

10.00 - 50.00

InstrumentName: HORIBA CA 60 Interpretation: Urea measurements are used in the diagnosis and treatment of certain renal and metabolic

diseases.

SERUM CREATININE

Methord:- Jaffe's Method

0.89

mg/dl

Males: 0.6-1.50 mg/dl

Females: 0.6 -1.40 mg/dl

Interpretation:

Creatinine is measured primarily to assess kidney function and has certain advantages over the measurement of urea. The plasma level of creatinine is relatively independent of protein ingestion, water intake, rate of urine production and exercise. Depressed levels of plasma creatinine are rare and not

clinically significant. SERUM URIC ACID

4.21

mg/dl

2.40 - 7.00

InstrumentName: HORIBA YUMIZEN CA60 Daytona plus Interpretation: Elevated Urate: High purine diet, Alcohol· Renal insufficiency, Drugs, Polycythaemia vera, Malignancies, Hypothyroidism, Rare enzyme defects , Downs syndrome, Metabolic syndrome, Pregnancy, Gout.

SODIUM Methord:- ISE	139.7	mmol/L	135.0 - 150.0
POTASSIUM Methord:- ISE	3.98	mmol/L	3.50 - 5.50
CHLORIDE Methord:- ISE	101.7	mmol/L	94.0 - 110.0
SERUM CALCIUM Methord:- Arsenazo III Method	10.10	mg/dL	8.80 - 10.20

InstrumentName:MISPA PLUS Interpretation: Serum calcium levels are believed to be controlled by parathyroid hormone and vitamin D. Increases in serum PTH or vitamin D are usually associated with hypercalcemia . Hypocalcemia may be observed in hypoparathyroidism, nephrosis and pancreatitis.

SERUM TOTAL PROTEIN Methord:- Direct Biuret Reagent	6.58	g/dl	6.00 - 8.40
SERUM ALBUMIN Methord:- Bromocresol Green	4.21	g/dl	3.50 - 5.50
SERUM GLOBULIN Methord:- CALCULATION	2.37	gm/dl	2.20 - 3.50
A/G RATIO	1.78		1.30 - 2.50

Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of dis

'iver, kidney and

Technologist age No: 10 of 15



O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⑥ +91 141 4824885 ⑤ maxcarediagnostics1@gmail.com





NAME :- Mrs. SAROJ DEVI

47 Yrs 2 Mon 25 Days Age :-

Female Sex :-

Patient ID: -12234987

Date :- 26/03/2024

10:30:52

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :-Mr.MEDIWHEEL

Final Authentication: 26/03/2024 17:37:27

BIOCHEMISTRY

bone marrow as well as other metabolic or nutritional disorders.

INTERPRETATION

Kidney function tests are group of tests that can be used to evaluate how well the kidneys are functioning. Creatinine is a waste product that comes from protein in the diet and also comes from the normal wear and tear of muscles of the body. In blood, it is a marker of GFR. in urine, it can remove the need for 24-hourcollections for many analytes or be used as a quality assurance tool to assess the accuracy of a 24-hour collection Higher levels may be a sign that the kidneys are not working properly. As kidney disease progresses, the level of creatinine and urea in the bloodincreases. Certain drugs are nephrotoxic hence KFT is done before and after initiation of treatment with these drugs.

Low serum creatinine values are rare, they almost always reflect low muscle mass.


Apart from renal failure Blood Urea can increase in dehydration and GI bleed



Technologist age No: 11 of 15



O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023





NAME :- Mrs. SAROJ DEVI

47 Yrs 2 Mon 25 Days Age :-

Sex :-Female Patient ID:-12234987

Date :- 26/03/2024

10:30:52

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company:-

Mr.MEDIWHEEL

Final Authentication: 26/03/2024 17:37:27

CLINICAL PATHOLOGY

URINE SUGAR (FASTING)
Collected Sample Received

Nil

Nil



Technologist MGR Page No: 13 of 15



P3 HEALTH SOLUTIONS L

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885 ⊕ maxcarediagnostics1@gmail.com





NAME :- Mrs. SAROJ DEVI

Age :-47 Yrs 2 Mon 25 Days

Female Sex :-

Patient ID: -12234987

Date :- 26/03/2024

10:30:52

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company:-

Mr.MEDIWHEEL

Final Authentication: 26/03/2024 17:37:27

TOTAL THYROID PROFILE

IMMUNOASSAY

Test Name	Value	Unit	Biological Ref Interval
THYROID-TRIIODOTHYRONINE T3 Methord:- ECLIA	0.91	ng/mL	0.70 - 2.04
THYROID - THYROXINE (T4) Methord:- ECLIA	6.44	ug/dl	5.10 - 14.10
TSH Methord:- ECLIA	1.982	μIU/mL	0.350 - 5.500

⁴th Generation Assay, Reference ranges vary between laboratories

PREGNANCY - REFERENCE RANGE for TSH IN ulU/mL (As per American Thyroid Association)

1st Trimester: 0.10-2.50 uIU/mL 2nd Trimester: 0.20-3.00 uIU/mL 3rd Trimester: 0.30-3.00 uIU/mL

The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy.

NOTE-TSH levels are subject to circardian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result.

INTERPRETATION

- 1.Primary hyperthyroidism is accompanied by ↑serum T3 & T4 values along with ↓ TSH level.
- 2.Primary hypothyroidism is accompanied by ↓ serum T3 and T4 values & ↑serum TSH levels
- Normal T4 levels accompanied by ↑ T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis
- 4. Normal or 1 T3 & †T4 levels indicate T4 Thyrotoxicosis (problem is conversion of T4 to T3)
- 5.Normal T3 & T4 along with \(\tau \) TSH indicate mild / Subclinical Hyperthyroidism
- . COMMENTS: Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test.

Disclaimer-TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age, and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly

. Reference ranges are from Teitz fundamental of clinical chemistry 8th ed (2018

Test performed by Instrument: Beckman coulter Dxi 800

Note: The result obtained relate only to the sample given/ received & tested. A single test result is not always indicative of a disease, it has to be correlated with clinical data for interpretation.

*** End of Report ***

Technologist MGR Page No: 15 of 15

MD (Pathology)

RMC No. 17226



O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⑥ +91 141 4824885 ② maxcarediagnostics1@gmail.com





NAME :- Mrs. SAROJ DEVI

47 Yrs 2 Mon 25 Days

Sex :-Female

Age :-

Patient ID: -12234987

Date :- 26/03/2024

10:30:52

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp:-

Company:-

Mr.MEDIWHEEL

Final Authentication: 26/03/2024 17:37:27

CLINICAL PATHOLOGY


Test Name	Value	Unit	Biological Ref Interval
Urine Routine PHYSICAL EXAMINATION			
PHYSICAL EXAMINATION COLOUR APPEARANCE CHEMICAL EXAMINATION REACTION(PH) SPECIFIC GRAVITY PROTEIN SUGAR BILIRUBIN UROBILINOGEN KETONES NITRITE MICROSCOPY EXAMINATION RBC/HPF	PALE YELL Clear 5.0 1.030 NIL NIL NEGATIVE NORMAL NEGATIVE NEGATIVE	OW /HPF	PALE YELLOW Clear 5.0 - 7.5 1.010 - 1.030 NIL NIL NEGATIVE NORMAL NEGATIVE NEGATIVE NEGATIVE
WBC/HPF EPITHELIAL CELLS CRYSTALS/HPF CAST/HPF AMORPHOUS SEDIMENT BACTERIAL FLORA YEAST CELL	2-3 2-3 ABSENT ABSENT ABSENT ABSENT ABSENT	/HPF /HPF	2-3 2-3 ABSENT ABSENT ABSENT ABSENT ABSENT
OTHER *	ABSENT		

Technologist MGR Page No: 12 of 15



B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885 ⊕ maxcarediagnostics1@gmail.com



ULTRASOUND OF WHOLE ABDOMEN

Liver is of normal size (14.2 cm) with increased echotexture. No focal space occupying lesion is seen within liver parenchyma. Intrahepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is well distended. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape (10.0 cm). Echotexture is normal. No focal lesion is seen.

Kidneys are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. Collecting system does not show any dilatation or calculus.

Right kidney is measuring approx. 10.9 x 4.0 cm.

Left kidney is measuring approx. 9.9 x 4.7 cm.

A simple, well-defined cortical cyst measuring 3.0 x 3.0 cm is noted in mid pole.

Urinary bladder does not show any calculus or mass lesion.

Uterus is anteverted and normal in size (measuring approx. 7.5 x 3.4 x 3.3 cm).

Myometrium shows normal echo-pattern. No focal space occupying lesion is seen. Endometrial echo is normal. Endometrial thickness is 5.4 mm.

Both ovaries are visualized and are normal. No adnexal mass lesion is seen.

No enlarged nodes are visualized. No retro-peritoneal lesion is identified. No significant free fluid is seen in pouch of Douglas.

IMPRESSION:

- Grade I fatty liver.
- Rest no significant abnormality is detected.


Shalln'

DR.SHALINI GOEL

M.B.B.S, D.N.B (Radiodiagnosis)

RMC no.: 21954

MBBS, DNB (- o ologist) RMC No. 21954 -3 Health Solutions LLP






128541925461287/Mrs Saroj Devi #P3 HEALTH SOLUTIONS LLP B-14, Vidhyadhar nahar , Jaipur lef.: BANK OF BARODA Test Date: 26-Mar-2024(2:23:36 P) Notch: 50Hz 0.05Hz - 35Hz iems (۲) Lta 47Yrs-11Months/Female Kgs/31 Cms 10mm/mV BP: 25mm/Sec mmHg

HR: 74 bpm

PR Interval: 144 ms QRS Duration: 86 ms

QT/QTc: 382/424ms P-QRS-T Axis: 55 - 67 - 25 (Deg)



B-14, Vidhyadhar Enclave-2, Vidhyadhar Nagar, Jaipur

oummary

Date: 26-Mar-2024 02:25:57 PM 10238594/MRS SAROJ DEVI 47 Yrs/Female 0 Kg/0 Cms

Stage 2 ExStart Findings: Supine ¥ Advice/Comments: Recovery Stage Medication: Nil Recovery Recovery Recovery Recovery PeakEx Stage 1 Objective: Ref. By : BANK OF BARODA Standing Max WorkLoad attained :7.7(Fair Effort Tolerance) Max BP : 150/85(mmHg) Max HR Attained **Exercise Time** StageTime PhaseTime Speed 5:00 4:00 3:00 2:00 0:34 3:01 3:01 1:00 6:35 6:02 3:02 :06:34 158 bpm 91% of Max Predictable HR 173 0.0 0.0 0.0 0.0 2.5 Grade 12.0 10.0 0.0 0.0 7.7 1.0 1.0 7.1 .0 4.7 1.0 **METS** . .2 .0 0 .0 131 158 145 # H.R. 128 (bpm) 93 97 89 77 90 73 79 Protocol : BRUCE History : Nil 140/85 140/85 130/80 120/80 120/80 130/80 150/85 120/80 120/80 140/85 120/80 120/80 B.P. (mmHg) Nogeraline R.P.P. 203 111 221 126 124 80 196 166 92 2 87 85 94 PVC Comments 3 R M -1.1 PeakEx PreEx V1 ۲, -0.2 AM avF avR MBBS, DIP. CARDIO (ESCORTS)
D.E.M. (RCGP-UK) **4** avL 16 **V**5 3 **V2** 4 Dr. Naresh Kumar Mohanka **TTS** Merymon 1 mm Money w homem -WIMMIN my from 0.5 mm/Div 12 15 18 21 Min.



B-14, Vidhyadhar Enclave-2, Vidhyadhar Nagar, Jaipur

TO DEALID SOLUTIONS LLF

(1) 6:01 2.5 mph (2) 3:01 12.0 % 145 bpm140/85 (1) 3:01 1.7 mph (2) 3:01 10.0 % 128 bpm130/80 (1) 0:00 0.0 mph (2) 0:00 0.0 % 73 bpm 120/80 (1) 0:00 0.0 mph (2) 0:00 0.0 % 71 bpm 120/80 Standing
(1) 0:00 0.0 mph (2) 0:00 0.0 %
79 bpm 120/80 (1) 0:00 0.0 mph (2) 0:00 0.0 % 77 bpm 120/80 Stage 2 10238594/MRS SAROJ DEVI 47 Yrs/Female 0 Kg/0 Cms Date: 26-Mar-2024 02:25:57 PM Stage 1 ¥ Supine 0.0 0.0 -0.1 0.0 0.0 ŧ -1.4 -0.1 0.0 -0.9 -0.2 -0.2 -0.2 H 0.0 -0.8 0.2 -0.2 -0.2 H 0.1 0.7 0.5 0.0 0.2 avR -0.2 -0.1 0.0 0.0 0.3 0.0 0.1 avL 0.2 0.0 0.1 -0.2 -0.2 -0.8 -0.2 avF 0.1 -0.5 -0.2 0.0 0.0 0.0 -0.2 -0.1 ****1 -0.2 0.1 0.6 0.3 -0.3 -0.5 -0.2 0.0 **Y2** 0.4 0.4 0.0 0.2 -0.3 0.2 0.2 **V**3 0.0 3.0 -0.4 -0.1 -0.1 0.5 **4**4 Ŧ -1.5 -0.2 -0.3 -0.6 -0.5 -0.3 -0.1 -0.3 55

0.1


-0.3 -0.3 -0.1 -0.2 -0.2 -0.1 -0.2

Ŧ

٧6

-0.9

Average







P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

- B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023
- ⊕ +91 141 4824885
 □ maxcarediagnostics1@gmail.com



NAME:	MRS. SAROJ SHARMA		AGE	47 YRS/F
REF.BY	BANK OF BARODA	3€3	DATE	26/03/2024

CHEST X-RAY (PA VIEW)

Bilateral lung fields appear clear.

Bilateral costo-phrenic angles appear clear.

Cardiothoracic ratio is normal.

Thoracic soft tissue and skeletal system appear unremarkable.

Soft tissue shadows appear normal.

IMPRESSION: No significant abnormality is detected.



DR.SHALINI GOEL

M.B.B.S, D.N.B (Radiodiagnosis)

RMC No.: 21954

