



Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.NEERAJ KUMAR	Registered On	: 15/Sep/2024 10:43:27
Age/Gender	: 32 Y 2 M 2 D /M	Collected	: 2024-09-15 11:06:34
UHID/MR NO	: ALDP.0000072040	Received	: 2024-09-15 11:06:34
Visit ID	: ALDP0217652425	Reported	: 19/Sep/2024 12:45:10
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

## DEPARTMENT OF CARDIOLOGY-ECG MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

## ECG / EKG

1. Machnism, Rhythm	Sinus, Regular	
2. Atrial Rate	78	/mt
3. Ventricular Rate	78	/mt
4. P - Wave	Normal	
5. P R Interval	Normal	
6. Q R S Axis : R/S Ratio : Configuration :	Normal Normal Normal	
7. Q T c Interval	Normal	
8. S - T Segment	Normal	
9. T – Wave <u>FINAL IMPRESSION</u>	Normal	_

ECG Within Normal Limits: Sinus Rhythm. Baseline artefacts. Please correlate clinically.











Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.NEERAJ KUMAR	Registered On	: 15/Sep/2024 10:43:26
Age/Gender	: 32 Y 2 M 2 D /M	Collected	: 15/Sep/2024 11:53:33
UHID/MR NO	: ALDP.0000072040	Received	: 15/Sep/2024 12:44:18
Visit ID	: ALDP0217652425	Reported	: 15/Sep/2024 15:06:10
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

## DEPARTMENT OF HAEMATOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
Blood Group (ABO & Rh typing), Blood				
Blood Group	В			ERYTHROCYTE MAGNETIZED TECHNOLOGY / TUBE AGGLUTINA
Rh ( Anti-D)	POSITIVE			ERYTHROCYTE MAGNETIZED TECHNOLOGY / TUBE AGGLUTINA
Complete Blood Count (CBC) , Whole Blood				
Haemoglobin	12.90	g/dl	1 Day- 14.5-22.5 g/dl 1 Wk- 13.5-19.5 g/dl 1 Mo- 10.0-18.0 g/dl 3-6 Mo- 9.5-13.5 g/dl 0.5-2 Yr- 10.5-13.5 g/dl 2-6 Yr- 11.5-15.5 g/dl 6-12 Yr- 11.5-15.5 g/dl 12-18 Yr 13.0-16.0 g/dl Male- 13.5-17.5 g/dl Female- 12.0-15.5 g/dl	COLORIMETRIC METHOD (CYANIDE-FREE REAGENT)
TLC (WBC) <u>DLC</u>	5,700.00	/Cu mm	4000-10000	IMPEDANCE METHOD
Polymorphs (Neutrophils )	60.00	%	40-80	FLOW CYTOMETRY
Lymphocytes	31.00	%	20-40	FLOW CYTOMETRY
Monocytes	4.00	%	2-10	FLOW CYTOMETRY
Eosinophils	5.00	%	1-6	FLOW CYTOMETRY
Basophils <b>ESR</b>	0.00	%	< 1-2	FLOW CYTOMETRY
Observed	16.00	MM/1H	10-19 Yr 8.0 20-29 Yr 10.8 30-39 Yr 10.4 40-49 Yr 13.6 50-59 Yr 14.2 60-69 Yr 16.0 70-79 Yr 16.5 80-91 Yr 15.8	



Page 2 of 14







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.NEERAJ KUMAR	Registered On	: 15/Sep/2024 10:43:26
Age/Gender	: 32 Y 2 M 2 D /M	Collected	: 15/Sep/2024 11:53:33
UHID/MR NO	: ALDP.0000072040	Received	: 15/Sep/2024 12:44:18
Visit ID	: ALDP0217652425	Reported	: 15/Sep/2024 15:06:10
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

## DEPARTMENT OF HAEMATOLOGY

## MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
			Pregnancy Early gestation - 48 (62 if anaemic) Leter gestation - 70 (95 if anaemic)	
Corrected	-	Mm for 1st hr.	< 9	
PCV (HCT)	37.00	%	40-54	
Platelet count				
Platelet Count	1.52	LACS/cu mm	1.5-4.0	ELECTRONIC IMPEDANCE/MICROSCOPIC
PDW (Platelet Distribution width)	16.60	fL	9-17	ELECTRONIC IMPEDANCE
P-LCR (Platelet Large Cell Ratio)	-	%	35-60	ELECTRONIC IMPEDANCE
PCT (Platelet Hematocrit)	0.20	%	0.108-0.282	ELECTRONIC IMPEDANCE
MPV (Mean Platelet Volume)	13.70	fL	6.5-12.0	ELECTRONIC IMPEDANCE
RBC Count				
RBC Count	4.06	Mill./cu mm	4.2-5.5	ELECTRONIC IMPEDANCE
Blood Indices (MCV, MCH, MCHC)				
MCV	93.40	fl	80-100	CALCULATED PARAMETER
MCH	31.70	pg	27-32	CALCULATED PARAMETER
MCHC	34.00	%	30-38	CALCULATED PARAMETER
RDW-CV	13.00	%	11-16	ELECTRONIC IMPEDANCE
RDW-SD	46.70	fL	35-60	ELECTRONIC IMPEDANCE
Absolute Neutrophils Count	3,420.00	/cu mm	3000-7000	
Absolute Eosinophils Count (AEC)	285.00	/cu mm	40-440	

AS

View Reports on

Chandan 24x7 App

Dr.Akanksha Singh (MD Pathology)

Page 3 of 14









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.NEERAJ KUMAR	Registered On	: 15/Sep/2024 10:43:27
Age/Gender	: 32 Y 2 M 2 D /M	Collected	: 15/Sep/2024 11:53:33
UHID/MR NO	: ALDP.0000072040	Received	: 15/Sep/2024 12:44:18
Visit ID	: ALDP0217652425	Reported	: 15/Sep/2024 14:07:18
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

## DEPARTMENT OF BIOCHEMISTRY

## MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interva	al Method
<b>GLUCOSE FASTING</b> , <i>Plasma</i> Glucose Fasting	90.50	10	00 Normal 0-125 Pre-diabetes <b>26 Diabetes</b>	GOD POD

## Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions.b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential.c) I.G.T = Impaired Glucose Tolerance.

**CLINICAL SIGNIFICANCE:-** Glucose is the major source of energy in the body. Lack of insulin or resistance to it section at the cellular level causes diabetes. Therefore, the blood glucose levels are very high. Elevated serum glucose levels are observed in diabetes mellitus and may be associated with pancreatitis, pituitary or thyroid dysfunction and liver disease. Hypoglycaemia occurs most frequently due to over dosage of insulin.

Glucose PP	119.00	mg/dl	<140 Normal	GOD POD
Sample:Plasma After Meal			140-199 Pre-diabetes	
			>200 Diabetes	

## Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions.b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential.c) I.G.T = Impaired Glucose Tolerance.

## GLYCOSYLATED HAEMOGLOBIN (HBA1C), EDTA BLOOD

Glycosylated Haemoglobin (HbA1c)	5.20	% NGSP	HPLC (NGSP)
Glycosylated Haemoglobin (HbA1c)	32.90	mmol/mol/IFCC	
Estimated Average Glucose (eAG)	102	mg/dl	

## Interpretation:

## NOTE:-

• eAG is directly related to A1c.



Page 4 of 14





Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.NEERAJ KUMAR	Registered On	: 15/Sep/2024 10:43:27
Age/Gender	: 32 Y 2 M 2 D /M	Collected	: 15/Sep/2024 11:53:33
UHID/MR NO	: ALDP.0000072040	Received	: 15/Sep/2024 12:44:18
Visit ID	: ALDP0217652425	Reported	: 15/Sep/2024 14:07:18
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

## **DEPARTMENT OF BIOCHEMISTRY**

## MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
-----------	--------	------	--------------------	--------

- An A1c of 7% -the goal for most people with diabetes-is the equivalent of an eAG of 154 mg/dl.
- eAG may help facilitate a better understanding of actual daily control helping you and your health care provider to make necessary changes to your diet and physical activity to improve overall diabetes mnagement.

The following ranges may be used for interpretation of results. However, factors such as duration of diabetes, adherence to therapy and the age of the patient should also be considered in assessing the degree of blood glucose control.

Haemoglobin A1C (%)NGSP	mmol/mol / IFCC Unit	eAG (mg/dl)	<b>Degree of Glucose Control Unit</b>
> 8	>63.9	>183	Action Suggested*
7-8	53.0 -63.9	154-183	Fair Control
< 7	<63.9	<154	Goal**
6-7	42.1 -63.9	126-154	Near-normal glycemia
< 6%	<42.1	<126	Non-diabetic level

\*High risk of developing long term complications such as Retinopathy, Nephropathy, Neuropathy, Cardiopathy, etc. \*\*Some danger of hypoglycemic reaction in Type 1diabetics. Some glucose intolerant individuals and "subclinical" diabetics may demonstrate HbA1C levels in this area.

N.B. : Test carried out on Automated VARIANT II TURBO HPLC Analyser.

## **<u>Clinical Implications:</u>**

\*Values are frequently increased in persons with poorly controlled or newly diagnosed diabetes.

\*With optimal control, the HbA 1c moves toward normal levels.

\*A diabetic patient who recently comes under good control may still show higher concentrations of glycosylated hemoglobin. This level declines gradually over several months as nearly normal glycosylated \*Increases in glycosylated hemoglobin occur in the following non-diabetic conditions: a. Iron-deficiency anemia b. Splenectomy

c. Alcohol toxicity d. Lead toxicity

\*Decreases in A 1c occur in the following non-diabetic conditions: a. Hemolytic anemia b. chronic blood loss

\*Pregnancy d. chronic renal failure. Interfering Factors:

\*Presence of Hb F and H causes falsely elevated values. 2. Presence of Hb S, C, E, D, G, and Lepore (autosomal recessive mutation resulting in a hemoglobinopathy) causes falsely decreased values.

BUN (Blood Urea Nitrogen)	14.20	mg/dL	7.0-23.0	CALCULATED
Sample:Serum				



View Reports on

Chandan 24x7 App







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Detterst Nieures						
Patient Name	: Mr.NEERAJ KUMAR		Registered	l On	: 15/Sep/2024 10:	
Age/Gender	: 32 Y 2 M 2 D /M		Collected		: 15/Sep/2024 11:	
UHID/MR NO Visit ID	: ALDP.0000072040 : ALDP0217652425		Received Reported		: 15/Sep/2024 12: : 15/Sep/2024 14:	
Ref Doctor	: Dr. MEDIWHEEL-ARCC CARE LTD -	DFEMI HEALTH	Status		: Final Report	07.10
		DEPARTMEN	T OF BIOCH	EMISTR	Y	
	MEDIWH	HEEL BANK OF	BARODA M	ALE AB	OVE 40 YRS	
Test Name		Result	U	nit	Bio. Ref. Interval	Method
	N levels can be seen in th	_	Contraintentia		le dine	
High-protein diet, De	hydration, Aging, Certain n	nedications, Burns,	Gastrointestin	nal (GI) bl	leeding.	
Low BUN levels ca	n be seen in the following	g:				
Low-protein diet, ove	erhydration, Liver disease.					
Creatinine Sample:Serum		1.02	mg/dl	0.7-1.3		DDIFIED JAFFES
<i>Interpretation:</i> The significance of si mass will have a high absolute creatinine co could be affected mile	ngle creatinine value must b ter creatinine concentration. oncentration. Serum creatini dly and may result in anoma	be interpreted in lig The trend of serun ine concentrations	ht of the patien n creatinine co may increase v	nts muscle ncentratio when an A	mass. A patient with ns over time is more i .CE inhibitor (ACE) i	a greater muscle important than is taken. The assay
<i>Interpretation:</i> The significance of si mass will have a high absolute creatinine co	ner creatinine concentration. Discontration. Serum creatini	be interpreted in lig The trend of serun ine concentrations	ht of the patien n creatinine co may increase v	nts muscle ncentratio when an A	mass. A patient with ns over time is more i CE inhibitor (ACE) i hilic antibodies, hemo	a greater muscle important than is taken. The assay
The significance of si mass will have a high absolute creatinine co could be affected mile lipemic. Jric Acid Cample:Serum Interpretation: Note:-	ner creatinine concentration. Discontration. Serum creatini	be interpreted in lig The trend of serun ine concentrations alous values if serun 5.57	ht of the patien n creatinine co may increase v m samples hav	nts muscle ncentratio when an A re heteropl	mass. A patient with ns over time is more i CE inhibitor (ACE) i hilic antibodies, hemo	a greater muscle important than is taken. The assay olyzed, icteric or
ample:Serum Interpretation: The significance of si mass will have a high absolute creatinine co could be affected mile lipemic. Jric Acid ample:Serum Interpretation: Note:- Elevated uric acid l	er creatinine concentration. oncentration. Serum creatini dly and may result in anoma	be interpreted in lig The trend of serun ine concentrations alous values if serun 5.57 <b>5.57</b>	ht of the patien n creatinine co may increase w m samples haw mg/dl	nts muscle ncentratio when an A re heteropl 3.4-7.0	mass. A patient with ns over time is more i CE inhibitor (ACE) i hilic antibodies, hemo	a greater muscle important than is taken. The assay olyzed, icteric or
The significance of si mass will have a high absolute creatinine co could be affected mild lipemic. Jric Acid Sample:Serum Interpretation: Note:- Elevated uric acid I	er creatinine concentration. oncentration. Serum creatini dly and may result in anoma levels can be seen in the f otein diet, alcohol), Chronic	be interpreted in lig The trend of serun ine concentrations alous values if serun 5.57 <b>5.57</b>	ht of the patien n creatinine co may increase w m samples haw mg/dl	nts muscle ncentratio when an A re heteropl 3.4-7.0	mass. A patient with ns over time is more i CE inhibitor (ACE) i hilic antibodies, hemo	a greater muscle important than is taken. The assay olyzed, icteric or
Interpretation: The significance of si mass will have a high absolute creatinine co could be affected mild lipemic. Jric Acid ample:Serum Interpretation: Note:- Elevated uric acid I Drugs, Diet (high-pro FT (WITH GAMMA	er creatinine concentration. oncentration. Serum creatini dly and may result in anoma levels can be seen in the f otein diet, alcohol), Chronic	be interpreted in lig The trend of serun ine concentrations alous values if serun 5.57 <b>5.57</b>	ht of the patien n creatinine co may increase w m samples haw mg/dl	nts muscle ncentratio when an A re heteropl 3.4-7.0	mass. A patient with ns over time is more i CE inhibitor (ACE) i hilic antibodies, hemo UR	a greater muscle important than is taken. The assay olyzed, icteric or
Interpretation: The significance of si mass will have a high absolute creatinine co could be affected mild lipemic. Jric Acid ample:Serum Interpretation: Note:- Elevated uric acid I Drugs, Diet (high-pro FT (WITH GAMIMA SGOT / Aspartate Am SGPT / Alanine Amir	er creatinine concentration. oncentration. Serum creatini dly and may result in anoma levels can be seen in the f otein diet, alcohol), Chronic ( GT) , <i>Serum</i> ninotransferase (AST)	be interpreted in lig The trend of serum ine concentrations alous values if serue 5.57 <b>Collowing:</b> kidney disease, H 32.40 <b>45.70</b>	ht of the patien n creatinine co may increase v m samples hav mg/dl ypertension, C U/L U/L	nts muscle ncentratio when an A /e heteropl 3.4-7.0 Desity. < 35 < 40	mass. A patient with ns over time is more i CE inhibitor (ACE) i hilic antibodies, hemo UR UR	a greater muscle important than is taken. The assay Jyzed, icteric or ICASE C WITHOUT P5P C WITHOUT P5P
Interpretation: The significance of si mass will have a high absolute creatinine co could be affected mild lipemic. Jric Acid ample:Serum Interpretation: Note:- Elevated uric acid I Drugs, Diet (high-pro FT (WITH GAMIMA SGOT / Aspartate Amir Gamma GT (GGT)	er creatinine concentration. oncentration. Serum creatini dly and may result in anoma levels can be seen in the f otein diet, alcohol), Chronic ( GT) , <i>Serum</i> ninotransferase (AST)	be interpreted in lig The trend of serun ine concentrations alous values if serun 5.57 <b>Collowing:</b> kidney disease, H 32.40 <b>45.70</b> 27.80	ht of the patien n creatinine co may increase v m samples hav mg/dl ypertension, C U/L U/L IU/L	nts muscle ncentratio when an A re heteropl 3.4-7.0 Desity. >Desity. < 35 < 40 11-50	mass. A patient with ns over time is more i CE inhibitor (ACE) i nilic antibodies, hemo UR UR IFC IFC	a greater muscle important than is taken. The assay Jyzed, icteric or ICASE C WITHOUT P5P C WITHOUT P5P TIMIZED SZAZING
Interpretation: The significance of si mass will have a high absolute creatinine co could be affected mild lipemic. Jric Acid ample:Serum Interpretation: Note:- Elevated uric acid I Drugs, Diet (high-pro FT (WITH GAMIMA SGOT / Aspartate An SGPT / Alanine Amir Gamma GT (GGT) Protein	er creatinine concentration. oncentration. Serum creatini dly and may result in anoma levels can be seen in the f otein diet, alcohol), Chronic ( GT) , <i>Serum</i> ninotransferase (AST)	be interpreted in lig The trend of serun ine concentrations alous values if serun 5.57 <b>Collowing:</b> kidney disease, H 32.40 <b>45.70</b> 27.80 <b>6.06</b>	ht of the patien n creatinine co may increase v m samples hav mg/dl ypertension, C U/L U/L U/L IU/L gm/dl	nts muscle ncentratio when an A re heteropl 3.4-7.0 Desity. < 35 < 40 11-50 6.2-8.0	mass. A patient with ns over time is more i CE inhibitor (ACE) i hilic antibodies, hemo UR IFC IFC OP BIL	a greater muscle important than is taken. The assay olyzed, icteric or ICASE C WITHOUT P5P C WITHOUT P5P TIMIZED SZAZING JRET
Interpretation: The significance of si mass will have a high absolute creatinine co could be affected mild lipemic. Jric Acid ample:Serum Interpretation: Note:- Elevated uric acid I Drugs, Diet (high-pro FT (WITH GAMIMA SGOT / Aspartate Amir SGPT / Alanine Amir Gamma GT (GGT)	er creatinine concentration. oncentration. Serum creatini dly and may result in anoma levels can be seen in the f otein diet, alcohol), Chronic ( GT) , <i>Serum</i> ninotransferase (AST)	be interpreted in lig The trend of serun ine concentrations alous values if serun 5.57 <b>Collowing:</b> kidney disease, H 32.40 <b>45.70</b> 27.80	ht of the patien n creatinine co may increase v m samples hav mg/dl ypertension, C U/L U/L IU/L	nts muscle ncentratio when an A re heteropl 3.4-7.0 Desity. >Desity. < 35 < 40 11-50	mass. A patient with ns over time is more i CE inhibitor (ACE) i hilic antibodies, hemo UR IFC IFC OP BIU B.C	a greater muscle important than is taken. The assay olyzed, icteric or ICASE C WITHOUT P5P C WITHOUT P5P TIMIZED SZAZING JRET



Chandan 24x7 App







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.NEERAJ KUMAR	Registered On	: 15/Sep/2024 10:43:27
Age/Gender	: 32 Y 2 M 2 D /M	Collected	: 15/Sep/2024 11:53:33
UHID/MR NO	: ALDP.0000072040	Received	: 15/Sep/2024 12:44:18
Visit ID	: ALDP0217652425	Reported	: 15/Sep/2024 14:07:18
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

# **DEPARTMENT OF BIOCHEMISTRY**

## MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	U	Init Bio. Ref. Ir	nterval Method
Alkaline Phosphatase (Total)	102.00	U/L	42.0-165.0	PNP/AMP KINETIC
Bilirubin (Total)	0.99	mg/dl	0.3-1.2	<b>JENDRASSIK &amp; GROF</b>
Bilirubin (Direct)	0.30	mg/dl	< 0.30	<b>JENDRASSIK &amp; GROF</b>
Bilirubin (Indirect)	0.69	mg/dl	< 0.8	JENDRASSIK & GROF
LIPID PROFILE (MINI), Serum				
Cholesterol (Total)	192.00	mg/dl	<200 Desirable 200-239 Borderline > 240 High	CHOD-PAP e High
HDL Cholesterol (Good Cholesterol)	63.30	mg/dl	30-70	DIRECT ENZYMATIC
LDL Cholesterol (Bad Cholesterol)	83	mg/dl	< 100 Optimal 100-129 Nr. Optimal/Above Op 130-159 Borderling 160-189 High > 190 Very High	
VLDL	46.08	mg/dl	10-33	CALCULATED
Triglycerides	230.40	mg/dl	< 150 Normal 150-199 Borderline 200-499 High >500 Very High	GPO-PAP e High

AS

View Reports on

Dr.Akanksha Singh (MD Pathology)









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.NEERAJ KUMAR	Registered On	: 15/Sep/2024 10:43:27
Age/Gender	: 32 Y 2 M 2 D /M	Collected	: 15/Sep/2024 16:04:15
UHID/MR NO	: ALDP.0000072040	Received	: 15/Sep/2024 17:04:25
Visit ID	: ALDP0217652425	Reported	: 15/Sep/2024 18:50:38
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

# DEPARTMENT OF CLINICAL PATHOLOGY

## MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
URINE EXAMINATION, ROUTINE , $\boldsymbol{\upsilon}$	Irine			
Color	PALE YELLOW			
Specific Gravity	1.020			
Reaction PH	Acidic ( 5.0 )			DIPSTICK
Appearance	CLEAR			
Protein	ABSENT	mg %	< 10 Absent 10-40 (+) 40-200 (++) 200-500 (+++) > 500 (++++)	DIPSTICK
Sugar	ABSENT	gms%	< 0.5 (+) 0.5-1.0 (++) 1-2 (+++) > 2 (++++)	DIPSTICK
Ketone	ABSENT	mg/dl	Serum-0.1-3.0 Urine-0.0-14.0	BIOCHEMISTRY
Bile Salts	ABSENT			
Bile Pigments	ABSENT			
Bilirubin	ABSENT			DIPSTICK
Leucocyte Esterase	ABSENT			DIPSTICK
Urobilinogen(1:20 dilution)	ABSENT			
Nitrite	ABSENT			DIPSTICK
Blood	ABSENT			DIPSTICK
Microscopic Examination:				
Epithelial cells	0-2/h.p.f			MICROSCOPIC EXAMINATION
Pus cells	0-2/h.p.f			
RBCs	ABSENT			MICROSCOPIC EXAMINATION
Cast	ABSENT			
Crystals	ABSENT			MICROSCOPIC EXAMINATION
Others	ABSENT			

Urine Microscopy is done on centrifuged urine sediment.









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.NEERAJ KUMAR	Registered On	: 15/Sep/2024 10:43:27
Age/Gender	: 32 Y 2 M 2 D /M	Collected	: 15/Sep/2024 16:04:15
UHID/MR NO	: ALDP.0000072040	Received	: 15/Sep/2024 17:04:25
Visit ID	: ALDP0217652425	Reported	: 15/Sep/2024 18:50:38
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

## **DEPARTMENT OF CLINICAL PATHOLOGY** MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

IVIEL	JIVITEEL DAINK OF DAK		ADOVE 40 TKS	
Test Name	Result	Unit	Bio. Ref. Interval	Method
SUGAR, FASTING STAGE, Urine				
Sugar, Fasting stage	ABSENT	gms%		
Interpretation:   (+) < 0.5				
SUGAR, PP STAGE, Urine				
Sugar, PP Stage	ABSENT			
Interpretation:				
(+) < 0.5 gms% (++) 0.5-1.0 gms% (+++) 1-2 gms%				

(++++) > 2 gms%

AS

View Reports on

Dr.Akanksha Singh (MD Pathology)









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.NEERAJ KUMAR	Registered On	: 15/Sep/2024 10:43:27
Age/Gender	: 32 Y 2 M 2 D /M	Collected	: 15/Sep/2024 11:53:33
UHID/MR NO	: ALDP.0000072040	Received	: 15/Sep/2024 12:44:18
Visit ID	: ALDP0217652425	Reported	: 15/Sep/2024 14:22:39
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

## DEPARTMENT OF IMMUNOLOGY

## MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
<b>PSA (Prostate Specific Antigen), Total</b> Sample:Serum	0.65	ng/mL	<4.1	CLIA

## Interpretation:

- 1. PSA is detected in the serum of males with normal, benign hypertrophic, and malignant prostate tissue.
- 2. Measurement of serum PSA levels is not recommended as a screening procedure for the diagnosis of cancer because elevated PSA levels also are observed in patients with benign prostatic hypertrophy. However, studies suggest that the measurement of PSA in conjunction with digital rectal examination (DRE) and ultrasound provide a better method of detecting prostate cancer than DRE alone<sup>-</sup>
- 3. PSA levels increase in men with cancer of the prostate, and after radical prostatectomy PSA levels routinely fall to the undetectable range.
- 4. If prostatic tissue remains after surgery or metastasis has occurred, PSA appears to be useful in detecting residual and early recurrence of tumor.
- 5. Therefore, serial PSA levels can help determine the success of prostatectomy, and the need for further treatment, such as radiation, endocrine or chemotherapy, and in the monitoring of the effectiveness of therapy.

## THYROID PROFILE - TOTAL , Serum

T3, Total (tri-iodothyronine)	119.00	ng/dl	84.61–201.7	CLIA
T4, Total (Thyroxine)	6.37	ug/dl	3.2-12.6	CLIA
TSH (Thyroid Stimulating Hormone)	1.880	μlU/mL	0.27 - 5.5	CLIA

## Interpretation:

0.3-4.5	µIU/mL	First Trimest	er
0.5-4.6	µIU/mL	Second Trim	ester
0.8-5.2	µIU/mL	Third Trimes	ter
0.5-8.9	µIU/mL	Adults	55-87 Years
0.7-27	µIU/mL	Premature	28-36 Week
2.3-13.2	µIU/mL	Cord Blood	> 37Week
0.7-64	µIU/mL	Child(21 wk	- 20 Yrs.)
1-39	µIU/mL	Child	0-4 Days
1.7-9.1	µIU/mL	Child	2-20 Week

1) Patients having low T3 and T4 levels but high TSH levels suffer from primary hypothyroidism, cretinism, juvenile myxedema or









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.NEERAJ KUMAR	Registered On	: 15/Sep/2024 10:43:27
Age/Gender	: 32 Y 2 M 2 D /M	Collected	: 15/Sep/2024 11:53:33
UHID/MR NO	: ALDP.0000072040	Received	: 15/Sep/2024 12:44:18
Visit ID	: ALDP0217652425	Reported	: 15/Sep/2024 14:22:39
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

## DEPARTMENT OF IMMUNOLOGY

## MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method

autoimmune disorders.

2) Patients having high T3 and T4 levels but low TSH levels suffer from Grave's disease, toxic adenoma or sub-acute thyroiditis.

**3**) Patients having either low or normal T3 and T4 levels but low TSH values suffer from iodine deficiency or secondary hypothyroidism.

**4**) Patients having high T3 and T4 levels but normal TSH levels may suffer from toxic multinodular goiter. This condition is mostly a symptomatic and may cause transient hyperthyroidism but no persistent symptoms.

5) Patients with high or normal T3 and T4 levels and low or normal TSH levels suffer either from T3 toxicosis or T4 toxicosis respectively.

**6)** In patients with non thyroidal illness abnormal test results are not necessarily indicative of thyroidism but may be due to adaptation to the catabolic state and may revert to normal when the patient recovers.

7) There are many drugs for eg. Glucocorticoids, Dopamine, Lithium, Iodides, Oral radiographic dyes, etc. which may affect the thyroid function tests.

**8**) Generally when total T3 and total T4 results are indecisive then Free T3 and Free T4 tests are recommended for further confirmation along with TSH levels.

Dr.Akanksha Singh (MD Pathology)









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.NEERAJ KUMAR	Registered On	: 15/Sep/2024 10:43:27
Age/Gender	: 32 Y 2 M 2 D /M	Collected	: 2024-09-15 10:55:21
UHID/MR NO	: ALDP.0000072040	Received	: 2024-09-15 10:55:21
Visit ID	: ALDP0217652425	Reported	: 15/Sep/2024 13:31:19
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

## DEPARTMENT OF X-RAY

## MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

## **X-RAY DIGITAL CHEST PA**

## <u>X-RAY REPORT</u> (300 mA COMPUTERISED UNIT SPOT FILM DEVICE) <u>CHEST P-A VIEW</u>

- Both lung field did not reveal any significant lesion.
- Costo-phrenic angles are bilaterally clear.
- Trachea is central in position.
- Cardiac size & contours are normal.
- Hilar shadows are normal.
- Soft tissue shadow appears normal.
- Bony cage is normal.

Please correlare clinically.

Crohoh

DR K N SINGH (MBBS, DMRE)



View Reports on **Chandan 24x7 App** 







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.NEERAJ KUMAR	Registered On	: 15/Sep/2024 10:43:27
Age/Gender	: 32 Y 2 M 2 D /M	Collected	: 2024-09-15 13:48:06
UHID/MR NO	: ALDP.0000072040	Received	: 2024-09-15 13:48:06
Visit ID	: ALDP0217652425	Reported	: 15/Sep/2024 13:51:50
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

## DEPARTMENT OF ULTRASOUND

## MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

## **ULTRASOUND WHOLE ABDOMEN (UPPER & LOWER)**

**LIVER**: - Normal in size (15.2 cm), shape and echogenicity. No focal lesion is seen. No intra hepatic biliary radicle dilation is seen.

**GALL BLADDER** :- Well distended. Normal wall thickness is seen. No evidence of calculus/focal mass lesion/pericholecystic fluid is seen.

**CBD** :- Normal in calibre at porta.

PORTAL VEIN: - Normal in calibre and colour uptake at porta.

**PANCREAS:** - Head is visualised, normal in size & echopattern. No evidence of ductal dilatation or calcification is seen. Rest of the pancreas is obscured by bowel gases.

SPLEEN: - Normal in size, shape and echogenicity. No evidence of mass lesion is seen.

**RIGHT KIDNEY**: - Normal in size (10.8 cm), shape and position. Cortical echogenicity is normal with maintained corticomedullary differentiation. No focal lesion or calculus is seen. Pelvicalyceal system is not dilated.

**LEFT KIDNEY**: - Normal in size (9.2 cm), shape and position. Cortical echogenicity is normal with maintained corticomedullary differentiation. **Few tiny concretions are seen.** Pelvicalyceal system is not dilated.

**URINARY BLADDER :-** Is adequately distended. No evidence of wall thickening/calculus is seen.

**PROSTATE :-** Normal in size (2.6 x 4.9 x 2.8 cm vol - 19.8 cc), shape and echo pattern.

**HIGH RESOLUTION** :- No evidence of bowel loop dilatation or abnormal wall thickening is seen. No significant retroperitoneal lymphadenopathy is seen. No free fluid is seen in the abdomen/pelvis.

## **IMPRESSION**: Left renal concretions.

Please correlate clinically

Contil

DR K N SINGH (MBBS, DMRE)











Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.NEERAJ KUMAR	Registered On	: 15/Sep/2024 10:43:28
Age/Gender	: 32 Y 2 M 2 D /M	Collected	: 2024-09-15 12:43:27
UHID/MR NO	: ALDP.0000072040	Received	: 2024-09-15 12:43:27
Visit ID	: ALDP0217652425	Reported	: 15/Sep/2024 16:54:57
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

## **DEPARTMENT OF TMT**

## MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

## Tread Mill Test (TMT)

NORMAL

\*\*\* End Of Report \*\*\*

Result/s to Follow: STOOL, ROUTINE EXAMINATION



This report is not for medico legal purpose. If clinical correlation is not established, kindly repeat the test at no additional cost within seven days

Facilities: Pathology, Bedside Sample Collection, Health Check-ups, Digital X-Ray, ECG (Bedside also), Allergy Testing, Test And Health Check-ups, Ultrasonography, Sonomammography, Bone Mineral Density (BMD), Doppler Studies, 2D Echo, CT Scan, MRI, Blood Bank, TMT, EEG, PFT, OPG, Endoscopy, Digital Mammography, Electromyography (EMG), Nerve Condition Velocity (NCV), Audiometry, Brainstem Evoked Response Audiometry (BERA), Colonoscopy, Ambulance Services, Online Booking Facilities for Diagnostics, Online Report Viewing \* 365 Days Open \*Facilities Available at Select Location Page 14 of 14



Home Sample Collection 08069366666





Dr. R K VERMA MBBS, PGDGM