

Land amas a



: :

Dr. PIYUSH GOYAL
MBBS, DMRDy Radiologist)
RMC No. 037041



- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- S +91 141 4824885 ⊠ p3healthsolutionsllp@gmail.com



### **General Physical Examination**

Date of Examination: 24/68/2029
Name: Lang Narayan Navaria Age: 42 DOB: 03/11/1981 Sex: Male
Referred By: Banda
Photo ID: Adhar (and ID#: 1750
Ht: 170 (cm) Wt: 60 (Kg)
Chest (Expiration): (cm) Abdomen Circumference: (cm)
Blood Pressure: 30/80 mm Hg PR: 78/min RR: 18/min Temp: Achaile
BMI
Eye Examination: RIE-GIGNIG, NCI3 LIE-GIGNIG, NCI3
Other:
Other:
On examination he/she appears physically and mentally fit: \\Yes\f\ No
Land away
Signature Of Examine: Name of Examinee: LAX MJ NA 120 YOUNGWART
Signature Medical Examiner: Name Medical Examiner 128.12 TYDSH CIONAL NO037041



 B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023

③ +91 141 4824885 ⊠ p3healthsolutionsllp@gmail.com



Patient ID	1224990 Patient Mob No.9166636200	Registered On	24/08/2024 11:31:46
NAME	Mr. LAXMI NARAYAN NAWARIA	Collected On	24/08/2024 12:11:44
Age / Sex	Male 42 Yrs 9 Mon 22 Days	Authorized On	25/08/2024 09:21:25
Ref. By	BANK OF BARODA	Printed On	25/08/2024 09:21:35
Lab/Hosp	Mr.MEDIWHEEL		

#### HAEMOGARAM

### **HAEMATOLOGY**

Value	Unit	Biological Ref Interval		
FULL BODY HEALTH CHECKUP ABOVE 40 MALE				
EO DALW DESCRIPTO	g/dI	13.0 - 17.0		
7.60	/cumm	4.00 - 10.00		
51.6	%	40.0 - 80.0		
44.5 H	%	20.0 - 40.0		
1.1	%	1.0 - 6.0		
2.8	%	2.0 - 10.0		
0.0	%	0.0 - 2.0		
4.81	x10^6/uL	4.50 - 5.50		
42.40	%	40.00 - 50.00		
88.0	fL	83.0 - 101.0		
28.8	pg	27.0 - 32.0		
32.7	g/dL	31.5 - 34.5		
122 L	x10^3/uL	150 - 410		
15.1 H	%	11.6 - 14.0		
	0 MALE 13.8 7.60 51.6 44.5 H 1.1 2.8 0.0 4.81 42.40 88.0 28.8 32.7 122 L	13.8 g/dL 7.60 /cumm 51.6 % 44.5 H % 1.1 % 2.8 % 0.0 % 4.81 x10^6/uL 42.40 % 88.0 fL 28.8 pg 32.7 g/dL 122 L x10^3/uL		

Technologist



 B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023

🕲 +91 141 4824885 🖂 p3healthsolutionsllp@gmail.com



Patient ID	1224990 Patient Mob No.9166636200	Registered On	24/08/2024 11:31:46
NAME	Mr. LAXMI NARAYAN NAWARIA	Collected On	24/08/2024 12:11:44
Age / Sex	Male 42 Yrs 9 Mon 22 Days	Authorized On	25/08/2024 09:21:25
Ref. By	BANK OF BARODA	Printed On	25/08/2024 09:21:35
Lab/Hosp	Mr.MEDIWHEEL		

#### HAEMATOLOGY

#### HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
Erythrocyte Sedimentation Rate (ESR) Methord:- Westergreen	<b>20</b> H	mm in 1st hr	00 - 15

The erythrocyte sedimentation rate (ESR or sed rate) is a relatively simple, inexpensive, non-specific test that has been used for many years to help detect inflammation associated with conditions such as infections, cancers, and autoimmune diseases. ESR is said to be a non-specific test because an elevated result often indicates the presence of inflammation but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other tests, such as C-reactive protein. ESR is used to help diagnose certain specific inflammatory diseases, including temporal arteritis, systemic vasculitis and polymyalgia rheumatica. (For more on these, read the article on Vasculitis.) A significantly elevated ESR is one of the main test results used to support the diagnosis. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as

Technologist



- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- 🕒 +91 141 4824885 🖾 p3healthsolutionsllp@gmail.com



Patient ID	1224990 Patient Mob No.9166636200	Registered On	24/08/2024 11:31:46
NAME	Mr. LAXMI NARAYAN NAWARIA		
Age / Sex	Male 42 Yrs 9 Mon 22 Days		
Ref. By	BANK OF BARODA	Printed On	25/08/2024 09:21:35
Lab/Hosp	Mr.MEDIWHEEL	27	

(CBC): Methodology: TLC,DLC Fluorescent Flow cytometry, HB SLS method,TRBC,PCV,PLT Hydrodynamically focused Impedance, and MCH,MCV,MCTIC,MENTZER INDEX are calculated. InstrumentName: Sysmex 6 part fully automatic analyzer XN-L,Japan



Page No: 3 of 17



 B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023

③ +91 141 4824885 ⋈ p3healthsolutionsllp@gmail.com



Patient ID	12 <b>24990</b> Patient Mob No.9166636200	Registered On	24/08/2024 11:31:46
NAME	Mr. LAXMI NARAYAN NAWARIA	Collected On	24/08/2024 12:11:44
Age / Sex	Male 42 Yrs 9 Mon 22 Days	Authorized On	25/08/2024 09:21:25
Ref. By	BANK OF BARODA	Printed On	25/08/2024 09:21:35
Lab/Hosp	Mr.MEDIWHEEL		

### **BIOCHEMISTRY**

Test Name	Value	Unit	Biological Ref Interval
FASTING BLOOD SUGAR (Plasma) Methord:- GLUCOSE OXIDASE/PEROXIDASE	99.4	mg/dl	70.0 - 115.0
Impaired glucose tolerance (IGT)	1	11 - 125 mg/dL	
Diabetes Mellitus (DM)	> 126 mg/dL		

Instrument Name: HORIBA CA60 Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm,

hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin

therapy or various liver diseases.

Technologist



- O B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ♦ +91 141 4824885 

  p3healthsolutionsllp@gmail.com



Patient ID	12 <b>24990</b> Patient Mob No.9166636200	Registered On	24/08/2024 11:31:46
NAME	Mr. LAXMI NARAYAN NAWARIA	Collected On	24/08/2024 12:11:44
Age / Sex	Male 42 Yrs 9 Mon 22 Days	Authorized On	25/08/2024 09:21:25
Ref. By	BANK OF BARODA	Printed On	25/08/2024 09:21:35

Lab/Hosp Mr.MEDIWHEEL

#### **HAEMATOLOGY**

Test Name	Value	Unit	Biological Ref Interval
GLYCOSYLATED HEMOGLOBIN (HbA	AIC)		
Methord:- CAPILLARY with EDTA	5.7	mg%	Non-Diabetic < 6.0 Good Control 6.0-7.0 Weak Control 7.0-8.0 Poor control > 8.0
MEAN PLASMA GLUCOSE Methord:- Calculated Parameter	117	mg/dL	68 - 125

#### INTERPRETATION

AS PER AMERICAN DIABETES ASSOCIATION (ADA)

Reference Group HbA1c in % Non diabetic adults >=18 years < 5.7 At risk (Prediabetes) 5.7 - 6.4 Diagnosing Diabetes >= 6.5

CLINICAL NOTES

In vitro quantitative determ nation of HbA1c in whole blood is utilized in long term monitoring of glycemia. The HbA1c level correlates with the mean glucose concentration prevailing in the course of the patient's recent history (approx - 6-8 weeks) and therefore provides much more reliable information for glycemia monitoring than do determinations of blood glucose or urinary glucose. It is recommended that the determination of HbA1c be performed at intervals of 4-6 weeks during Diabetes Mellitus therapy. Results of HbA1c should be assessed in conjunction with the patient's medical history, clinical examinations and other findings.

Some of the factors that influence HbA1c and its measurement [Adapted from Gallagher et al.]

Erythropoiesis

- Increased HbA1c; iron sitamin B12 deficiency, decreased erythropoiesis.

- Decreased HbA1c administration of erythropoietin, iron, vitamin B12, reticulocytosis, chronic liver disease 2. Altered Haemoglobin Genetic or chemical alterations in hemoglobin; hemoglobinopathies, HbF, methemoglobin, may increase or decrease HbA1c.

3. Glycation

Increased HbA1c\_alconomy, chronic renal failure, decreased intraerythrocytic pH.
 Decreased HbA1c\_cenam hemoglobinopathies, increased intra-erythrocyte pH

.4. Erythrocyte destruction

- Increased HbA1c increased erythrocyte life span: Splenectomy.

- Decreased A1c increased RBC life span: hemoglobinopathies, splenomegaly, rheumatoid arthritis or drugs such as antiretrovirals, ribavirin & dapsone.

Increased HbA1c, hyperbilingbinemia, carbamylated hemoglobin, alcoholism, large doses of aspirin, chronic opiate use chronic renal failure

- Decreased HbA1c: hyganniglyceridemia, reticulocytosis, chronic liver disease, aspirin, vitamin C and E, splenomegaly, rheumatoid arthritis or drugs

Technologist

Janu DR.TANU RUNGTA MD (Pathology) RMC No. 17226

This Report Is Not Valid For Medico Legal Purpose



- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ♦ +91 141 4824885 

  p3healthsolutionsllp@gmail.com



Patient ID	1224990	Patient Mob No.9166636200
· acionic no		

NAME Mr. Age / Sex Male

Mr. LAXMI NARAYAN NAWARIA

Age / Sex Male 42 Yrs 9 Mon 22 Days Ref. By BANK OF BARODA

Lab/Hosp Mr.MEDIWHEEL

Registered On

24/08/2024 11:31:46

Collected On Authorized On 24/08/2024 12:11:44 25/08/2024 09:21:25

Printed On

25/08/2024 09:21:35

### HAEMATOLOGY

#### HAEMATOLOGY

Test Name	Value	TI-:+	Biological Ref Interval
Test Name	value	Unit	Diological Kei Interval

BLOOD GROUP ABO Methord - Haemagghamation reaction "B" POSITIVE

Technologist



Lab/Hosp

# P3 HEALTH SOLUTIONS LLP

B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023

🕒 +91 141 4824885 🖂 p3healthsolutionsllp@gmail.com



Patient ID	1224990	Patient Mob No.9166636200	Registered On	24/08/2024 11:31:46
NAME	™r. LAXN	II NARAYAN NAWARIA	Collected On	24/08/2024 12:11:44
Age / Sex	Male	42 Yrs 9 Mon 22 Days	Authorized On	25/08/2024 09:21:25
Ref. By	BANK OF	BARODA	Printed On	25/08/2024 09:21:35

BI	OCI	MI	CT	DV

Test Name	Value	Unit	Biological Ref Interval
LIPID PROFILE			
SERUM TOTAL CHOLESTEROL Methord - CHOLESTEROL OXIDASE/PEROXIDASE	218.00	mg/dl	Desirable <200 Borderline 200-239 High> 240
InstrumentName HORIBA Interpretation: Chol disorders	esterol measurements are	used in the diagnosis and tro	eatments of lipid lipoprotein metabolism
SERUM TRIGI YCERIDES Methord - GLYCEROL PHOSPHATE OXIDASE/PREOXIDAS	121.50 SE	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500
InstrumentName Randox Rx Imola Interpretati metabolism and various endocrine disorders e.g. diab			osis and treatment of diseases involving lipid
DIRECT HDL CHOLESTEROL Methord - Direct cleatance Method	45.80	mg/dl	
			MALE- 30-70 FEMALE - 30-85

Instrument Name Rx Daytona plus Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods LDL CHOLESTEROL 151.95

Mr.MEDIWHEEL

Methord - Calculated Method

mg/dl

Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190

Interpretation: Accurate measurement of LDL-Cholesterol is of vital importance in therapies which focus on lipid reduction to prevent atheroselerosis or reduce its progress and to avoid plaque rupture.

VLDL CHOLL STEROL

24.30

mg/dl

0.00 - 80.00

Janu

Technologist,



Lab/Hosp

Mr.MEDIWHEEL

# P3 HEALTH SOLUTIONS LLP

- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ♦ +91 141 4824885 

  p3healthsolutionsllp@gmail.com



Patient ID	1224990	Patient Mob No.9166636200	Registered On	24/08/2024 11:31:46
NAME	Mr. LAXM	II NARAYAN NAWARIA	Collected On	24/08/2024 12:11:44
Age / Sex	Male	42 Yrs 9 Mon 22 Days	Authorized On	25/08/2024 09:21:25
Ref. By	BANK OF	BARODA	Printed On	25/08/2024 09:21:35

#### **BIOCHEMISTRY**

#### **BIOCHEMISTRY**

Test Name	Value	Unit	Biological Ref Interval
T.CHOLESTEROL/HDL CHOLESTEROL RATIO Methord:- Calculated	4.76		0.00 - 4.90
LDL/HDL CHOLESTEROL RATIO Methord - Calculated	3.32		0.00 - 3.50
TOTAL LIPID Methord: CALCULATED	633.80	mg/dl	400.00 - 1000.00

- 1. Measurements in the same patient can show physiological& analytical variations. Three serialsamples I week apart are recommended for Total Cholesterol. Triglycerides, HDL& LDL Cholesterol.
- 2. As per NCEP guide nes, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 20 ears with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended
- 3. Low HDL levels are associated with Coronary Heart Disease due to insufficient HDL being available to participate in reverse cholesterol transport the process by which cholesterol is eliminated from peripheral tissues.

Technologist



- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ③ +91 141 4824885 ⊠ p3healthsolutionsllp@gmail.com



Patient ID	12 <b>24990</b> Patient Mob No.9166636200	Registered On	24/08/2024 11:31:46
NAME	Mr. LAXMI NARAYAN NAWARIA	Collected On	24/08/2024 12:11:44
Age / Sex	Male 42 Yrs 9 Mon 22 Days	Authorized On	25/08/2024 09:21:25
Ref. By	BANK OF BARODA	Printed On	25/08/2024 09:21:35
Lab/Hosp	Mr.MEDIWHEEL		

### **BIOCHEMISTRY**

### **BIOCHEMISTRY**

Test Name	Value	Unit	Biological Ref Interval
LIVER PROFILE WITH GGT		-	
SERUM BILIRUBIN (TOTAL) Methord: DIAZOTIZ TO SULFANILIC	0.62	mg/dL	Infants: 0.2-8.0 mg/dL Adult - Up to - 1.2 mg/dL
SERUM BILIRUBIN (DIRECT) Methord:- DIAZOTIZED SULFANILIC	0.16	mg/dL	Up to 0.40 mg/dL
SERUM BILIRUBIN (INDIRECT) Methord - Calculated	0.46	mg/dl	0.30-0.70
SGOT Methord - ITCC	29.5	U/L	0.0 - 40.0
SGPT Methord - IFCC	22.6	U/L	0.0 - 40.0
SERUM ALKALINE PHOSPHATASE Methord - DGKC - SCT	70.00	U/L	53.00 - 141.00
SERUM GAMMA GT Methord: Nzasz methodology Instrument Name Ran for its Impla Interpretation: Elevations in GGT levels are seen earlier and more pronounce	17.00	U/L zymes in cases of obstructive jaundice and	10.00 - 45.00
metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or phepatic bilians obstruction. Only moderate elevations in the enzyme level (2)		with infectious hepatitis.	
SERUM TOTAL PROTEIN Methord: BIURET	6.73	g/dl	6.00 - 8.40
SERUM ALBUMIN Methord:- BROMOCKI SOL GREEN	4.84	g/dl	3.50 - 5.50
SERUM GLOBULIN Methorit: CALCULATION	1.89 L	gm/dl	2.20 - 3.50
A/G RATIO	2.56 H	s.	1.30 - 2.50

Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

Note: These are group of tests that can be used to detect the presence of liver disease, distinguish among different types of liver disorders, gauge the extent of known liver damage, and monitor the response to treatment. Most liver diseases cause only mild symptoms initially, but these diseases must be detected early. Some tests are associated with functionality (e.g.,

Trehnologist



- O B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ♦ +91 141 4824885 

  p3healthsolutionsllp@gmail.com

42 Yrs 9 Mon 22 Days



1224990 Patient ID Patient Mob No.9166636200

NAME

Mr. LAXMI NARAYAN NAWARIA

Age / Sex Ref. By

BANK OF BARODA

Lab/Hosp Mr MEDIWHEEL Registered On

24/08/2024 11:31:46

Collected On Authorized On 24/08/2024 12:11:44

25/08/2024 09:21:25

Printed On

25/08/2024 09:21:35

#### **BIOCHEMISTRY**

### BIOCHEMISTRY

Test Name Value **Biological Ref Interval** 

albumin), some with collular integrity (e.g., transaminase), and some with conditions linked to the biliary tract (gamma-glutamyl transferase and alkaline phosphatase). Conditions with elevated levels of ALT and AST include hepatitis A,B,C ,paracetamol toxicity etc Several biochemical tests are useful in the evaluation and management of patients with hepatic dysfunction. Some or all of these measurements are also carried out (usually about twice a year for routine cases) on those individuals taking certain medications, such as anticonvulsants, to ensure that the medications are not adversely impacting the person's liver



Technologist,

DR.TANU RUNGTA MD (Pathology) RMC No. 17226

This Report Is Not Valid For Medico Legal Purpose



 B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023

🕲 +91 141 4824885 ⊠ p3healthsolutionsllp@gmail.com



Patient ID	12 <b>24990</b> Patient Mob No.9166636200	Registered On	24/08/2024 11:31:46
NAME	Mr. LAXMI NARAYAN NAWARIA	Collected On	24/08/2024 12:11:44
Age / Sex	Male 42 Yrs 9 Mon 22 Days	Authorized On	25/08/2024 09:21:25
Ref. By	BANK OF BARODA	Printed On	25/08/2024 09:21:35
Lab/Hosp	Mr.MEDIWHEEL		

#### BIOCHEMISTRY

BIOCHEMISTRY					
BIOCHEMISTRY					
Test Name	Value	Unit	Biological Ref Interval		
RET / KET WITH ELECTROLYTES					
SERUM UREA Methord:-UREASE GLUTAMATE DEHYDROGENASE	23.90	mg/dl	10.00 - 50.00		
InstrumentName: HORIBA CA 60 Interpretation : diseases.	Urea measurements a	ire used in the diagnosis and	treatment of certain renal and metabolic		
SERUM CREATININE Methord: JAFFI	0.69	mg/dl	Males : 0.6-1.50 mg/dl Females : 0.6 -1.40 mg/dl		
Interpretation: Creatinine is measured primarily to assess kidney function relatively independent of protein ingestion, water intake					
clinically significant. SERUM URIC ACID Methord: URICANE/PEROXIDASE	6.60	mg/dl	2.40 - 7.00		
InstrumentName.HORIBA YUMIZEN CA60 Daytor Polycythaemia vera, Malignancies,Hypothyroidism,Rare					
SODIUM Methord:- Jon-Selective Electrode with Serum	139.7	mmol/L	135.0 - 145.0		
POTASSIUM Methord: Ion-Selective Electrode with Serum	3.44 L	mmol/L	3.50 - 5.00		
CHLORIDE Methord: Jon-Selective Electrode with Serum	103.3	mmol/L	97.0 - 107.0		
SERUM CALCIUM Medoord: Arsenaze III Method	10.00	mg/dL	8.80 - 10.20		
InstrumentName:MISPA PLUS Interpretation: So Increases in serum PTH or vitamin D are usually asso nephrosis and pancreatitis.	erum calcium levels a peiated with hypercal	are believed to be controlled deemia .Hypocalcemia may	by parathyroid hormone and vitamin D. be observed in hypoparathyroidism,		
SERUM TOTAL PROTEIN	6.73	g/dl ·	6.00 - 8.40		
SUM ALIJUMIN M. rd-Bromockesol green	4.84	g/dl	3.50 - 5.50		

Jechnologist,



- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ♦ +91 141 4824885 
  p3healthsolutionsllp@gmail.com



Patient ID	1224990 Patient Mob No.9166636200	Registered On	24/08/2024 11:31:46
NAME	Mr. LAXMI NARAYAN NAWARIA	Collected On	24/08/2024 12:11:44
Age / Sex	Male 42 Yrs 9 Mon 22 Days	Authorized On	25/08/2024 09:21:25
Ref. By	BANK OF BARODA	Printed On	25/08/2024 09:21:35
Lab/Hosp	Mr.MEDIWHEEL		

#### **BIOCHEMISTRY**

#### BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
SF 79 UM GLOBULIN Methods - CALCULATION	1.89 L	gm/dl	2.20 - 3.50
A/G RATIO	2.56 H		1.30 - 2.50

Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

#### INTERPRETATION

Killing function tests are group of tests that can be used to evaluate how well the kidneys are functioning. Creatinine is a waste product that comes from protein in the diet and also comes from the normal wear and tear of muscles of the body. In blood, it is a marker of GFR in urine, it can remove the need for 24-hourcollections for many analytes or be used as a quality assurance tool to assess the accuracy of a 24-hour collection Higher levels may be a sign that the kidneys are not working properly. As kidney disease progresses, the level of creatinine and research the blood accesses. Certain drugs are nephrotoxic hence KFT is done before and after initiation of treatment with these drugs.

Low serum creatmine values are rare, they almost always reflect low muscle mass

Apart from renal failure Blood Urea can increase in dehydration and GI bleed

Technologist-



- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ♦ +91 141 4824885 ⊠ p3healthsolutionsllp@gmail.com



Patient ID	12 <b>24990</b> Patient Mob No.9166636200	Registered On	24/08/2024 11:31:46
NA**E	Mr. LAXMI NARAYAN NAWARIA	Collected On	24/08/2024 12:11:44
Age / Sex	Male 42 Yrs 9 Mon 22 Days	Authorized On	25/08/2024 09:21:25
Ref. By	BANK OF BARODA	Printed On	25/08/2024 09:21:35
Lah/Hosp	Mr MEDIMHEEL		

### **CLINICAL PATHOLOGY**

### **CLINICAL PATHOLOGY**

Tort Name	Value	Unit	Biological Ref Interval
UP'NE SUGAR (FASTING) Collected Sample Received	Nil		Nil
URINE SUGAR PP Collected Sample Received	Nil		Nil

Technologist7



Lab/Hosp

# P3 HEALTH SOLUTIONS LLP

- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- 🕲 +91 141 4824885 🖂 p3healthsolutionsllp@gmail.com



Patient ID	12 <b>24990</b> Patient Mob No.9166636200	Registered On	24/08/2024 11:31:46
NAME	Mr. LAXMI NARAYAN NAWARIA	Collected On	24/08/2024 12:11:44
Age / Sex	Male 42 Yrs 9 Mon 22 Days	Authorized On	25/08/2024 09:21:25
Ref. By	BANK OF BARODA	Printed On	25/08/2024 09:21:35

#### **IMMUNOASSAY**

Test Name	Value	Unit	Biological Ref Interval
PSA (PROSTATE SPECIFIC ANTIGEN) -TOTAL Methord: -Methodology CLIA	2.190	ng/mL	0.00-4.00

CLINICAL NOTES - Prostate-specific antigen (PSA)is a 34-kD glycoprotein produced almost exclusively by the prostate gland.

PSA is normally present in the blood at very low levels. Increased levels of PSA may suggest the presence of prostate cancer.

- 1.Immediate PSA testing following digital rectal examination, ejaculation, prostatic massage, indwelling catheterization, ultrasonography and needle biopsy of prostate is not
- recommended as they falsely elevate levels

Mr.MEDIWHEEL

- 2. PSA values regardless of levels should not be interpreted as absolute evidence of the presence or absence of disease. All values should be correlated with clinical
- findings and other investigations
- 3. Physiological decrease in PSA level by 18% has been observed in sedentary patients either due to supine position or suspended sexual activity

#### Clinical Use

- An aid in the early detection of Prostate cancer when used in conjunction with Digital rectal examination in males more than 50 years of age and in those with two or more
- affected first degree relatives.
- · Follow up and management of Prostate cancer patients
- . Detect metastatic or persistent disease in patients following surgical or medical treatment of Prostate cancer

#### NOTE

PSA levels can be also increased by prostatitis, irritation, benign prostatic hyperplasia (BPH), and recent ejaculation, producing a false positive result. Digital rectal examination (DRE) has been shown in several studies to produce an increase in PSA. However, the effect is clinically insignificant, since DRE causes the most substantial increases in patients with PSA levels already elevated over 4.0 ng/mL.

Ohee'ty has been reported to reduce serum PSA levels. Delayed early detection may partially explain worse outcomes in obese men with early provide cancer. Aftertreatment, higher BMI also correlates to higher risk of recurrence.

Technologist,



- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- 🕲 +91 141 4824885 🖂 p3healthsolutionsllp@gmail.com



Patient ID	1224990 Patient Mob No.9166636200	Registered On	24/08/2024 11:31:46
NAME	Mr. LAXMI NARAYAN NAWARIA	Collected On	24/08/2024 12:11:44
Age / Sex	Male 42 Yrs 9 Mon 22 Days	Authorized On	25/08/2024 09:21:25
Ref. By	BANK OF BARODA	Printed On	25/08/2024 09:21:35
Lab/Hosp	Mr.MEDIWHEEL		

#### **IMMUNOASSAY**

#### **IMMUNOASSAY**

Tref Name	Value	Unit	Biological Ref Interval
TOTAL THYROID PROFILE		40	
THYROID-TRIIODOTHYRONINE T3 Methord: Chemiluminescence	1.38	ng/ml	0.69 - 2.15
TUVROID - THYROXINE (T4) Methord - Chemiluminescence	5.89	ug/dl	5.20 - 12.70
TSH Methord - Chemiluminescence	4.470	μIU/mL	0.470 - 4.680

#### Note:

- 1. TSH levels are subject to circadian variation, reaching peak levels between 2 4.a.m. and at a minimum between 6-10 pm. The variation is of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 2. Recommended test for T3 and T4 is unbound fraction or free levels as it is metabolically active.
- 3. Physiological rise in Total T3 / T4 levels is seen in pregnancy and in patients on steroid therapy. Clinical Uso
- in infancy and early childhood

\*\*\* End of Report \*\*\*

\*\*\* End of Report \*\*\*

Technologist 7



- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ③ +91 141 4824885 ⊠ p3healthsolutionsllp@gmail.com



Pa	tieni	

1224990 Patient Mob No.9166636200

Mr. LAXMI NARAYAN NAWARIA

NAME Age / Sex

42 Yrs 9 Mon 22 Days

Ref. By

BANK OF BARODA

Lab/Hosp

Mr.MEDIWHEEL

Registered On

Collected On Authorized On

Printed On

24/08/2024 11:31:46

24/08/2024 12:11:44

25/08/2024 09:21:25

25/08/2024 09:21:35

### **CLINICAL PATHOLOGY**

Value	Unit	Biological Ref Interval
	. No	
PALE YELLO	OW	PALE YELLOW
		Clear
, <del>- 11 - 2 - 41 -</del>		
7.0		5.0 - 7.5
		1.010 - 1.030
NIL		NIL
NIL		NIL
NEGATIVE		NEGATIVE
NORMAL		NORMAL
NEGATIVE		NEGATIVE
NEGATIVE		NEGATIVE
NIL	/HPF	NIL
2-3	/HPF	2-3
0-2	/HPF	2-3
ABSENT		ABSENT
ABSENT		
	PALE YELLA Clear  7.0 1.020 NIL NIL NEGATIVE NORMAL NEGATIVE NEGATIVE NEGATIVE ABSENT ABSENT ABSENT ABSENT ABSENT ABSENT	PALE YELLOW Clear  7.0 1.020 NIL NIL NEGATIVE NORMAL NEGATIVE NEGATIVE NEGATIVE NEGATIVE NEGATIVE ABSENT ABSENT ABSENT ABSENT ABSENT ABSENT ABSENT

Technologist7



- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- 🕒 +91 141 4824885 🖂 p3healthsolutionsllp@gmail.com



NAME:	MR. LAXMI NARAYAN NAWARIA	AGE	42 YRS/M
REF.BY	BANK OF BARODA	DATE	24/08/2024

### **CHEST X RAY (PA VIEW)**

Bilateral lung fields appear clear.

Bilateral costo-phrenic angles appear clear.

Cardiothoracic ratio is normal.


Thoracic soft tissue and skeletal system appear unremarkable.

Soft tissue shadows appear normal.

IMPRESSION: No significant abnormality is detected

Som

DR. ROHAN GAUR
M.B.B.S, M.D (Radiodiagnosis)
RMC no. 17887





- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ③ +91 141 4824885 ⊠ maxcarediagnostics1@gmail.com



MR. LAXMI NARAYAN NAWARIA	42 Y/M
Registration Date: 24/08/2024	Ref. by: BANK OF BARODA

### 2D-ECHOCARDIOGRAPHY M.MODE WITH DOPPLER STUDY: FAIR TRANSTHORACIC ECHOCARIDIOGRAPHIC WINDOW MORPHOLOGY:

MITRAL VALVE		NO	NORMAL		TRI	TRICUSPID VALVE			AL .	
<b>AORTIC VALVE</b>		NO	NORMAL		PUI	PULMONARY VALVE			NORMAL	
				M.MOD	E EXAMITAT	TION:				
AO	2.6	Cm	LA		2.7	cm	IVS-D	0.8	cm	
IVS-S	1.2	cm	LVII	D	4.2	cm	LVSD	2.7	cm	
LVPW-D	0.8	cm	LVP	W-S	1.2	cm	RV		cm	
RVWT		cm	EDV	1		MI	LVVS		ml	
LVEF	55-60%	ó	A		RWM	Α	ABSENT			
			4	<u>c</u>	HAMBERS:		No.			
LA	NOR	MAL	59	RA			NORMAL			
LV	NOR	MAL	47	RV		(27) A	NORMAL			
PERICARDIUM		F.S.		NORMA		NOT CHEF AN				
		25		COLO	OUR DOPPLE	R:				
		MITRA	L VALVE							
E VELOCITY		0.97	m/se	c PEA	K GRADIENT	RADIENT		Mm/hg		
A VELOCITY		0.68	m/se	c MEA	N GRADIEN	RADIENT		Mm/hg		
MVA BY PHT		100	Cm2	Cm2 MVA BY		PLANIMETRY		Cm2	Cm2	
MITRAL REGUE	RGITATION	18				ABSENT	1			
		AORTIC	VALVE				All Control			
PEAK VELOCITY	Y	1.47	The state of	m/sec	PEAK G	RADIENT	4	mm	/hg	
AR VMAX				m/sec	MEAN	MEAN GRADIENT		mm	mm/hg	
<b>AORTIC REGUR</b>	RGITATION		- 3		ABSENT					
		TRICUSE	PID VAL	<b>VE</b>	Series -					
PEAK VELOCITY	Υ			m/sec	PEAK G	PEAK GRADIENT			nm/hg	
MEAN VELOCIT	TY			m/sec MEAN GRADIENT			r	nm/hg		
VMax VELOCI	TY									
				1	ABSEN	т				
TRICUSPID REG	URGITATIO		DALA DV.	/413/5	ABSEN					
		PULMO	DNARY \		Malana	PEAK GRADI	ENT		Mm/h	
PEAK VELOCIT			1.04		M/sec.	MEAN GRAD			Mm/h	
MEAN VALOCI PULMONARY						ABSENT	VICIAL .		17111/11	

#### Impression—

- NORMAL LV SIZE & CONTRACTILITY.
- NO RWMA, LVEF 55-60%.
- NORMAL DIASTOLIC FUNCTION.
- ALL CARDIAC VALVES ARE NORMAL.
- NO CLOT, NO VEGETATION, NO PERICARDIAL EFFUSION.

(Cardiologist)







- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- 🕲 +91 141 4824885 🖂 maxcarediagnostics1@gmail.com



MR. LAXMI NARAYAN NAWARIA	42 Y/M		
Registration Date: 24/08/2024	Ref. by: BANK OF BARODA		

### **ULTRASOUND OF WHOLE ABDOMEN**

**Liver** is of normal size (13.0 cm). Echo-texture is normal. No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

**Gall bladder** is well distended. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

**Spleen** is of normal size and shape. Echotexture is normal. No focal lesion is seen.

**Kidneys** are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. Collecting system does not show any calculus or dilatation.

Right kidney is measuring approx. 9.1 x 3.9 cm.

Left kidney is measuring approx. 9.3 x 4.3 cm.

Urinary bladder is well distended and does not show any calculus or mass lesion.

Prostate is normal in size (17.0 cc) with normal echotexture and outline.

No enlarged nodes are visualized. No retro-peritoneal lesion is identified. No significant free fluid is seen in pelvis.

### IMPRESSION:-

No significant abnormality is detected.

DR. ROHAN GAUR

M.B.B.S, M.D (Radiodiagnosis)

RMC no. 17887

