# Dr. Goyals Path Lab & Imaging Centre B-51, Ganesh Nagar, Near Metro Piller No. General Physical Examination Sondala Jainur-302019

Name: Devended Kumout Dharbwal. Age: 47 Sex:	95
DOB: 94-08-19-16  Referred By: BOR (Mediwheel)	
Referred By:BOB (Medi Wheel)	
Photo ID: ID #:	_
FIIULU ID. 14 (/ 5)	
Ht: 172 (cm) Wt: 82 (Kg)	
Chest (Expiration):(cm) Abdomen Circumference:	(
Blood Pressure: 108/59 mm Hg PR: 83/min	
BMI 27 7	
Eye Examination: Dis. Nisson 6/6 6/24 . Near vision.  Bl eyes (with spees). Mossocal Color vision  Other: Not significant	
Bleges (with spees). Mossocial Color 112100	
Other: Not significant	
•	
On examination he/she appears physically and mentally fit: Yes / No	
Signature Of Examine: Name of Examinee:	
Signature Of Examine: Name of Examinee: Name of	
Signature Medical Examiner:	



Tiles 1000

Dr. R.B. R. R. D. O. T. O. S. O. S. C. R. S. O. S. O. T. O. S. O.

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Near Metro Piller No. General Physical Examination
Sodala, Jaipur-302019

Tele: 0141-2293346, 4049787 8887049787

Name: Devendry Kymon, Dhourwal, Age: 47 Sex: Mg/c
DOB: 24-08-19-16
Referred By:
Photo ID: ID#:
Ht:(cm) Wt:(Kg)
Chest (Expiration):(cm)
Blood Pressure: 108 69 mm Hg PR: 83/min
BMI 27 T
Eye Examination: Dis. Vision 6/6 6/24. Near vision, M/6 Bl eyes (with spees). Mossocal Color vision
Bleges (with spees). Normal Color vision
Other: Lod significant
On examination he/she appears physically and mentally fit: Yes / No
Signature Of Examine:  Name of Examinee:  Name of Examinee:  Name Medical Examiner:
Signature Medical Examiner: Name Medical Examiner



Tille

Dr. R. B. B. B. No. O. To S. S.



B-51, Ganesh Nagar, Near Metro Pillar No. 109-110, New Sanganer Road, Jaipur

Tele: 0141-2293346, 4049787, 9887049787



Date

:- 23/03/2024 10:26:13

NAME :- Mr. DEVENDER KUMAR

Sex / Age :- Male

47 Yrs 6 Mon 30 Days

Company :- MediWheel

Patient ID :-12236564 Ref. By Doctor:-BOB

Lab/Hosp:-

Final Authentication: 23/03/2024 18:52:58

**BOB PACKAGE ABOVE 40MALE** 

#### X RAY CHEST PA VIEW:

Both lung fields appears clear.

Bronchovascular markings appear normal.

Trachea is in midline.

Both the hilar shadows are normal.

Both the C.P.angles is clear.

Both the domes of diaphragm are normally placed.

Bony cage and soft tissue shadows are normal.

Heart shadows appear normal.

Impression: - Normal Study

(Please correlate clinically and with relevant further investigations)

DR. POORVI MALIK MBBS, MD, DNB (RADIO DIAGNOSIS) RMC REG. NO. 21505

End of Report \*

Dr. Piyush Goyal ( D.M.R.D.) **BILAL** 

Transcript by.

M.B.B.S., D.M.R.D.

Page No: 1 of 1

Dr. Ashish Choudhary MBBS, MD (Radio Diagnosis) Fetal Medicine Consultant

Dr. Abhishek Jain

RMC No. 21687

Dr. Navneet Agarwal RMC No. 33613/14911

Dr. Poorvi Malik MBBS, DNB, (Radio-Diagnosis) MD, DNB (Radio Diagnosis) MBBS, MD, DNB (Radio Diagnosis) RMC No. 21505

FMF ID - 260517 | RMC No 22430 his report is not valid for medico-legal purpose



B-51. Ganesh Nagar. Near Metro Pillar No. 109-110, New Sanganer Road, Jaipur Tele: 0141-2293346, 4049787, 9887049787



Date :- 23/03/2024 10:26:13

NAME :- Mr. DEVENDER KUMAR

Sex / Age :- Male

47 Yrs 6 Mon 30 Days

Company :-MediWheel Patient ID: -12236564 Ref. By Doctor:-BOB

Lab/Hosp:-

Final Authentication: 23/03/2024 12:20:10

**BOB PACKAGE ABOVE 40MALE** 

#### **USG WHOLE ABDOMEN**

Liver is borderline enlarged in size (~14.9cm) and shows mildly raised parenchymal echogenicity. No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is of normal size. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within

Spleen is of normal size and shape. Echotexture is normal. No focal lesion is seen.

Kidneys are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. No focal lesion is seen. Collecting system does not show any dilatation or calculus.

Urinary bladder is well distended and showing smooth wall with normal thickness. Urinary bladder does not show any calculus or mass lesion.

Prostate is normal in size with normal echo-texture and outline. No significant free fluid is seen in peritoneal cavity.

#### **IMPRESSION:**

\* Borderline hepatomegaly with grade I fatty changes.

Needs clinical correlation.

End of Report \*\*\*

BILAL

Page No: 1 of 1

Dr. Poorvi Malik

Transcript by.



B-51, Ganesh Nagar, Near Metro Pillar No. 109-110, New Sanganer Road, Jaipur

Tele: 0141-2293346, 4049787, 9887049787



:- 23/03/2024 10:26:13

NAME :- Mr. DEVENDER KUMAR Sex / Age :- Male

47 Yrs 6 Mon 30 Days

Company :- MediWheel

Patient ID: -12236564 Ref. By Doctor:-BOB

Lab/Hosp :-

Final Authentication: 23/03/2024 12:54:37

**BOB PACKAGE ABOVE 40MALE** 2D ECHO OPTION TMT (ADULT/CHILD)

#### **2D-ECHOCARDIOGRAPHY M.MODE WITH DOPPLER STUDY:**

\_FAIR TRANSTHORACIC ECHOCARIDIOGRAPHIC WINDOW MORPHOLOGY:

MITRAL VALVE	NORMAL		TRICUS	TRICUSPID VALVE				
AORTIC VALVE	NORMAL		PULMO	ONARY VALVE	NORMAL			
	N	1.MODE E	XAMITATION:					
AO	27	mm	LA	33	Mm	IVS-D	9	mm
IVS-S	18	mm	LVID	39	Mm	LVSD	24	mm
LVPW-D	10	mm	LVPW-S	16	Mm	RV		mm
RVWT		mm	EDV		MI	LVVS		ml
LVEF	67%			RWMA		ABSENT		
				CHA	AMBERS:			
LA	NORMAL		RA			NORMAL		

		CHAMBERS					
LA	NORMAL	RA	NORMAL				
LV	NORMAL	RV	NORMAL				
PERICARDII	UM	NORMAL					

#### **COLOUR DOPPLER:**

		MITRA	L VAL	/E						
E VELOCITY	0.79		m/sec	P	EAK G	RADIENT			Mm/hg	
A VELOCITY	0.58		m/sec	N	IEAN (	GRADIEN	r		Mm/hg	
MVA BY PHT			Cm2	N	IVA BY	PLANIM	ETRY		Cm2	
MITRAL REGURGITATIO	N						ABSENT			
	-	AORTI	C VAL	/E					1.	
PEAK VELOCITY	1.1		m	/sec		PEAK GR	ADIENT		mm	/hg
AR VMAX			m	/sec		MEAN GRADIENT mm/hg				/hg
AORTIC REGURGITATIO	N					ABSENT				
	TF	RICUSI	PID VA	LVE						
PEAK VELOCITY	0.	.48		m/s	ec	PEAK G	RADIENT		r	nm/hg
MEAN VELOCITY				m/s	ec	MEAN (	GRADIENT		r	nm/hg
VMax VELOCITY										
TRICUSPID REGURGITA						ABSENT				
	F	PULMO	ONARY	VAL						
PEAK VELOCITY					N	√/sec.	PEAK GRADIEN	Г		Mm/hg
MEAN VALOCITY							MEAN GRADIEN	IT		Mm/hg
PULMONARY REGURGI	TATION						ABSENT		•	

Page No: 1 of 2

**AHSAN** 

Transcript by.

Dr. Piyush Goyal M.B.B.S., D.M.R.D. RMC Reg No. 017996

Dr. Ashish Choudhary MBBS, MD (Radio Diagnosis) Fetal Medicine Consultant

FMF ID - 260517 | RMC No 22430

Dr. Abhishek Jain

RMC No. 21687

Dr. Navneet Agarwal RMC No. 33613/14911

Dr. Poorvi Malik MBBS, DNB, (Radio-Diagnosis) MD, DNB (Radio Diagnosis) MBBS, MD, DNB (Radio Diagnosis) RMC No. 21505



B-51, Ganesh Nagar, Near Metro Pillar No. 109-110, New Sanganer Road, Jaipur

Tele: 0141-2293346, 4049787, 9887049787



:- 23/03/2024 10:26:13

NAME :- Mr. DEVENDER KUMAR

Sex / Age :- Male

47 Yrs 6 Mon 30 Days

Company :- MediWheel

Patient ID: -12236564 Ref. By Doctor:-BOB

Lab/Hosp:-

Final Authentication: 23/03/2024 12:54:37

#### Impression--

- 1. Normal LV size & contractility
- LVEF 67 %. 2. No RWMA,
- 3. Normal cardiac chamber.
- 4. Normal valve
- 5. No clot, no vegetation, no pericardial effusion.

(Cardiologist)

\*\*\* End of Report \*\*\*

Page No: 2 of 2

**AHSAN** 

Transcript by.



B-51, Ganesh Nagar, Near Metro Piller No. 109-110, New Sanganer Road,

Sodala, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www. drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com | Date :- 23/03/2024 10:26:13

NAME :- Mr. DEVENDER KUMAR

47 Yrs 6 Mon 30 Days Sex / Age :- Male

Company :-MediWheel

Sample Type :- EDTA

Sample Collected Time 23/03/2024 10:39:42

Final Authentication: 23/03/2024 13:12:49

HAEMATOLOGY

**Test Name** Value Unit **Biological Ref Interval** 

Lab/Hosp:-

**BOB PACKAGE ABOVE 40MALE** 

GLYCOSYLATED HEMOGLOBIN (HbA1C)

5.9

Patient ID: -12236564

Ref. By Dr:- BOB

Non-diabetic: < 5.7 Pre-diabetics: 5.7-6.4 Diabetics: = 6.5 or higher ADA Target: 7.0

Action suggested: > 6.5

Instrument name: ARKRAY's ADAMS Lite HA 8380V, JAPAN.

Test Interpretation:

HbA1C is formed by the condensation of glucose with n-terminal valine residue of each beta chain of HbA to form an unstable schiff base. It is the major fraction, constituting approximately 80% of HbA1c. Formation of glycated hemoglobin (GHb) is essentially irreversible and the concentration in the blood depends on both the lifespan of the red blood cells (RBC) (120 days) and the blood glucose concentration. The GHb concentration represents the integrated values for glucose overthe period of 6 to 8 weeks. GHb values are free of day to day glucose fluctuations and are unaffected by recent exercise or food ingestion. Concentration of plasmaglucose concentration in GHb depends on the time interval, with more recent values providing a larger contribution than earlier values. The interpretation of GHbdepends on RBC having a normal life span. Patients with hemolytic disease or other conditions with shortened RBC survival exhibit a substantial reduction of GHb.High GHb have been reported in iron deficiency anemia. GHb has been firmly established as an index of long term blood glucose concentrations and as a measureof the risk for the development of complications in patients with diabetes mellitus. The absolute risk of retinopathy and nephropathy are directly proportional to themean of HbA1C.Genetic variants (e.g. HbS trait, HbC trait), elevated HbF and chemically modified derivatives of hemoglobin can affect the accuracy of HbA1cmeasurements. The effects vary depending on the specific Hb vatiant or derivative and the specific HbA1c method.

Ref by ADA 2020

MEAN PLASMA GLUCOSE

Method:- Calculated Paramete

123

mg/dL

Non Diabetic < 100 mg/dL Prediabetic 100- 125 mg/dL Diabetic 126 mg/dL or Higher

**AJAYSINGH Technologist** 

Page No: 1 of 13







B-51, Ganesh Nagar, Near Metro Piller No. 109-110, New Sanganer Road,

Sodala, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www. drgoyalspathleb.com | E-mail: drgoyalpiyush@gmail.com | Date :- 23/03/2024 10:26:13

NAME :- Mr. DEVENDER KUMAR

Sex / Age :- Male 47 Yrs 6 Mon 30 Days

Company :- MediWheel

Sample Type :- EDTA

Patient ID :-12236564

Ref. By Dr:- BOB

Lab/Hosp :-

Final Authentication: 23/03/2024 13:12:49

HAEMATOLOGY

Sample Collected Time 23/03/2024 10:39:42

HAEMATOLOGY						
Value	Unit	Biological Ref Interval				
13.7	g/dL	13.0 - 17.0				
5.33	/cumm	4.00 - 10.00				
52.8	%	40.0 - 80.0				
41.7 H	%	20.0 - 40.0				
1.8	%	1.0 - 6.0				
3.5	%	2.0 - 10.0				
0.2	%	0.0 - 2.0				
2.82	10^3/uL	1.50 - 7.00				
2.23	10^3/uL	1.00 - 3.70				
0.09	10^3/uL	0.00 - 0.40				
0.18	10^3/uL	0.00 - 0.70				
0.01	10^3/uL	0.00 - 0.10				
4.90	x10^6/uL	4.50 - 5.50				
43.40	%	40.00 - 50.00				
88.5	fL	83.0 - 101.0				
27.9	pg .	27.0 - 32.0				
31.6	g/dL	31.5 - 34.5				
195	x10^3/uL	150 - 410				
14.2 H	%	11.6 - 14.0				
18.06						
	13.7 5.33 52.8 41.7 H 1.8 3.5 0.2 2.82 2.23 0.09 0.18 0.01 4.90 43.40 88.5 27.9 31.6 195 14.2 H	Value         Unit           13.7         g/dL           5.33         /cumm           52.8         %           41.7 H         %           1.8         %           3.5         %           0.2         %           2.82         10^3/uL           0.09         10^3/uL           0.18         10^3/uL           0.01         10^3/uL           4.90         x10^6/uL           43.40         %           88.5         fL           27.9         pg           31.6         g/dL           195         x10^3/uL           14.2 H         %				

The Mentzer index is used to differentiate iron deficiency anemia from beta thalassemia trait. If a CBC indicates microcytic anemia, these are two of the most likely causes, making it necessary to distinguish between them.

If the quotient of the mean corpuscular volume divided by the red blood cell count is less than 13, thalassemia is more likely. If the result is greater than 13, then iron-deficiency anemia is more likely.

AJAYSINGH **Technologist** 

Page No: 2 of 13



#### Path Lab & Imaging Centre

.B-51, Ganesh Nagar, Near Metro Piller No. 109-110, New Sanganer Road,

Sodala, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www. drgoyalspathlab.com | E-majl, drgoyalpiyush@gmall.com Date - 23/03/2024 10:26:13

Patient ID :-12236564

NAME :- Mr. DEVENDER KUMAR

Ref. By Dr:- BOB

Lab/Hosp :-

Company :- MediWheel

Sex / Age :- Male

Sample Type :- EDTA

Sample Collected Time 23/03/2024 10:39:42

Final Authentication: 23/03/2024 13:12:49

**HAEMATOLOGY** 

Test Name Value Unit Biological Ref Interval

Erythrocyte Sedimentation Rate (ESR)

07

mm/hr.

00 - 13

(ESR) Methodology: Measurment of ESR by cells aggregation.

47 Yrs 6 Mon 30 Days

Instrument Name : Indepedent form Hematocrit value by Automated Analyzer (Roller-20)

Interpretation : ESR test is a non-specific indicator of inflammatory disease and abnormal protein states.

The test in used to detect, follow course of a certain disease (e.g-tuberculosis, rheumatic fever, myocardial infarction

Levels are higher in pregnency due to hyperfibrinogenaemia.

The "3-figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia (CBC) in the the delegated by the control of the control o

AJAYSINGH Technologist

Page No: 3 of 13







MC- 5509

B-51, Ganesh Nagar, Near Metro Piller No. 109-110, New Sanganer Road,

Sodala, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www. drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com | Date : -23/03/2024 10:26:13

NAME :- Mr. DEVENDER KUMAR

47 Yrs 6 Mon 30 Days Sex / Age :- Male

Company :- MediWheel Sample Type :- PLAIN/SERUM Patient ID: -12236564

Ref. By Dr:- BOB

Lab/Hosp:-

Final Authentication: 24/03/2024 12:11:13

#### **BIOCHEMISTRY**

Sample Collected Time 23/03/2024 10:39:42

Test Name	Value	Unit	Biological Ref Interval
LIPID PROFILE			
TOTAL CHOLESTEROL Method:- Enzymatic Endpoint Method	189.37	mg/dl	Desirable <200 Borderline 200-239 High> 240
TRIGLYCERIDES Method:- GPO-PAP	166.57 H	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500
DIRECT HDL CHOLESTEROL Method:- Direct clearance Method	35.83	mg/dl	Low < 40 High > 60
DIRECT LDL CHOLESTEROL Method:- Direct clearance Method	125.78	mg/dl .	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190
VLDL CHOLESTEROL Method:- Calculated	33.31	mg/dl	0.00 - 80.00
T.CHOLESTEROL/HDL CHOLESTEROL RATIO Method:- Calculated	<b>5.29</b> H		0.00 - 4.90
LDL / HDL CHOLESTEROL RATIO Method:- Calculated	3.51 H		0.00 - 3.50
TOTAL LIPID  Method:- CALCULATED  TOTAL CHOLESTEROL InstrumentName: Randox Rx Imola	613.88	mg/dl	400.00 - 1000.00  diagnosis and treatments of lipid lipoprotein metabolism

disorders

TRIGLYCERIDES InstrumentName: Randox Rx Imola Interpretation: Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction

DIRECT HDLCHOLESTERO InstrumentName:Randox Rx Imola Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.

 $\textbf{DIRECT LDL-CHOLESTEROL InstrumentName}: Randox\ Rx\ Imola\ \ \textbf{Interpretation}: Accurate\ measurement\ of\ LDL-Cholesterol\ is\ of\ vital\ importance\ in\ therapies\ which\ focus\ on\ lipid\ and\ on\ lipid\ on\ lipid$ reduction to prevent atherosclerosis or reduce its progress and to avoid plaque rupture

TOTAL LIPID AND VLDL ARE CALCULATED

SURENDRAKHANGA

Page No: 4 of 13







B-51, Ganesh Nagar, Near Metro Piller No. 109-110, New Sanganer Road,

Sodala, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www. drgoyalspathlab.com | E-mail: drgoyalplyush@gmall.com | Date 23/03/2024 10:26:13

NAME :- Mr. DEVENDER KUMAR

47 Yrs 6 Mon 30 Days Sex / Age :- Male

Company:- MediWheel

Sample Type :- PLAIN/SERUM

Patient ID :-12236564

Ref. By Dr:- BOB

Lab/Hosp :-

Final Authentication: 24/03/2024 12:11:13

#### BIOCHEMISTRY

Sample Collected Time 23/03/2024 10:39:42

DIOCHEMIST KT								
Test Name	Value	Unit	Biological Ref Interval					
LIVER PROFILE WITH GGT								
SERUM BILIRUBIN (TOTAL) Method:- Colorimetric method	0.57	mg/dl	Up to - 1.0 Cord blood <2 Premature < 6 days <16 Full-term < 6 days= 12 1month - <12 months <2 1-19 years <1.5 Adult Up to - 1.2 Ref-(ACCP 2020)					
SERUM BILIRUBIN (DIRECT) Method:- Colorimetric Method	0.13	mg/dL	Adult - Up to 0.25 Newborn - <0.6 >- 1 month - <0.2					
SERUM BILIRUBIN (INDIRECT) Method:- Calculated	0.44	. mg/dl	0.30-0.70					
SGOT Method:- IFCC	26.7	U/L	Men- Up to - 37.0 Women - Up to - 31.0					
SGPT Method:- IFCC	30.1	U/L	Men- Up to - 40.0 Women - Up to - 31.0					
SERUM ALKALINE PHOSPHATASE Method:- AMP Buffer	57.60	IU/L	30.00 - 120.00					
SERUM GAMMA GT Method:- IFCC	30.30	U/L	11.00 - 50.00					
SERUM TOTAL PROTEIN Method:- Biuret Reagent	7.36	g/dl	6.40 - 8.30					
SERUM ALBUMIN Method:- Bromocresol Green	4.50	g/dl	3.80 - 5.00					
SERUM GLOBULIN Method:-CALCULATION	2.86	gm/dl	2.20 - 3.50					
A/G RATIO	1.57		1.30 - 2.50					

Total BilirubinMethodology:Colorimetric method InstrumentName:Randox Rx Imola Interpretation An increase in bilirubin concentration in the serum occurs in toxic or infectious diseases of the liver e.g. hepatitis B or obstruction of the bile duct and in rhesus incompatible babies. High levels of unconjugated bilirubin indicate that too much haemoglobin is being destroyed or that the liver is not actively treating the haemoglobin it is receiving.

AST Aspartate Aminotransferase Methodology: IFCC InstrumentName:Randox Rx Imola Interpretation: Elevated levels of AST can signal myocardial infarction, hepatic disease, muscular dystrophy and organ damage. Although heart muscle is found to have the most activity of the enzyme, significant activity has also been seen in the brain, liver, gastric mucosa, adipose tissue and kidneys of humans. ALT Alanine Aminotransferase Methodology: IFCCInstrumentName:Randox Rx Imola Interpretation: The enzyme ALT has been found to be in highest concentrations in the liver, with decreasing concentrations found in kidney, heart, skeletal muscle, pancreas, spleen and lung tissue respectively. Elevated levels of the transaminases can indicate myocardial infarction, hepatic disease, muscular

Alkaline Phosphatase Methodology; AMP Buffer InstrumentName; Randox Rx Imola Interpretation; Measurements of alkaline phosphatase are of use in the diagnosis, treatment and investigation of Hospitalist Victoria (1997) and in bone disease associated with increased osteoblastic activity. Alkaline phosphatase is also used in the diagnosis of parathyroid and intestinal disease TOTAL PROTEIN Methodology:Biuret Reagent InstrumentName:Randox Rx Imola Interpretation: Measurements obtained by this method are used in the

is and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

ALBUMIN (ALB) Methodology: Bromocresol Green InstrumentName:Randox Rx Imola Interpretation: Albumin measurements are used in the diagnosis and treatment of numerous diseases involving primarily the liver or kidneys. Globulin & A/G ratio is calculated.

Instrument Name Randox Rx Imola Interpretation: Elevations in GGT levels are seen earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or post-hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal)

SURENDRAKHANGA

Page No: 5 of 13







B-51, Ganesh Nagar, Near Metro Piller No. 109-110, New Sanganer Road,

Sodala, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www. drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com | Date :- 23/03/2024 10:26:13

NAME :- Mr. DEVENDER KUMAR

Sex / Age :- Male 47 Yrs 6 Mon 30 Days

Company :- MediWheel

Sample Type :- PLAIN/SERUM

Patient ID :-12236564

Ref. By Dr:- BOB

Lab/Hosp :-

Final Authentication: 23/03/2024 12:40:21

Sample Collected Time 23/03/2024 10:39:42

IMMIINOASSAY

HAMONOASSAI								
Test Name	Value	Unit	Biological Ref Interval					
TOTAL THYROID PROFILE								
SERUM TOTAL T3 Method:- Chemiluminescence(Competitive immunoassay)	1.140	ng/ml	0.970 - 1.690					
SERUM TOTAL T4 Method:- Chemiluminescence(Competitive immunoassay)	7.690	ug/dl	6.530 - 13.210					
SERUM TSH ULTRA Method:- Enhanced Chemiluminescence Immunoassay	2.410	$\mu IU/mL$	0.350 - 5.500					

Interpretation: Triiodothyronine (T3) contributes to the maintenance of the euthyroid state. A decrease in T3 concentration of up to 50% occurs in a variety of clinical situations, including acute and chronic disease. Although T3 results alone cannot be used to diagnose hypothyroidism, T3 concentration may be more sensitive than thyroxine (T4) for hyperthyroidism. Consequently, the total T3 assay can be used in conjunction with other assays to aid in the differential diagnosis of thyroid disease. T3 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, Free T3 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake, or T4 uptake can be used with the total T3 result to calculate the free T3 index and estimate the concentration of free T3.

Interpretation: The measurement of Total T4 aids in the differential diagnosis of thyroid disease. While >99.9% of T4 is protein-bound, primarily to thyroxine-binding globulin (TBG), it is the free fraction that is biologically active. In most patients, the total T4 concentration is a good indicator of thyroid status. T4 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, free T4 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake may be used with the total T4 result to calculate the free T4 index (FT4I) and estimate the concentration of free T4. Some drugs and some nonthyroidal patient conditions are known to alter TT4 concentrations in vivo.

Interpretation: TSH stimulates the production of thyroxine (T4) and triiodothyronine (T3) by the thyroid gland. The diagnosis of overt hypothyroidism by the finding of a low total T4 or free T4 concentration is readily confirmed by a raised TSH concentration. Measurement of low or undetectable TSH concentrations may assist the diagnosis of hyperthyroidism, where concentrations of T4 and T3 are elevated and TSH secretion is suppressed. These have the advantage of discriminating between the concentrations of TSH observed in thyrotoxicosis, compared with the low, but detectable, concentrations that occur in subclinical hyperthyroidism. The performance of this assay has not been established for neonatal specimens. Some drugs and some nonthyroidal patient conditions are known to alter TSH concentrations in vivo.

#### INTERPRETATION

PREGNANCY	REFERENCE RANGE FOR TSH IN uIU/mL (As per American Thyroid
	Association)
1st Trimester	0.10-2.50
2nd Trimester	0.20-3.00
3rd Trimester	0.30-3.00

NARENDRAKUMAR **Technologist** 

Page No: 6 of 13



#### Path Lab & Imaging Centre



B-51, Ganesh Nagar, Near Metro Piller No. 109-110, New Sanganer Road,

Sodala, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www. drgoyalspathlah.com | E-mail: drgoyalpiyush@gmall.com | Date :- 23/03/2024 10:26:13

NAME :- Mr. DEVENDER KUMAR

Sex / Age :- Male 47 Yrs 6 Mon 30 Days

Method:- Reagent Strip(Double indicatior blue reaction)

Company :- MediWheel

Sample Type :- URINE

REACTION(PH)

Patient ID: -12236564

Ref. By Dr:- BOB

Lab/Hosp :-

Fir

5.0 - 7.5

Final Authentication: 23/03/2024 14:08:09

Sample Collected Time 23/03/2024 10:39:42

CLINICAL PATHOLOGY

Test Name	Value Unit	Biological Ref Interval
Urine Routine	*	
PHYSICAL EXAMINATION		
COLOUR	PALE YELLOW	PALE YELLOW
APPEARANCE	Clear	Clear
CHEMICAL EXAMINATION		

SPECIFIC GRAVITY
Method:- Reagent Strip(bromthymol blue)

PROTEIN

1.025

1.010 - 1.030

NIL

NIL

6.0

Method:- Reagent Strip (Sulphosalicylic acid test)

GLUCOSE NIL NIL

Method:- Reagent Strip (Glu.Oxidase Peroxidase Benedict)

BILIRUBIN NEGATIVE NEGATIVE Method:- Reagent Strip (Azo-coupling reaction)

UROBILINOGEN NORMAL NORMAL

Method:- Reagent Strip (Modified ehrlich reaction)

KETONES NEGATIVE NEGATIVE

Method:- Reagent Strip (Sodium Nitropruside) Rothera's

NITRITE

NEGATIVE

NEGATIVE

Method:- Reagent Strip (Diazotization reaction)

RBC NIL NIL

Method:- Reagent Strip (Peroxidase like activity)

MICROSCOPY EXAMINATION

WIICKOSCOPY EXAMINATION			
RBC/HPF	NIL	/HPF	NIL
WBC/HPF	2-3	/HPF	2-3
EPITHELIAL CELLS	1-2	/HPF	2-3
CRYSTALS/HPF	ABSENT		ABSENT
CAST/HPF	ABSENT		ABSENT
AMORPHOUS SEDIMENT	ABSENT		ABSENT
BACTERIAL FLORA	ABSENT		ABSENT
YEAST CELL	ABSENT		ABSENT
OTHER	ABSENT		

VIJENDRAMEENA **Technologist** 

Page No: 7 of 13





B-51, Ganesh Nagar, Near Metro Piller No. 109-110, New Sanganer Road,

Sodala, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www. drgoyalspathlab.com | E-mail: drgoyalpiyush@gmall.com | Date :- 23/03/2024 10:26:13

Patient ID: -12236564

NAME :- Mr. DEVENDER KUMAR

Ref. By Dr:- BOB

Lab/Hosp:-

Sex / Age :- Male

47 Yrs 6 Mon 30 Days Company :- MediWheel

Final Authentication: 24/03/2024 12:11:13

Sample Type :- KOx/Na FLUORIDE-F, KOx/Na Sabbore 1061-PR-PLTAIN-856-783/2024 10:39:42

**BIOCHEMISTRY** 

Test Name	Value	Unit	Biological Ref Interval
FASTING BLOOD SUGAR (Plasma) Method:- GOD PAP	96.7	mg/dl	75.0 - 115.0

Impaired glucose tolerance (IG1)	111 - 125 mg/dL	
Diabetes Mellitus (DM)	> 126 mg/dL	

Instrument Name: Randox Rx Imola Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels(hypoglycemia) may result from excessive insulin therapy or various liver diseases

BLOOD SUGAR PP (Plasma) Method:- GOD PAP

108.8

mg/dl

Instrument Name: Randox Rx Imola Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels(hypoglycemia) may result from excessive insulin therapy or various liver diseases

SERUM CREATININE Method:- Colorimetric Method SERUM URIC ACID Method:- Enzymatic colorimetric 0.95

mg/dl

Men - 0.6-1.30 Women - 0.5-1.20

8.77 H

mg/dl

Men - 3.4-7.0

Women - 2.4-5.7

SURENDRAKHANGA

Page No: 9 of 13



#### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Near Metro Piller No. 109-110, New Sanganer Road,

Sodala, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www. drgoyalspathlab.com | E-majl: drgoyalplyush@gmall.com | Date : 23/03/2024 10:26:13

47 Yrs 6 Mon 30 Days

Patient ID: -12236564

Unit

NAME :- Mr. DEVENDER KUMAR

Ref. By Dr:- BOB

Sex / Age :- Male

Company :- MediWheel

Lab/Hosp:-

Sample Collected Time 23/03/2024 10:39:42

Final Authentication: 23/03/2024 14:08:09

Sample Type :- EDTA, URINE

**HAEMATOLOGY** 

**Biological Ref Interval** 

**BLOOD GROUP ABO** 

**Test Name** 

"A" NEGATIVE

BLOOD GROUP ABO Methodology: Haemagglutination reaction Kit Name: Monoclonal agglutinating antibodies (Span clone).

URINE SUGAR (FASTING) Collected Sample Received

Nil

Value

Nil

AJAYSINGH, VIJENDRAMEENA **Technologist** 

Page No: 11 of 13





B-51, Ganesh Nagar, Near Metro Piller No. 109-110, New Sanganer Road,

Sodala, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www. drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com Date - 23/03/2024 | 10:26:13

47 Yrs 6 Mon 30 Days

Patient ID :-12236564

mg/dl

NAME :- Mr. DEVENDER KUMAR

BLOOD UREA NITROGEN (BUN)

Ref. By Dr:- BOB

Lab/Hosp :-

Company :- MediWheel

Sample Type :- PLAIN/SERUM

Sex / Age :- Male

Sample Collected Time 23/03/2024 10:39:42

Final Authentication: 24/03/2024 12:11:13.

0.0 - 23.0

**BIOCHEMISTRY** 

	Diochen	DICCHEMISTRI		
Test Name	Value	Unit	Biological Ref Interval	
		*		

SURENDRAKHANGA

Page No: 12 of 13





B-51, Ganesh Nagar, Near Metro Piller No. 109-110, New Sanganer Road,

Sodala, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Sex / Age :- Male

Website: www. drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

47 Yrs 6 Mon 30 Days

NAME :- Mr. DEVENDER KUMAR

Patient ID :-12236564

Ref. By Dr:- BOB

Lab/Hosp:-

Company:- MediWheel Sample Type :- PLAIN/SERUM

Sample Collected Time 23/03/2024 10:39:42

Final Authentication: 23/03/2024:12:40:21

**IMMUNOASSAY** 

**Test Name** Value Unit **Biological Ref Interval** 

TOTAL PSA

0.420

ng/ml

0.000 - 4.000

InstrumentName: ADVIA CENTAUR CP Interpretation: Elevated serum PSA concentrations are found in men with prostate cancer, benign prostatic hypertrophy (BHP) or inflammatory conditions of other adjacent genitourinary tissues, but not in apparently healthy men or in men with cancers other than prostate cancer.PSA has been demonstrated to be an accurate marker for monitoring advancing clinical stage in untreated patients and for monitoring response to therapy by radical prostatectomy, radiation therapy and anti-androgen therapy. PSA is also important in determining the potential and actual effectiveness of surgery or other therapies. Progressive disease is defined by an increase of at least 25%. Sampling should be repeated within two to four weeks for additional evidence Different assay methods cannot be used interchangeably.

\*\*\* End of Report \*\*\*

NARENDRAKUMAR **Technologist** 

Page No: 13 of 13



