

CODE/NAME & ADDRESS : C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0061XB000117 PATIENT ID : KARIF02029461

CLIENT PATIENT ID: ABHA NO

AGE/SEX :30 Years Female :02/02/2024 15:06:00 RECEIVED: 02/02/2024 15:07:21 REPORTED :06/02/2024 14:31:18

Test Report Status Results **Biological Reference Interval** Units <u>Final</u>

HAEMATOLOGY - CBC					
MEDI WHEEL FULL BODY HEALTH CHECKUP BE	LOW 40FEMALE				
BLOOD COUNTS,EDTA WHOLE BLOOD					
HEMOGLOBIN (HB)	13.1	12.0 - 15.0	g/dL		
RED BLOOD CELL (RBC) COUNT	4.63	3.8 - 4.8	mil/μL		
WHITE BLOOD CELL (WBC) COUNT	7.64	4.0 - 10.0	thou/µL		
PLATELET COUNT	311	150 - 410	thou/µL		
RBC AND PLATELET INDICES					
HEMATOCRIT (PCV)	40.4	36 - 46	%		
MEAN CORPUSCULAR VOLUME (MCV)	87.3	83 - 101	fL		
MEAN CORPUSCULAR HEMOGLOBIN (MCH)	28.3	27.0 - 32.0	pg		
MEAN CORPUSCULAR HEMOGLOBIN	32.4	31.5 - 34.5	g/dL		
CONCENTRATION (MCHC) RED CELL DISTRIBUTION WIDTH (RDW)	13.6	11.6 - 14.0	%		
MENTZER INDEX	18.9	11.0 - 14.0	70		
	18.9 11.0 High	6.8 - 10.9	fL		
MEAN PLATELET VOLUME (MPV)	11.0 High	0.6 - 10.9	IL.		
WBC DIFFERENTIAL COUNT					
NEUTROPHILS	65	40 - 80	%		
LYMPHOCYTES	30	20 - 40	%		
MONOCYTES	04	2 - 10	%		
EOSINOPHILS	01	1 - 6	%		
BASOPHILS	00	< 1 - 2	%		

BLOOD COUNTS, EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

Dr. Itisha Dhiman **Pathologist**





Page 1 Of 18

View Report

PERFORMED AT:

Agilus Diagnostics Ltd.

M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School

Jodhpur, 342001 Rajasthan, India





Female

PATIENT NAME: KARISHMA REF. DOCTOR: SELF

CODE/NAME & ADDRESS: C000138375

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL
F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHÍ

NEW DELHI 110030

8800465156

ACCESSION NO: 0061XB000117

PATIENT ID : KARIF02029461

CLIENT PATIENT ID: ABHA NO : DRAWN :02/02/2024 15:06:00 RECEIVED :02/02/2024 15:07:21

:30 Years

AGE/SEX

REPORTED :06/02/2024 14:31:18

Test Report Status <u>Final</u> Results Biological Reference Interval Units

RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.

3.3, COVID-19 patients tend to show mild disease.
(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients A.-P. Yang, et al. International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope.

Itisha.

Dr. Itisha Dhiman Pathologist





Page 2 Of 18

View Details

View Report

PERFORMED AT :

Agilus Diagnostics Ltd. M/S S.S. Wellness Centre,Ground Floor,C-22,Shastri Nagar,Near Central Academy School Jodhpur, 342001





mm at 1 hr

REF. DOCTOR: SELF PATIENT NAME: KARISHMA

CODE/NAME & ADDRESS : C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO: 0061XB000117 PATIENT ID : KARIF02029461

CLIENT PATIENT ID:

AGE/SEX :30 Years Female :02/02/2024 15:06:00 DRAWN RECEIVED: 02/02/2024 15:07:21

REPORTED :06/02/2024 14:31:18

Test Report Status Biological Reference Interval Final Results Units

ABHA NO

HAEMATOLOGY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

ERYTHROCYTE SEDIMENTATION RATE (ESR), EDTA BLOOD

10 0 - 20E.S.R

METHOD: WESTERGREN METHOD

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE **BLOOD**

HBA1C 5.3 Non-diabetic: < 5.7 %

> Pre-diabetics: 5.7 - 6.4 Diabetics: > or = 6.5ADA Target: 7.0

Action suggested: > 8.0

ESTIMATED AVERAGE GLUCOSE(EAG) 105.4 < 116.0 mg/dL

Interpretation(s)

ERYTHROCYTE SEDIMENTATION RATE (ESR),EDTA BLOOD-TEST DESCRIPTION :-

Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change.

TEST INTERPRETATION

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy, Estrogen medication, Aging

Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias,

Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis). In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum. Decreased in: Polycythermia vera, Sickle cell anemia

LIMITATIONS

False elevated ESR: Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia False Decreased: Poikilocytosis, (SickleCells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs(Quinine,

salicylates)

REFERENCE:

. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis,10th edition.

Dr. Itisha Dhiman **Pathologist**





Page 3 Of 18

View Report

PERFORMED AT:

Agilus Diagnostics Ltd.

M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School

Jodhpur, 342001 Rajasthan, India





Female

REF. DOCTOR: SELF PATIENT NAME: KARISHMA

CODE/NAME & ADDRESS: C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO: 0061XB000117

PATIENT ID : KARIF02029461

CLIENT PATIENT ID: ABHA NO

:30 Years :02/02/2024 15:06:00 DRAWN

AGE/SEX

RECEIVED: 02/02/2024 15:07:21 REPORTED :06/02/2024 14:31:18

Test Report Status Results **Biological Reference Interval** Units <u>Final</u>

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-Used For:

- 1. Evaluating the long-term control of blood glucose concentrations in diabetic patients.
- 2. Diagnosing diabetes.

3. Identifying patients at increased risk for diabetes (prediabetes).
The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for well-controlled type 2 diabetic patients) to determine whether a patients metabolic control has remained continuously within the target range.

1. eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.

- eAG gives an evaluation of blood glucose levels for the last couple of months.
 eAG is calculated as eAG (mg/dl) = 28.7 * HbA1c 46.7

- **HbA1c Estimation can get affected due to :**1. Shortened Erythrocyte survival : Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss, hemolytic anemia) will falsely lower HbA1c test résults.Fructosamine is recommended in these patients which indicates diabetes control over 15 days.
- 2.Vitamin C & E are reported to falsely lower test results. (possibly by inhibiting glycation of hemoglobin.
 3. Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates
- addiction are reported to interfere with some assay methods, falsely increasing results.

 4. Interference of hemoglobinopathies in HbA1c estimation is seen in
- a) Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c.
- b) Heterozygous state detected (D10 is corrected for HbS & HbC trait.)
- c) HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c.Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy

Dr. Itisha Dhiman

Pathologist



Page 4 Of 18

Agilus Diagnostics Ltd. M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School Jodhpur, 342001

Rajasthan, India





CODE/NAME & ADDRESS: C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO: 0061XB000117

PATIENT ID : KARIF02029461

CLIENT PATIENT ID: ABHA NO

AGE/SEX :30 Years Female :02/02/2024 15:06:00 RECEIVED: 02/02/2024 15:07:21

REPORTED :06/02/2024 14:31:18

Test Report Status Results **Biological Reference Interval** Units <u>Final</u>

IMMUNOHAEMATOLOGY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

TYPE B **ABO GROUP**

METHOD: FORWARD/REVERSE

POSITIVE RH TYPE

METHOD: FORWARD/REVERSE

Interpretation(s)

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same.

The test is performed by both forward as well as reverse grouping methods.

Dr. Itisha Dhiman **Pathologist**





Page 5 Of 18



Agilus Diagnostics Ltd. M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School Jodhpur, 342001

Rajasthan, India





CODE/NAME & ADDRESS: C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO: 0061XB000117 PATIENT ID : KARIF02029461

CLIENT PATIENT ID:

AGE/SEX :30 Years Female :02/02/2024 15:06:00 RECEIVED: 02/02/2024 15:07:21 REPORTED :06/02/2024 14:31:18

mg/dL

Test Report Status Results Biological Reference Interval Units <u>Final</u>

ABHA NO

BIOCHEMISTRY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

GLUCOSE FASTING, FLUORIDE PLASMA

FBS (FASTING BLOOD SUGAR) 95 Normal : < 100

Pre-diabetes: 100-125 Diabetes: >/=126

METHOD: SPECTROPHOTOMETRY

GLUCOSE, POST-PRANDIAL, PLASMA

PPBS(POST PRANDIAL BLOOD SUGAR) 98 70 - 140 mg/dL

METHOD: SPECTROPHOTOMETRY

LIPID PROFILE WITH CALCULATED LDL

CHOLESTEROL, TOTAL 174 < 200 Desirable mg/dL

200 - 239 Borderline High

>/= 240 High METHOD: SPECTROPHOTOMETRY

TRIGLYCERIDES 48 < 150 Normal mg/dL

150 - 199 Borderline High

200 - 499 High >/=500 Very High

METHOD: SPECTROPHOTOMETRY

55 mg/dL HDL CHOLESTEROL < 40 Low

>/=60 High

METHOD: SPECTROPHOTOMETRY

CHOLESTEROL LDL 109 High < 100 Optimal mg/dL

100 - 129

Near optimal/ above optimal

130 - 159 Borderline High 160 - 189 High >/= 190 Very High

Dr. Itisha Dhiman **Pathologist**





Page 6 Of 18



Agilus Diagnostics Ltd. M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School Jodhpur, 342001

Rajasthan, India





PATIENT NAME: KARISHMA REF. DOCTOR: SELF CODE/NAME & ADDRESS : C000138375 ACCESSION NO: 0061XB000117 AGE/SEX :30 Years Female ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID : KARIF02029461 :02/02/2024 15:06:00 F-703, LADO SARAI, MEHRAULISOUTH WEST CLIENT PATIENT ID: RECEIVED: 02/02/2024 15:07:21 DELHI REPORTED :06/02/2024 14:31:18 ABHA NO **NEW DELHI 110030** 8800465156

Test Report Status <u>Final</u>	Results	Biological Reference Interval Units
NON HDL CHOLESTEROL	119	Desirable: Less than 130 mg/dL Above Desirable: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very high: > or = 220
VERY LOW DENSITY LIPOPROTEIN	9.6	= 30.0 mg/dL</td
CHOL/HDL RATIO	3.2 Low	3.3 - 4.4 Low Risk 4.5 - 7.0 Average Risk 7.1 - 11.0 Moderate Risk > 11.0 High Risk
LDL/HDL RATIO	2.0	0.5 - 3.0 Desirable/Low Risk 3.1 - 6.0 Borderline/Moderate Risk >6.0 High Risk

Interpretation(s)

Serum lipid profile is measured for cardiovascular risk prediction. Lipid Association of India recommends LDL-C as primary target and Non HDL-C as co-primary treatment target.

Risk Stratification for ASCVD (Atherosclerotic cardiovascular disease) by Lipid Association of India

A.CAD with > 1 feature of high risk group			
B. CAD with > 1 feature of Very high risk g	group or recurrent ACS (within 1 year) despite LDL-C < or =		
50 mg/dl or polyvascular disease			
	najor risk factors or evidence of end organ damage 3.		
Familial Homozygous Hypercholesterolemia	a		
1. Three major ASCVD risk factors. 2. Dia	betes with 1 major risk factor or no evidence of end organ		
damage. 3. CKD stage 3B or 4. 4. LDL >190 mg/dl 5. Extreme of a single risk factor. 6. Coronary			
Artery Calcium - CAC >300 AU. 7. Lipoprotein a >/= 50mg/dl 8. Non stenotic carotid plaque			
2 major ASCVD risk factors			
0-1 major ASCVD risk factors			
Major ASCVD (Atherosclerotic cardiovascular disease) Risk Factors			
1. Age > or = 45 years in males and > or = 55 years in females 3. Current Cigarette smoking or tobacco use			
2. Family history of premature ASCVD 4. High blood pressure			
5. Low HDL			
	B. CAD with > 1 feature of Very high risk g 50 mg/dl or polyvascular disease 1. Established ASCVD 2. Diabetes with 2 r Familial Homozygous Hypercholesterolemi 1. Three major ASCVD risk factors. 2. Dia damage. 3. CKD stage 3B or 4. 4. LDL > 1 Artery Calcium - CAC > 300 AU. 7. Lipopr 2 major ASCVD risk factors 0-1 major ASCVD risk factors erosclerotic cardiovascular disease) Risk Fa s in males and > or = 55 years in females		

Dr. Itisha Dhiman Pathologist



Page 7 Of 18

View Details

View Report

PERFORMED AT:

Agilus Diagnostics Ltd.

M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School

Jodhpur, 342001 Rajasthan, India





PATIENT NAME: KARISHMA REF. DOCTOR: SELF CODE/NAME & ADDRESS : C000138375 ACCESSION NO: 0061XB000117 AGE/SEX :30 Years Female ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID DRAWN :02/02/2024 15:06:00 : KARIF02029461 F-703, LADO SARAI, MEHRAULISOUTH WEST CLIENT PATIENT ID: RECEIVED: 02/02/2024 15:07:21 DELHI ABHA NO REPORTED :06/02/2024 14:31:18 **NEW DELHI 110030** 8800465156

Test Report Status <u>Final</u> Results Biological Reference Interval Units

Newer treatment goals and statin initiation thresholds based on the risk categories proposed by LAI in 2020.

Risk Group	Treatment Goals		Consider Drug Therapy	
	LDL-C (mg/dl)	Non-HDL (mg/dl)	LDL-C (mg/dl)	Non-HDL (mg/dl)
Extreme Risk Group Category A	<50 (Optional goal < OR = 30)	< 80 (Optional goal <or 60)<="" =="" td=""><td>>OR = 50</td><td>>OR = 80</td></or>	>OR = 50	>OR = 80
Extreme Risk Group Category B	<OR = 30	<or 60<="" =="" td=""><td>> 30</td><td>>60</td></or>	> 30	>60
Very High Risk	<50	<80	>OR= 50	>OR= 80
High Risk	<70	<100	>OR= 70	>OR= 100
Moderate Risk	<100	<130	>OR= 100	>OR= 130
Low Risk	<100	<130	>OR= 130*	>OR= 160

^{*}After an adequate non-pharmacological intervention for at least 3 months.

References: Management of Dyslipidaemia for the Prevention of Stroke: Clinical Practice Recommendations from the Lipid Association of India. Current Vascular Pharmacology, 2022, 20, 134-155.

LIVER FUNCTION PROFILE, SERUM

BILIRUBIN, TOTAL	0.60	0.2 - 1.0	mg/dL
METHOD: SPECTROPHOTOMETRY			
BILIRUBIN, DIRECT	0.2	0.0 - 0.2	mg/dL
METHOD: SPECTROPHOTOMETRY			
BILIRUBIN, INDIRECT	0.4	0.1 - 1.0	mg/dL
METHOD: SPECTROPHOTOMETRY			
TOTAL PROTEIN	7.5	6.4 - 8.2	g/dL
METHOD: SPECTROPHOTOMETRY			
ALBUMIN	3.9	3.4 - 5.0	g/dL
METHOD: SPECTROPHOTOMETRY			
GLOBULIN	3.6	2.0 - 4.1	g/dL
METHOD: CALCULATED PARAMETER			
ALBUMIN/GLOBULIN RATIO	1.1	1.0 - 2.1	RATIO
METHOD: CALCULATED PARAMETER			
ASPARTATE AMINOTRANSFERASE(AST/SGOT)	45 High	15 - 37	U/L
METHOD: SPECTROPHOTOMETRY			
ALANINE AMINOTRANSFERASE (ALT/SGPT)	63 High	< 34.0	U/L
METHOD: SPECTROPHOTOMETRY			
ALKALINE PHOSPHATASE	96	30 - 120	U/L
METHOD: SPECTROPHOTOMETRY			
GAMMA GLUTAMYL TRANSFERASE (GGT)	33	5 - 55	U/L
METHOD: SPECTROPHOTOMETRY			
LACTATE DEHYDROGENASE	145	81 - 234	U/L
METHOD: SPECTROPHOTOMETRY			

700 /

Dr. Itisha Dhiman Pathologist





Page 8 Of 18

View Details

View Report

PERFORMED AT:

Agilus Diagnostics Ltd.

M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School

Jodhpur, 342001 Rajasthan, India





CODE/NAME & ADDRESS: C000138375

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL
F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHÍ

NEW DELHI 110030 8800465156 ACCESSION NO: **0061XB000117**PATIENT ID: KARIF02029461

CLIENT PATIENT ID: ABHA NO : AGE/SEX :30 Years Female
DRAWN :02/02/2024 15:06:00
RECEIVED :02/02/2024 15:07:21
REPORTED :06/02/2024 14:31:18

Test Report Status <u>Final</u>	Results	Biological Reference I	Interval Units
BLOOD UREA NITROGEN (BUN), SERUM			
BLOOD UREA NITROGEN METHOD: SPECTROPHOTOMETRY	6	6 - 20	mg/dL
CREATININE, SERUM			
CREATININE METHOD: SPECTROPHOTOMETRY	0.61	0.60 - 1.10	mg/dL
BUN/CREAT RATIO			
BUN/CREAT RATIO METHOD: SPECTROPHOTOMETRY	9.84	5.00 - 15.00	
URIC ACID, SERUM			
URIC ACID METHOD: SPECTROPHOTOMETRY	3.6	2.6 - 6.0	mg/dL
TOTAL PROTEIN, SERUM			
TOTAL PROTEIN METHOD: SPECTROPHOTOMETRY	7.5	6.4 - 8.2	g/dL
ALBUMIN, SERUM			
ALBUMIN METHOD: SPECTROPHOTOMETRY	3.9	3.4 - 5.0	g/dL
GLOBULIN			
GLOBULIN METHOD: CALCULATED PARAMETER	3.6	2.0 - 4.1	g/dL

Dr. Itisha Dhiman Pathologist



Page 9 Of 18

View Details

View Report

PERFORMED AT :

Agilus Diagnostics Ltd. M/S S.S. Wellness Centre,Ground Floor,C-22,Shastri Nagar,Near Central Academy School Jodhpur, 342001





CODE/NAME & ADDRESS: C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO: 0061XB000117 : KARIF02029461

PATIENT ID CLIENT PATIENT ID:

ABHA NO

AGE/SEX :30 Years Female :02/02/2024 15:06:00 RECEIVED: 02/02/2024 15:07:21 REPORTED :06/02/2024 14:31:18

Test Report Status Results **Biological Reference Interval** Units <u>Final</u>

ELECTROLYTES (NA/K/CL), SERUM

SODIUM, SERUM 139 136 - 145 mmol/L

METHOD: ION SELECTIVE ELECTRODE TECHNOLOGY

POTASSIUM, SERUM 3.50 - 5.10 mmol/L 4.4

METHOD: ION SELECTIVE ELECTRODE TECHNOLOGY

110 High 98 - 107 mmol/L CHLORIDE, SERUM

METHOD: ION SELECTIVE ELECTRODE TECHNOLOGY

Interpretation(s)

Sodium	Potassium	Chloride
Decreased in: CCF, cirrhosis,	Decreased in: Low potassium	Decreased in: Vomiting, diarrhea,
vomiting, diarrhea, excessive	intake,prolonged vomiting or diarrhea,	renal failure combined with salt
sweating, salt-losing	RTA types I and II,	deprivation, over-treatment with
nephropathy, adrenal insufficiency,	hyperaldosteronism, Cushing's	diuretics, chronic respiratory acidosis,
nephrotic syndrome, water	syndrome,osmotic diuresis (e.g.,	diabetic ketoacidosis, excessive
intoxication, SIADH. Drugs:	hyperglycemia),alkalosis, familial	sweating, SIADH, salt-losing
thiazides, diuretics, ACE inhibitors,	periodic paralysis,trauma	nephropathy, porphyria, expansion of
chlorpropamide,carbamazepine,anti	(transient).Drugs: Adrenergic agents,	extracellular fluid volume,
depressants (SSRI), antipsychotics.	diuretics.	adrenalinsufficiency,
		hyperaldosteronism, metabolic
		alkalosis. Drugs: chronic
		laxative,corticosteroids, diuretics.
Increased in: Dehydration	Increased in: Massive hemolysis,	Increased in: Renal failure, nephrotic
(excessivesweating, severe	severe tissue damage, rhabdomyolysis,	syndrome, RTA,dehydration,
vomiting or diarrhea),diabetes	acidosis, dehydration,renal failure,	overtreatment with
mellitus, diabetesinsipidus,	Addison's disease, RTA type IV,	saline,hyperparathyroidism, diabetes
hyperaldosteronism, inadequate	hyperkalemic familial periodic	insipidus, metabolic acidosis from
water intake. Drugs: steroids,	paralysis. Drugs: potassium salts,	diarrhea (Loss of HCO3-), respiratory
licorice,oral contraceptives.	potassium- sparing diuretics,NSAIDs,	alkalosis,hyperadrenocorticism.
	beta-blockers, ACE inhibitors, high-	Drugs: acetazolamide,androgens,
	dose trimethoprim-sulfamethoxazole.	hydrochlorothiazide, salicylates.
Interferences: Severe lipemia or	Interferences: Hemolysis of sample,	Interferences:Test is helpful in
hyperproteinemi, if sodium analysis	delayed separation of serum,	assessing normal and increased anion
involves a dilution step can cause	prolonged fist clenching during blood	gap metabolic acidosis and in
spurious results. The serum sodium	drawing, and prolonged tourniquet	distinguishing hypercalcemia due to
falls about 1.6 mEq/L for each 100	placement. Very high WBC/PLT counts	hyperparathyroidism (high serum
mg/dL increase in blood glucose.	may cause spurious. Plasma potassium	chloride) from that due to malignancy
	levels are normal.	(Normal serum chloride)

Interpretation(s)
GLUCOSE FASTING,FLUORIDE PLASMA-TEST DESCRIPTION

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in the

Dr. Itisha Dhiman **Pathologist**





Page 10 Of 18

PERFORMED AT:

Agilus Diagnostics Ltd.

M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School

Jodhpur, 342001 Rajasthan, India





Female

REF. DOCTOR: SELF PATIENT NAME: KARISHMA

CODE/NAME & ADDRESS: C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO : 0061XB000117

PATIENT ID : KARIF02029461

CLIENT PATIENT ID: ABHA NO

:02/02/2024 15:06:00 DRAWN RECEIVED: 02/02/2024 15:07:21

:30 Years

AGE/SEX

REPORTED :06/02/2024 14:31:18

Test Report Status Results **Biological Reference Interval Final** Units

Increased in:Diabetes mellitus, Cushing's syndrome (10 – 15%), chronic pancreatitis (30%). Drugs:corticosteroids,phenytoin, estrogen, thiazides.

Decreased in:Pancreatic islet cell disease with increased insulin,insulinoma,adrenocortical insufficiency,hypopituitarism,diffuse liver disease, malignancy(adrenocortical,stomach,fibrosarcoma),infant of a diabetic mother,enzyme deficiency diseases(e.g.galactosemia),Drugs-insulin,ethanol,propranolol

sulfonylureas,tolbutamide,and other oral hypoglycemic agents.

NOTE: While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values),there is wide fluctuation within

individuals.Thus, glycosylated hemoglobin(HbA1c) levels are favored to monitor glycemic control.

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment,Renal Glyosuria,Glycaemic

index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.

GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc. Additional test HbA1c LIVER FUNCTION PROFILE, SERUM
Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give

yellow discoloration in jaundice. **Elevated levels** results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis,

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Pagets disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilsons disease.

GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease,high alcohol consumption and use of enzyme-inducing drugs etc. **Total Protein** also known as total protein,is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and

globulin.Higher-than-normal levels may be due to:Chronic inflammation or infection,including HIV and hepatitis B or C,Multiple myeloma,Waldenstrom disease.Lower-than-normal levels may be due to: Agammaglobulinemia,Bleeding (hemorrhage),Burns,Glomerulonephritis,Liver disease, Malabsorption,Malnutrition,Nephrotic syndrome, Protein-losing enteropathy etc.

syndrome, Protein-losing enteropathy etc.

Albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc

BLOOD UREA NITROGEN (BUN), SERUM-Causes of Increased levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)

Causes of decreased level include Liver disease, SIADH.

CREATININE, SERUM-Higher than normal level may be due to:

• Blockage in the urinary tract, Kidney problems, such as kidney damage or failure, infection, or reduced blood flow, Loss of body fluid (dehydration), Muscle problems, such as breakdown of muscle fibers, Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:• Myasthenia Gravis, Muscuophy

URIC ACID, SERUM-Causes of Increased levels:-Dietary(High Protein Intake,Prolonged Fasting,Rapid weight loss),Gout,Lesch nyhan syndrome,Type 2 DM,Metabolic syndrome Causes of decreased levels-Low Zinc intake,OCP,Multiple Sclerosis

TOTAL PROTEIN, SERUM-is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease.

Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic

syndrome, Protein-losing enteropathy etc.
ALBUMIN, SERUM-Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance,malnutrition and wasting etc.

Dr. Itisha Dhiman

Pathologist

Page 11 Of 18





View Report

PERFORMED AT:

Agilus Diagnostics Ltd. M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School Jodhpur, 342001

CIN - U74899PB1995PLC045956 Email: srl.jodhpur@gmail.com





CODE/NAME & ADDRESS: C000138375

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL
F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHÍ

NEW DELHI 110030 8800465156 ACCESSION NO: **0061XB000117**PATIENT ID : KARIF02029461

CLIENT PATIENT ID: ABHA NO : AGE/SEX :30 Years Female
DRAWN :02/02/2024 15:06:00
RECEIVED :02/02/2024 15:07:21
REPORTED :06/02/2024 14:31:18

Test Report Status <u>Final</u> Results Biological Reference Interval Units

CLINICAL PATH - URINALYSIS

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

PHYSICAL EXAMINATION, URINE

COLOR PALE YELLOW

APPEARANCE HAZY

CHEMICAL EXAMINATION, URINE

PH	7.0	4.7 - 7.5
SPECIFIC GRAVITY	1.020	1.003 - 1.035
PROTEIN	NOT DETECTED	NOT DETECTED
GLUCOSE	NOT DETECTED	NOT DETECTED
KETONES	NOT DETECTED	NOT DETECTED
BLOOD	NOT DETECTED	NOT DETECTED
BILIRUBIN	NOT DETECTED	NOT DETECTED
UROBILINOGEN	NORMAL	NORMAL
NITRITE	NOT DETECTED	NOT DETECTED
LEUKOCYTE ESTERASE	DETECTED (+)	NOT DETECTED

MICROSCOPIC EXAMINATION, URINE

EPITHELIAL CELLS	20-30	0-5	/HPF
PUS CELL (WBC'S)	3-5	0-5	/HPF
RED BLOOD CELLS	NOT DETECTED	NOT DETECTED	/HPF

CASTS NOT DETECTED
CRYSTALS NOT DETECTED

BACTERIA **DETECTED (++)** NOT DETECTED

 ${\tt METHOD}: {\tt MICROSCOPIC} \ {\tt EXAMINATION}$

YEAST NOT DETECTED NOT DETECTED

Dr. Itisha Dhiman Pathologist



Page 12 Of 18

View Details

View Repor

PERFORMED AT :

Agilus Diagnostics Ltd. M/S S.S. Wellness Centre,Ground Floor,C-22,Shastri Nagar,Near Central Academy School Jodhpur, 342001

Rajasthan, India





PATIENT NAME: KARISHMA REF. DOCTOR: SELF CODE/NAME & ADDRESS : C000138375 ACCESSION NO: 0061XB000117 AGE/SEX :30 Years Female ARCOFEMI HEALTHCARE LTD (MEDIWHEEL :02/02/2024 15:06:00 PATIENT ID : KARIF02029461 DRAWN F-703, LADO SARAI, MEHRAULISOUTH WEST CLIENT PATIENT ID: RECEIVED : 02/02/2024 15:07:21 DELHI REPORTED :06/02/2024 14:31:18 ABHA NO **NEW DELHI 110030** 8800465156

Test Report Status <u>Final</u> Results Biological Reference Interval Units

Interpretation(s)

The following table describes the probable conditions, in which the analytes are present in urine

Presence of	Conditions	
Proteins	Inflammation or immune illnesses	
Pus (White Blood Cells)	Urinary tract infection, urinary tract or kidney stone, tumors or any kind	
·	of kidney impairment	
Glucose	Diabetes or kidney disease	
Ketones	Diabetic ketoacidosis (DKA), starvation or thirst	
Urobilinogen	Liver disease such as hepatitis or cirrhosis	
Blood	Renal or genital disorders/trauma	
Bilirubin	Liver disease	
Erythrocytes	Urological diseases (e.g. kidney and bladder cancer, urolithiasis), urinary tract infection and glomerular diseases	
Leukocytes	Urinary tract infection, glomerulonephritis, interstitial nephritis either acute or chronic, polycystic kidney disease, urolithiasis, contamination by genital secretions	
Epithelial cells	Urolithiasis, bladder carcinoma or hydronephrosis, ureteric stents or bladder catheters for prolonged periods of time	
Granular Casts	Low intratubular pH, high urine osmolality and sodium concentration, interaction with Bence-Jones protein	
Hyaline casts	Physical stress, fever, dehydration, acute congestive heart failure, renal diseases	
Calcium oxalate	Metabolic stone disease, primary or secondary hyperoxaluria, intravenous infusion of large doses of vitamin C, the use of vasodilator naftidrofuryl oxalate or the gastrointestinal lipase inhibitor orlistat, ingestion of ethylene glycol or of star fruit (Averrhoa carambola) or its juice	
Uric acid	arthritis	
Bacteria	Urinary infectionwhen present in significant numbers & with pus cells.	
Trichomonas vaginalis	Vaginitis, cervicitis or salpingitis	

Dr. Itisha Dhiman Pathologist



Page 13 Of 18

View Details



PERFORMED AT:

Agilus Diagnostics Ltd. M/S S.S. Wellness Centre,Ground Floor,C-22,Shastri Nagar,Near Central Academy School Jodhpur, 342001

Rajasthan, India





CODE/NAME & ADDRESS: C000138375

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL
F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHÍ

NEW DELHI 110030

8800465156

ACCESSION NO : 0061XB000117

PATIENT ID : KARIF02029461 CLIENT PATIENT ID:

ABHA NO :

AGE/SEX :30 Years Female
DRAWN :02/02/2024 15:06:00
RECEIVED :02/02/2024 15:07:21
REPORTED :06/02/2024 14:31:18

Test Report Status <u>Final</u> Results Biological Reference Interval Units

CYTOLOGY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

PAPANICOLAOU SMEAR

TEST METHOD CONVENTIONAL GYNEC CYTOLOGY

SPECIMEN TYPE TWO UNSTAINED CERVICAL SMEARS RECEIVED

REPORTING SYSTEM 2014 BETHESDA SYSTEM FOR REPORTING CERVICAL CYTOLOGY

SPECIMEN ADEQUACY SMEARS ARE SATISFACTORY FOR EVALUATION.

MICROSCOPY SMEARS SHOW SUPERFICIAL AND INTERMEDIATE SQUAMOUS

CELLS.

BACKGROUND SHOW LACTOBACILLI AND IS CLEAR.

ENDOCERVICAL COMPONENT ABSENT.

METHOD: MANUAL

INTERPRETATION / RESULT NEGATIVE FOR INTRAEPITHELIAL LESION OR MALIGNANCY

grisha.

Dr. Itisha Dhiman Pathologist





Page 14 Of 18

View Details

View Report



Agilus Diagnostics Ltd. M/S S.S. Wellness Centre,Ground Floor,C-22,Shastri Nagar,Near Central Academy School Jodhpur, 342001





CODE/NAME & ADDRESS: C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO: 0061XB000117 PATIENT ID : KARIF02029461

CLIENT PATIENT ID:

AGE/SEX :30 Years Female :02/02/2024 15:06:00 RECEIVED: 02/02/2024 15:07:21

REPORTED :06/02/2024 14:31:18

Test Report Status Results **Biological Reference Interval Units** <u>Final</u>

ABHA NO

CLINICAL PATH - STOOL ANALYSIS

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE PHYSICAL EXAMINATION, STOOL

COLOUR SAMPLE NOT RECEIVED

Dr. Itisha Dhiman

Pathologist



Page 15 Of 18





Agilus Diagnostics Ltd. M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School Jodhpur, 342001

Rajasthan, India





REF. DOCTOR: SELF PATIENT NAME: KARISHMA

CODE/NAME & ADDRESS: C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO: 0061XB000117 PATIENT ID : KARIF02029461

CLIENT PATIENT ID: ABHA NO

AGE/SEX :30 Years Female DRAWN :02/02/2024 15:06:00 RECEIVED: 02/02/2024 15:07:21 REPORTED :06/02/2024 14:31:18

Test Report Status Results **Biological Reference Interval** Units **Final**

SPECIALISED CHEMISTRY - HORMONE

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

THYROID PANEL, SERUM			
ТЗ	112.30	Non-Pregnant Women 80.0 - 200.0 Pregnant Women 1st Trimester:105.0 - 230.0 2nd Trimester:129.0 - 262.0 3rd Trimester:135.0 - 262.0	
T4	7.59	Non-Pregnant Women 5.10 - 14.10 Pregnant Women 1st Trimester: 7.33 - 14.80 2nd Trimester: 7.93 - 16.10 3rd Trimester: 6.95 - 15.70	μg/dL
TSH (ULTRASENSITIVE)	4.010	Non Pregnant Women 0.27 - 4.20 Pregnant Women (As per American Thyroid Associatio 1st Trimester 0.100 - 2.500 2nd Trimester 0.200 - 3.000 3rd Trimester 0.300 - 3.000	

Interpretation(s)

Triiodothyronine T3, Thyroxine T4, and Thyroid Stimulating Hormone TSH are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hyporthyroidism, TSH levels are low. Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3. Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, Free T4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Total T4 FT4 Total T3

Dr. Itisha Dhiman **Pathologist**



Page 16 Of 18

PERFORMED AT:

Agilus Diagnostics Ltd.

M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School Jodhpur, 342001





PATIENT NAME: KARISHMA REF. DOCTOR: SELF CODE/NAME & ADDRESS: C000138375 ACCESSION NO: 0061XB000117 AGE/SEX :30 Years Female ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID : KARIF02029461 DRAWN :02/02/2024 15:06:00 F-703, LADO SARAI, MEHRAULISOUTH WEST CLIENT PATIENT ID: RECEIVED: 02/02/2024 15:07:21 DELHI ABHA NO REPORTED :06/02/2024 14:31:18 **NEW DELHI 110030** 8800465156

Test Report Status Final Results Biological Reference Interval Units

1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3)
1	Ingh	Low	Low	Low	Post Thyroidectomy (4) Post Radio-Iodine treatment
2	High	Normal	Normal	Normal	(1)Subclinical Hypothyroidism (2) Patient with insufficient thyroid hormone replacement therapy (3) In cases of Autoimmune/Hashimoto thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical inflammation, drugs like amphetamines, Iodine containing drug and dopamine antagonist e.g. domperidone and other physiological reasons.
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre (3)Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4 replacement therapy (7) First trimester of Pregnancy
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism
6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent treatment for Hyperthyroidism
8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies

REF: 1. TIETZ Fundamentals of Clinical chemistry 2.Guidlines of the American Thyroid association during pregnancy and Postpartum, 2011. **NOTE: It is advisable to detect Free T3,FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.**TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

End Of Report
Please visit www.agilusdiagnostics.com for related Test Information for this accession

Dr. Itisha Dhiman

Pathologist



Page 17 Of 18

View Details

View Report

PERFORMED AT:

Agilus Diagnostics Ltd. M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School Jodhpur, 342001

Tel : 0291-2646000, 2644000, Fax CIN - U74899PB1995PLC045956 Email : srl.jodhpur@gmail.com



Rajasthan, India Tel: 0291-2646000, 2644000, Fax:



REF. DOCTOR: SELF PATIENT NAME: KARISHMA

CODE/NAME & ADDRESS : C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0061XB000117 PATIENT ID : KARIF02029461

CLIENT PATIENT ID: ABHA NO

AGE/SEX :30 Years Female :02/02/2024 15:06:00 RECEIVED: 02/02/2024 15:07:21 REPORTED :06/02/2024 14:31:18

Test Report Status Results Biological Reference Interval Units **Final**

CONDITIONS OF LABORATORY TESTING & REPORTING

- 1. It is presumed that the test sample belongs to the patient named or identified in the test requisition form.
- 2. All tests are performed and reported as per the turnaround time stated in the AGILUS Directory of Services.
- 3. Result delays could occur due to unforeseen circumstances such as non-availability of kits / equipment breakdown / natural calamities / technical downtime or any other unforeseen event.
- 4. A requested test might not be performed if:
 - i. Specimen received is insufficient or inappropriate
 - ii. Specimen quality is unsatisfactory
 - iii. Incorrect specimen type
 - iv. Discrepancy between identification on specimen container label and test requisition form

- 5. AGILUS Diagnostics confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity.
- 6. Laboratory results should not be interpreted in isolation; it must be correlated with clinical information and be interpreted by registered medical practitioners only to determine final diagnosis.
- Test results may vary based on time of collection, physiological condition of the patient, current medication or nutritional and dietary changes. Please consult your doctor or call us for any clarification.
- Test results cannot be used for Medico legal purposes.
- 9. In case of queries please call customer care (91115 91115) within 48 hours of the report.

Agilus Diagnostics Ltd

Fortis Hospital, Sector 62, Phase VIII, Mohali 160062

Dr. Itisha Dhiman

Pathologist

Page 18 Of 18





Agilus Diagnostics Ltd. M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School Jodhpur, 342001

Tel: 0291-2646000, 2644000, Fax: CIN - U74899PB1995PLC045956



