PID No.
 : MED112152459
 Register On
 : 20/04/2024 10:38 AM

 SID No.
 : 424023746
 Collection On
 : 20/04/2024 11:03 AM

 Age / Sex
 : 31 Year(s) / Male
 Report On
 : 20/04/2024 5:47 PM

 Type
 : OP
 Printed On
 : 22/04/2024 12:31 PM

Ref. Dr : MediWheel

Investigation BLOOD GROUPING AND Rh TYPING (EDTA Blood/Agglutination)	Observed Value 'O' 'Positive'	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood'Spectrophotometry)	14.7	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	43.7	%	42 - 52
RBC Count (EDTA Blood)	4.26	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (EDTA Blood)	102.8	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood)	34.6	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood)	33.7	g/dL	32 - 36
RDW-CV (EDTA Blood)	14.2	%	11.5 - 16.0
RDW-SD (EDTA Blood)	51.6	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood)	8600	cells/cu.mm	4000 - 11000
Neutrophils (EDTA Blood)	58.4	%	40 - 75
Lymphocytes (EDTA Blood)	28.0	%	20 - 45
Eosinophils (EDTA Blood)	4.8	%	01 - 06
Monocytes (EDTA Blood)	8.5	%	01 - 10





 PID No.
 : MED112152459
 Register On
 : 20/04/2024 10:38 AM

 SID No.
 : 424023746
 Collection On
 : 20/04/2024 11:03 AM

 Age / Sex
 : 31 Year(s) / Male
 Report On
 : 20/04/2024 5:47 PM

 Type
 : OP
 Printed On
 : 22/04/2024 12:31 PM

Ref. Dr : MediWheel

Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Basophils (Blood)	0.3	%	00 - 02
INTERPRETATION: Tests done on Automated I	Five Part cell count	er. All abnormal results a	re reviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood)	5.0	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood)	2.4	10^3 / μ1	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood)	0.4	10^3 / µl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood)	0.7	10^3 / µl	< 1.0
Absolute Basophil count (EDTA Blood)	0.0	10^3 / μl	< 0.2
Platelet Count (EDTA Blood)	263	10^3 / µl	150 - 450
MPV (EDTA Blood)	10.0	fL	7.9 - 13.7
PCT (EDTA Blood/Automated Blood cell Counter)	0.264	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood)	15	mm/hr	< 15
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	90.88	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine) (Urine - F/GOD - POD)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD-PAP)	87.43	mg/dL	70 - 140





 PID No.
 : MED112152459
 Register On
 : 20/04/2024 10:38 AM

 SID No.
 : 424023746
 Collection On
 : 20/04/2024 11:03 AM

 Age / Sex
 : 31 Year(s) / Male
 Report On
 : 20/04/2024 5:47 PM

 Type
 : OP
 Printed On
 : 22/04/2024 12:31 PM

Ref. Dr : MediWheel

<u>Investigation</u>	Observed Unit	<u>Biological</u>
-	<u>Value</u>	Reference Interval

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	6.4	mg/dL	7.0 - 21
Creatinine (Serum/Modified Jaffe)	0.96	mg/dL	0.9 - 1.3

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetylcyteine, chemotherapeutic agent such as flucytosine etc.

Uric Acid (Serum/Enzymatic) <u>Liver Function Test</u>	6.50	mg/dL	3.5 - 7.2
Bilirubin(Total) (Serum/DCA with ATCS)	0.60	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.18	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.42	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	21.73	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	34.21	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	32.87	U/L	< 55
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	114.5	U/L	53 - 128





APPROVED BY

The results pertain to sample tested.

 PID No.
 : MED112152459
 Register On
 : 20/04/2024 10:38 AM

 SID No.
 : 424023746
 Collection On
 : 20/04/2024 11:03 AM

 Age / Sex
 : 31 Year(s) / Male
 Report On
 : 20/04/2024 5:47 PM

 Type
 : OP
 Printed On
 : 22/04/2024 12:31 PM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Total Protein (Serum/Biuret)	7.07	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.52	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	2.55	gm/dL	2.3 - 3.6
A : G RATIO (Serum/Derived)	1.77		1.1 - 2.2
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	169.37	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	147.70	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the õusualö circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	40.60	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	99.3	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	29.5	mg/dL	< 30





 PID No.
 : MED112152459
 Register On
 : 20/04/2024 10:38 AM

 SID No.
 : 424023746
 Collection On
 : 20/04/2024 11:03 AM

 Age / Sex
 : 31 Year(s) / Male
 Report On
 : 20/04/2024 5:47 PM

 Type
 : OP
 Printed On
 : 22/04/2024 12:31 PM

Ref. Dr : MediWheel

<u>Investigation</u>	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Non HDL Cholesterol (Serum/Calculated)	128.8	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2. It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

4.2

	Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
3.6	Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
2.4	Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0

Glycosylated Haemoglobin (HbA1c)

Total Cholesterol/HDL Cholesterol

HbA1C 5.2 % Normal: 4.5 - 5.6 (Whole Blood/HPLC) Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 102.54 mg/dL

(Whole Blood)





Optimal: < 3.3

 PID No.
 : MED112152459
 Register On
 : 20/04/2024 10:38 AM

 SID No.
 : 424023746
 Collection On
 : 20/04/2024 11:03 AM

 Age / Sex
 : 31 Year(s) / Male
 Report On
 : 20/04/2024 5:47 PM

 Type
 : OP
 Printed On
 : 22/04/2024 12:31 PM

Ref. Dr : MediWheel

Investigation Observed Unit Biological Value Reference Interval

INTERPRETATION: Comments

(Serum/Manometric method)

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.

Prostate specific antigen - Total(PSA)

0.825

ng/ml

Normal: 0.0 - 4.0 Inflammatory & Non Malignant conditions of Prostate & genitourinary system: 4.01 - 10.0 Suspicious of Malignant disease of

Prostate: > 10.0

INTERPRETATION: Analytical sensitivity: 0.008 - 100 ng/mL

PSA is a tumor marker for screening of prostate cancer. Increased levels of PSA are associated with prostate cancer and benign conditions like bacterial infection, inflammation of prostate gland and benign hypertrophy of prostate/ benign prostatic hyperplasia (BPH).

Transient elevation of PSA levels are seen following digital rectal examination, rigorous physical activity like bicycle riding, ejaculation within 24 hours.

PSA levels tend to increase in all men as they age.

Clinical Utility of PSA:

ÉIn the early detection of Prostate cancer.

ÉAs an aid in discriminating between Prostate cancer and Benign Prostatic disease.

ÉTo detect cancer recurrence or disease progression.

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.01 ng/ml 0.7 - 2.04

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Tyroxine) - Total $6.79 \mu g/dl$ 4.2 - 12.0

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.





APPROVED BY

The results pertain to sample tested.

Page 6 of 10

 PID No.
 : MED112152459
 Register On
 : 20/04/2024 10:38 AM

 SID No.
 : 424023746
 Collection On
 : 20/04/2024 11:03 AM

 Age / Sex
 : 31 Year(s) / Male
 Report On
 : 20/04/2024 5:47 PM

 Type
 : OP
 Printed On
 : 22/04/2024 12:31 PM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
TSH (Thyroid Stimulating Hormone)	2.34	μIU/mL	0.35 - 5.50

(Serum Leen 1)

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester: 0.3-3.0

INTERPRETATION:

(Indian Thyroid Society Guidelines)

Comment:

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.

2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.

3. Values&lt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.

PHYSICAL EXAMINATION (URINE COMPLETE)

Colour	Yellow	Yellow to Amber
(Urine)		
Appearance	Clear	Clear
(Urine)		
Volume(CLU)	20	
(Urine)		
CHEMICAL EXAMINATION (<u>URINE</u>	
<u>COMPLETE)</u>		
pН	5	4.5 - 8.0
(Urine)		
Specific Gravity	1.020	1.002 - 1.035
(Urine)		
Ketone	Negative	Negative
(Urine)		
Urobilinogen	Normal	Normal
(Urine)		
Blood	Negative	Negative
(Urine)		





APPROVED BY

The results pertain to sample tested.

Page 7 of 10

 PID No.
 : MED112152459
 Register On
 : 20/04/2024 10:38 AM

 SID No.
 : 424023746
 Collection On
 : 20/04/2024 11:03 AM

 Age / Sex
 : 31 Year(s) / Male
 Report On
 : 20/04/2024 5:47 PM

 Type
 : OP
 Printed On
 : 22/04/2024 12:31 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Unit</u> <u>Value</u>	<u>Biological</u> <u>Reference Interval</u>
Nitrite	Negative	Negative

(Urine)

Bilirubin Negative Negative

(Urine)

Protein Negative Negative

(Urine)

Glucose Negative Negative

(Urine/GOD - POD)

Leukocytes(CP) Negative

(Urine)

MICROSCOPIC EXAMINATION (URINE COMPLETE)

Pus Cells 0-1 /hpf NIL

(Urine)

Epithelial Cells 0-1 /hpf NIL

(Urine)

RBCs NIL /HPF NIL

(Urine)

Others

(Urine)

INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.

Casts NIL /hpf NIL

(Urine)

Crystals NIL /hpf NIL

(Urine)





 PID No.
 : MED112152459
 Register On
 : 20/04/2024 10:38 AM

 SID No.
 : 424023746
 Collection On
 : 20/04/2024 11:03 AM

 Age / Sex
 : 31 Year(s) / Male
 Report On
 : 20/04/2024 5:47 PM

 Type
 : OP
 Printed On
 : 22/04/2024 12:31 PM

Ref. Dr : MediWheel

InvestigationObserved ValueUnit ValueBiological Reference IntervalBUN / Creatinine Ratio6.666.0 - 22.0





 PID No.
 : MED112152459
 Register On
 : 20/04/2024 10:38 AM

 SID No.
 : 424023746
 Collection On
 : 20/04/2024 11:03 AM

 Age / Sex
 : 31 Year(s) / Male
 Report On
 : 20/04/2024 5:47 PM

 Type
 : OP
 Printed On
 : 22/04/2024 12:31 PM

Ref. Dr : MediWheel

InvestigationObservedUnitBiologicalValueReference Interval

URINE ROUTINE





APPROVED BY

-- End of Report --

Name	MR.VIKAS KUMAR SRIVASTAVA	ID	MED112152459
Age & Gender	31Y/MALE	Visit Date	20 Apr 2024
Ref Doctor Name	MediWheel		

2 D ECHOCARDIOGRAPHIC STUDY

M mode measurement:

AORTA : 3.2cms

LEFT ATRIUM : 3.3cms

AVS :----

LEFT VENTRICLE (DIASTOLE) : 4.0cms

(SYSTOLE) : 2.6cms

VENTRICULAR SEPTUM (DIASTOLE) : 0.9cms

(SYSTOLE) : 1.5cms

POSTERIOR WALL (DIASTOLE) : 0.9cms

(SYSTOLE) : 1.6cms

: 35%

EDV : 71ml

ESV : 25ml

EJECTION FRACTION : 65%

EPSS :---

RVID : 1.9cms

DOPPLER MEASUREMENTS:

FRACTIONAL SHORTENING

MITRAL VALVE : E' 0.94 m/s A' 0.63 m/s NO MR

AORTIC VALVE : 0.99 m/s NO AR

TRICUSPID VALVE : E' - m/s A' - m/s NO TR

PULMONARY VALVE : 1.06 m/s NO PR

Name	MR.VIKAS KUMAR SRIVASTAVA	ID	MED112152459
Age & Gender	31Y/MALE	Visit Date	20 Apr 2024
Ref Doctor Name	MediWheel	-	

2D ECHOCARDIOGRAPHY FINDINGS:

Left ventricle : Normal size, Normal systolic function.

No regional wall motion abnormalities.

Left Atrium : Normal.

Right Ventricle : Normal.

Right Atrium : Normal.

Mitral valve : Normal, No mitral valve prolapsed.

Aortic valve : Normal, Trileaflet.

Tricuspid valve : Normal.

Pulmonary valve : Normal.

IAS : Intact.

IVS : Intact.

Pericardium : No pericardial effusion.

IMPRESSION:

- > NORMAL SIZED CARDIAC CHAMBERS.
- > NORMAL LV SYSTOLIC FUNCTION. EF:65 %.
- > NO REGIONAL WALL MOTION ABNORMALITIES.
- > NORMAL VALVES.
- > NO CLOTS / PERICARDIAL EFFUSION / VEGETATION.

DR. K.S. SUBRAMANI. MBBS, MD, DM (CARDIOLOGY) FESC SENIOR CONSULTANT INTERVENTIONAL CARDIOLOGIST SRI JAYADEVA INSTITUTE OF CARDIOLOGY, BANGALORE

Kss/da

Note:

- * Report to be interpreted by qualified medical professional.
- * To be correlated with other clinical findings.
- * Parameters may be subjected to inter and intra observer variations.
- *Any discrepancy in reports due to typing errors should be corrected as soon as possible.

Name	MR.VIKAS KUMAR SRIVASTAVA	ID	MED112152459
Age & Gender	31Y/MALE	Visit Date	20 Apr 2024
Ref Doctor Name	MediWheel	-	

Name	MR.VIKAS KUMAR SRIVASTAVA	ID	MED112152459
Age & Gender	31Y/MALE	Visit Date	20 Apr 2024
Ref Doctor Name	MediWheel		

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in shape, size and has uniform echopattern. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER shows normal shape and has clear contents. Gall bladder wall is of normal thickness. CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN shows normal shape, size and echopattern.

No demonstrable Para -aortic lymphadenopathy.

KIDNEYS move well with respiration and have normal shape, size and echopattern. Cortico- medullary differentiations are well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

·	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney	10.5	1.2
Left Kidney	9.0	1.2

URINARY BLADDER shows normal shape and wall thickness. It has clear contents. No evidence of diverticula.

PROSTATE shows normal shape, size and echopattern. It measures 2.6 x 2.5 x 3.1cms (Vol:11cc).

No evidence of ascites / pleural effusion.

IMPRESSION:

> NO SIGNIFICANT ABNORMALITY DETECTED.

DR. APARNA CONSULTANT RADIOLOGIST A/da

Name	MR.VIKAS KUMAR SRIVASTAVA	ID	MED112152459
Age & Gender	31Y/MALE	Visit Date	20 Apr 2024
Ref Doctor Name	MediWheel		

Name	Mr. Vikas Kumar Srivastava	Customer ID	MED112152459
Age & Gender	31Y/M	Visit Date	Apr 20 2024 10:37AM
Ref Doctor	MediWheel		

X - RAY CHEST PA VIEW

FINDINGS:

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.


Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

IMPRESSION:

No significant abnormality detected.

Dr.Nitash Prakash MBBS.,MD Consultant Radiologist



SRI PARVATHI OPTICS Multi Branded Opticals Store

Computerized Eye Testing & Spectacles Clinic

#333. 8th Main 5th Cross Near Combridge & Miranda School, HAL 3rd Stage, Behind Vishveshvarioh Park New Thipposondro, Bangolore - 560075 email: parvathiopticals@gmail.com

SPECTACLE PRESCRIPTION

Name: Vileas Kungo Solvestale?

Mobil No:

Age / Gender 314/H .

Date: 20 4/2009 Ref. No. 11 2/52454

	RIGHT EYE					LE	FT EY	E
	SPH	CYL	AXIS	VISION	SPH	CYL	AXIS	VISION
	6.50	-	120	014		-	In	0/0
DESTRUCE			- 61	9/0		7.0	18	0/60

PD GURLAP

Advice to use glasses for:

☐ DISTANCE ☐ FAR & NEAR ☐ READING ☐ COMPUTER PURPOSE

We Care Your Eyes

SRI PARVATHI OPTICS NEW THEFASANDRA



CLUMAX DIAGNOSTICS

MEDALL HEALTHCARE PUT LTD
CUSTOMER CHECKLIST



(HARLO)

Com	pany Name	: MediWheel		20/0	04/2024
Pack	age Name :	Mediwheel Full Body Health Checku Study			
S.N	o Modality	Study Study	p Male Above 40		
1	LAB	BLOOD UREA NITROGEN (BUN)	AccessionNo	Time	Signatur
2	LAB	CREATININE (BUN)			- grintar
3	LAB	GLUCOSE - FASTING	-		
4	LAB	GLUCOSE - POSTPRANDIAL (2 HRS)			
5	LAB	GLYCOSYLATED HAEMOGLOBIN (HbA1c)			
6	LAB	URIC ACID		1	
7	LAB	LIPID PROFILE			
8	LAB	LIVER FUNCTION TEST (LFT)			
9	LAB	TOTAL PROSTATE SPECIFIC ANTIGEN			
10	LAB				
11	LAB	THYROID PROFILE/ TFT(T3, T4, TSH) URINE GLUCOSE - FASTING			-
12	LAB	URINE GLUCOSE - POSTPRANDIAL (2 Hrs)			
13	LAB	COMPLETE BLOOD COUNT WITH ESR			1
14	LAB	STOOL ANALYSIS - ROUTINE			
15	LAB	URINE ROUTINE			
16	LAB	BUN/CREATININE RATIO			
17	LAB	BLOOD GROUP & RH TYPE (Forward Reverse)			
18	ECG	ECG .		7	
	OTHERS	Treadmill / 2D Echo	IND14613421138		
	OTHERS	physical examination	IND146134214690		6
1	US	ULTRASOUND ABDOMEN	IND146134215279		ttom
	OTHERS	Dental Consultation	IND146134215292		
3	OTHERS	EYE CHECKUP	IND146134216289		-
	GRAY	X RAY CHECT	IND146134217756		
5 (THERS	Oppultation Division	IND146134218659	_	
-		riysicián	IND146134218736		

