



CODE/NAME & ADDRESS: C000138383 ACCESSION NO: 0080XC008158 AGE/SEX :37 Years Female

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

**NEW DELHI 110030** 

8800465156

PATIENT ID : PURIF17018780

CLIENT PATIENT ID: ABHA NO

DRAWN

RECEIVED: 23/03/2024 08:16:00

REPORTED :23/03/2024 18:35:51

%

fL

pg

g/dL

%

fL

**Test Report Status** Results Biological Reference Interval Units <u>Final</u>

**HAEMATOLOGY - CBC** 

MEDI WHEEL FULL BODY HEALTH CI	HECKUP BELOW 40FEMALE		
BLOOD COUNTS, EDTA WHOLE BLOO	D		
HEMOCLOPIN (HP)	10.0 Low	12.0 15.0	a/dl

HEMOGRODIN (HD)	10.9 LOW	12.0 - 15.0	g/uL
METHOD: CYANMETHEMOGLOBIN METHOD			
RED BLOOD CELL (RBC) COUNT	4.48	3.8 - 4.8	mil/μL
METHOD: ELECTRICAL IMPEDANCE			
WHITE BLOOD CELL (WBC) COUNT	7.39	4.0 - 10.0	thou/µL
METHOD : ELECTRICAL IMPEDANCE			
PLATELET COUNT	296	150 - 410	thou/µL

METHOD: ELECTRICAL IMPEDANCE

RBC AND PLATELET INDICES	

HEMATOCRIT (PCV)	36.6	36 - 46
METHOD: ELECTRICAL IMPEDANCE		
MEAN CORPUSCULAR VOLUME (MCV)	81.6 Low	83 - 101
METHOD: CALCULATED PARAMETER		
MEAN CORPUSCULAR HEMOGLOBIN (MCH)	24.4 Low	27.0 - 32.0
METHOD: CALCULATED PARAMETER		

29.8 Low MEAN CORPUSCULAR HEMOGLOBIN 31.5 - 34.5 CONCENTRATION (MCHC)

METHOD: CALCULATED PARAMETER RED CELL DISTRIBUTION WIDTH (RDW) 15.9 High 11.6 - 14.0

METHOD: CALCULATED PARAMETER MENTZER INDEX 18.2

6.8 - 10.9 13.0 High MEAN PLATELET VOLUME (MPV) METHOD: CALCULATED PARAMETER

# **WBC DIFFERENTIAL COUNT**

NEUTROPHILS	61	40 - 80	%
METHOD: LIGHT ABSORBANCE OF CYTOCHEMICAL S	STAINED CELLS IMPEDANCE		
LYMPHOCYTES	30	20 - 40	%
METHOD: LIGHT ABSORBANCE OF CYTOCHEMICAL S	STAINED CELLS IMPEDANCE		
MONOCYTES	7	2 - 10	%

(Desaraht

Chardni garg

**DR.CHANDNI GARG** 

**CONSULTANT PATHOLOGIST** 



Page 1 Of 19





Dr.Pranjali Vasisht

**LAB HEAD** 

Agilus Diagnostics Ltd. 24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India







**PATIENT NAME: PURI HEENA REF. DOCTOR: SELF** 

CODE/NAME & ADDRESS: C000138383 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

**NEW DELHI 110030** 

8800465156

ACCESSION NO: 0080XC008158

PATIENT ID : PURIF17018780

CLIENT PATIENT ID: ABHA NO

AGE/SEX :37 Years

RECEIVED: 23/03/2024 08:16:00 REPORTED :23/03/2024 18:35:51

Test Report Status <u>Final</u>	Results	Biological Reference	Interval Units
METIOD LYGUE ADGODDANGE OF GEOGRAPHICAL STANKED OF LA	C MADED ANGE		
METHOD: LIGHT ABSORBANCE OF CYTOCHEMICAL STAINED CELL EOSINOPHILS	S IMPEDANCE  7	1 - 6	%
	2		
BASOPHILS	0	0 - 2	%
METHOD: LIGHT ABSORBANCE OF CYTOCHEMICAL STAINED CELL	S IMPEDANCE		
ABSOLUTE NEUTROPHIL COUNT	4.51	2.0 - 7.0	thou/µL
ABSOLUTE LYMPHOCYTE COUNT	2.22	1 - 3	thou/µL
ABSOLUTE MONOCYTE COUNT	0.52	0.20 - 1.00	thou/µL
METHOD: CALCULATED PARAMETER			
ABSOLUTE EOSINOPHIL COUNT	0.15	0.02 - 0.50	thou/µL
ABSOLUTE BASOPHIL COUNT	0.00 Low	0.02 - 0.10	thou/µL
NEUTROPHIL LYMPHOCYTE RATIO (NLR)	2.0		

METHOD: CALCULATED PARAMETER

#### Interpretation(s)

BLOOD COUNTS, EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13)

from Beta thalassaemia trait

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for

diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR = 3.0 years old and N

This ratio element is a calculated parameter and out of NABL scope.

(Desaraht

Dr. Pranjali Vasisht LAB HEAD

Chandri Garg

**DR.CHANDNI GARG CONSULTANT PATHOLOGIST** 





Page 2 Of 19



Agilus Diagnostics Ltd. 24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India







mm at 1 hr

**REF. DOCTOR: SELF PATIENT NAME: PURI HEENA** 

CODE/NAME & ADDRESS: C000138383 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

**NEW DELHI 110030** 

8800465156

ACCESSION NO: 0080XC008158

PATIENT ID : PURIF17018780

CLIENT PATIENT ID: ABHA NO

AGE/SEX

RECEIVED: 23/03/2024 08:16:00 REPORTED :23/03/2024 18:35:51

:37 Years

**Test Report Status** Results **Biological Reference Interval Final** Units

**HAEMATOLOGY** 

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

**ERYTHROCYTE SEDIMENTATION RATE (ESR), EDTA BLOOD** 

E.S.R 40 High 0 - 20

METHOD: MODIFIED WESTERGREN

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE **BLOOD** 

Non-diabetic Adult < 5.7 HBA1C 4.9 %

Pre-diabetes 5.7 - 6.4

Diabetes diagnosis: > or = 6.5Therapeutic goals: < 7.0 Action suggested : > 8.0

(ADA Guideline 2021)

ESTIMATED AVERAGE GLUCOSE(EAG) 93.9 < 116.0 mg/dL

Interpretation(s)

ERYTHROCYTE SEDIMENTATION RATE (ESR),EDTA BLOOD-TEST DESCRIPTION :-

Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change. **TEST INTERPRETATION** 

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy,

Estrogen medication, Aging.
Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias,

Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis).

In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum. Decreased in: Polycythermia vera, Sickle cell anemia

LIMITATIONS

False elevated ESR: Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia

False Decreased: Poikilocytosis, (SickleCells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs (Quinine,

salicylates)

REFERENCE:

Chardni Garg DR.CHANDNI GARG

(Desaraht

Dr.Pranjali Vasisht LAB HEAD

Page 3 Of 19

CONSULTANT PATHOLOGIST







Agilus Diagnostics Ltd. 24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India Tel: 9111591115, Fax:

CIN - U74899PB1995PLC045956







**REF. DOCTOR: SELF PATIENT NAME: PURI HEENA** 

CODE/NAME & ADDRESS: C000138383 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

**NEW DELHI 110030** 

8800465156

ACCESSION NO: 0080XC008158

: PURIF17018780

AGE/SEX :37 Years Female

RECEIVED: 23/03/2024 08:16:00

REPORTED :23/03/2024 18:35:51

**Test Report Status** Results **Biological Reference Interval** <u>Final</u> Units

CLIENT PATIENT ID:

PATIENT ID

ABHA NO

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis,10th edition. GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-**Used For**:

- 1. Evaluating the long-term control of blood glucose concentrations in diabetic patients. 2. Diagnosing diabetes.

3. Identifying patients at increased risk for diabetes (prediabetes).

The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for well-controlled type 2 diabetic patients) to determine whether a patients metabolic control has remained continuously within the target range.

- eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.
   eAG gives an evaluation of blood glucose levels for the last couple of months.
- 3. eAG is calculated as eAG (mg/dl) = 28.7 \* HbA1c 46.7

### HbA1c Estimation can get affected due to :

- 1. Shortened Erythrocyte survival: Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss,hemolytic anemia) will falsely lower HbA1c test results. Fructosamine is recommended in these patients which indicates diabetes control over 15 days.
- 2.Vitamin C & E are reported to falsely lower test results. (possibly by inhibiting glycation of hemoglobin.

  3. Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates addiction are reported to interfere with some assay methods, falsely increasing results.
- 4. Interference of hemoglobinopathies in HbA1c estimation is seen in
- a) Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c. b) Heterozygous state detected (D10 is corrected for HbS & HbC trait.)
- c) HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c. Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy

Chardni Garg

DR.CHANDNI GARG CONSULTANT PATHOLOGIST Dr. Pranjali Vasisht LAB HEAD

(Poraralit

Page 4 Of 19



24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956



Agilus Diagnostics Ltd.





CODE/NAME & ADDRESS: C000138383 ACCESSION NO: 0080XC008158 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID

F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI

**NEW DELHI 110030** 8800465156

: PURIF17018780

CLIENT PATIENT ID: ABHA NO

AGE/SEX :37 Years Female

RECEIVED: 23/03/2024 08:16:00 REPORTED :23/03/2024 18:35:51

**Test Report Status** Results **Biological Reference Interval** Units <u>Final</u>

#### **IMMUNOHAEMATOLOGY**

### MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

**ABO GROUP** TYPE O

METHOD : SLIDE AGGLUTINATION

RH TYPE **NEGATIVE** 

METHOD: SLIDE AGGLUTINATION

Interpretation(s)

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same.

The test is performed by both forward as well as reverse grouping methods.

Coracult

Dr. Pranjali Vasisht **LAB HEAD** 

Chandri Garg

**DR.CHANDNI GARG CONSULTANT PATHOLOGIST** 





Page 5 Of 19



Agilus Diagnostics Ltd. 24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India







**PATIENT NAME: PURI HEENA REF. DOCTOR: SELF** 

CODE/NAME & ADDRESS : C000138383 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

**NEW DELHI 110030** 8800465156

ACCESSION NO: 0080XC008158

PATIENT ID : PURIF17018780

CLIENT PATIENT ID: ABHA NO

AGE/SEX :37 Years

RECEIVED: 23/03/2024 08:16:00

REPORTED :23/03/2024 18:35:51

**Test Report Status** Results Biological Reference Interval Units <u>Final</u>

**BIOCHEMISTRY** 

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

GLUCOSE FASTING, FLUORIDE PLASMA

FBS (FASTING BLOOD SUGAR)

91

74 - 106

mg/dL

**GLUCOSE, POST-PRANDIAL, PLASMA** 

PPBS(POST PRANDIAL BLOOD SUGAR)

91

Non-Diabetes 70 - 140

mg/dL

METHOD: HEXOKINASE

METHOD: HEXOKINASE

LIPID PROFILE WITH CALCULATED LDL, SERUM

CHOLESTEROL, TOTAL 168 < 200 Desirable

200 - 239 Borderline High

mg/dL

METHOD: CHOLESTEROL OXIDASE, ESTERASE, PEROXIDASE

TRIGLYCERIDES

93

< 150 Normal

>/= 240 High

mg/dL

150 - 199 Borderline High

200 - 499 High >/= 500 Very High

METHOD: ENZYMATIC ASSAY

METHOD: DIRECT MEASURE - PEG CHOLESTEROL LDL

HDL CHOLESTEROL 49 < 40 Low >/=60 High mg/dL

mg/dL

100

< 100 Optimal

100 - 129

Near or above optimal

130 - 159 Borderline High 160 - 189

High >/= 190 Very High

METHOD: CHOLESTEROL OXIDASE, ESTERASE, PEROXIDASE

Chardni garg

Parallet

Dr.Pranjali Vasisht LAB HEAD

Page 6 Of 19

DR.CHANDNI GARG CONSULTANT PATHOLOGIST







Agilus Diagnostics Ltd. 24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India



8800465156





PATIENT NAME: PURI HEENA REF. DOCTOR: SELF

CODE/NAME & ADDRESS : C000138383 ACCESSION NO : **0080XC008158** AGE/SEX : 37 Years Female

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID : PURIF17018780

F-703, LADO SARAI, MEHRAULISOUTH WEST CLIENT PATIENT ID: RECEIVED : 23/03/2024 08:16:00

DELHI NEW DELHI 110030 | CLIENT PATIENT ID: | RECEIVED : 23/03/2024 08:16:00 | REPORTED : 23/03/2024 18:35:51

	<u> </u>	<u> </u>
Test Report Status <u>Final</u>	Results	Biological Reference Interval Units
NON HDL CHOLESTEROL	49	Desirable: Less than 130 mg/dL Above Desirable: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very high: > or = 220
METHOD: CALCULATED PARAMETER		
VERY LOW DENSITY LIPOPROTEIN	18.6	Desirable value : mg/dL 10 - 35
METHOD: CALCULATED PARAMETER		
CHOL/HDL RATIO	3.4	3.3-4.4 Low Risk 4.5-7.0 Average Risk 7.1-11.0 Moderate Risk > 11.0 High Risk
METHOD: CALCULATED PARAMETER		
LDL/HDL RATIO	2.0	0.5 - 3.0 Desirable/Low Risk 3.1 - 6.0 Borderline/Moderate Risk >6.0 High Risk
METHOD: CALCULATED PARAMETER		

### Interpretation(s)

Serum lipid profile is measured for cardiovascular risk prediction. Lipid Association of India recommends LDL-C as primary target and Non HDL-C as co-primary treatment target.

Risk Stratification for ASCVD (Atherosclerotic cardiovascular disease) by Lipid Association of India

Risk Category			
Extreme risk group	A.CAD with > 1 feature of high risk group		
	B. CAD with > 1 feature of Very high risk g	roup or recurrent ACS (within 1 year) despite LDL-C < or =	
	50 mg/dl or polyvascular disease		
Very High Risk	1. Established ASCVD 2. Diabetes with 2 r	najor risk factors or evidence of end organ damage 3.	
	Familial Homozygous Hypercholesterolemia	a	
High Risk		betes with 1 major risk factor or no evidence of end organ	
	damage. 3. CKD stage 3B or 4. 4. LDL >190 mg/dl 5. Extreme of a single risk factor. 6. Coronary		
	Artery Calcium - CAC >300 AU. 7. Lipoprotein a >/= 50mg/dl 8. Non stenotic carotid plaque		
Moderate Risk	2 major ASCVD risk factors		
Low Risk	0-1 major ASCVD risk factors		
Major ASCVD (Athe	Major ASCVD (Atherosclerotic cardiovascular disease) Risk Factors		
1. Age $>$ or $=$ 45 years	1. Age > or = 45 years in males and > or = 55 years in females  3. Current Cigarette smoking or tobacco use		
2. Family history of p	2. Family history of premature ASCVD 4. High blood pressure		
5. Low HDL			

Chardni Garg

Coraculit

DR.CHANDNI GARG
CONSULTANT PATHOLOGIST

Dr.Pranjali Vasisht LAB HEAD





Page 7 Of 19

View Details





Agilus Diagnostics Ltd. 24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India







CODE/NAME & ADDRESS : C000138383 ACCESSION NO: 0080XC008158 AGE/SEX :37 Years Female

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID : PURIF17018780

DRAWN F-703, LADO SARAI, MEHRAULISOUTH WEST

CLIENT PATIENT ID: RECEIVED: 23/03/2024 08:16:00 DELHI ABHA NO REPORTED :23/03/2024 18:35:51 **NEW DELHI 110030** 8800465156

**Test Report Status Results Biological Reference Interval** Units <u>Final</u>

Newer treatment goals and statin initiation thresholds based on the risk categories proposed by LAI in 2020

Risk Group	Treatment Goals		Consider Drug Therapy	
	LDL-C (mg/dl)	Non-HDL (mg/dl)	LDL-C (mg/dl)	Non-HDL (mg/dl)
Extreme Risk Group Category A	<50 (Optional goal < OR = 30 )	< 80 (Optional goal <or 60)<="" =="" td=""><td>&gt;OR = 50</td><td>&gt;OR = 80</td></or>	>OR = 50	>OR = 80
Extreme Risk Group Category B	<or 30<="" =="" td=""><td><or 60<="" =="" td=""><td>&gt; 30</td><td>&gt;60</td></or></td></or>	<or 60<="" =="" td=""><td>&gt; 30</td><td>&gt;60</td></or>	> 30	>60
Very High Risk	<50	<80	>OR= 50	>OR= 80
High Risk	<70	<100	>OR= 70	>OR= 100
Moderate Risk	<100	<130	>OR= 100	>OR= 130
Low Risk	<100	<130	>OR= 130*	>OR= 160

<sup>\*</sup>After an adequate non-pharmacological intervention for at least 3 months.

References: Management of Dyslipidaemia for the Prevention of Stroke: Clinical Practice Recommendations from the Lipid Association of India. Current Vascular Pharmacology, 2022, 20, 134-155.

### LIVER FUNCTION PROFILE, SERUM

· · · · · · · · · · · · · · · · · · ·			
BILIRUBIN, TOTAL	0.48	UPTO 1.2	mg/dL
METHOD: DIAZONIUM ION, BLANKED (ROCHE) BILIRUBIN, DIRECT	0.16	0.00 - 0.30	mg/dL
METHOD : DIAZOTIZATION			-
BILIRUBIN, INDIRECT  METHOD: CALCULATED PARAMETER	0.32	0.00 - 0.60	mg/dL
TOTAL PROTEIN	7.7	6.6 - 8.7	g/dL
METHOD : BIURET			
ALBUMIN METHOD: BROMOCRESOL GREEN	4.3	3.97 - 4.94	g/dL
GLOBULIN	3.4	2.0 - 4.0 Neonates - Pre Mature: 0.29 - 1.04	g/dL
METHOD: CALCULATED PARAMETER			
ALBUMIN/GLOBULIN RATIO  METHOD: CALCULATED PARAMETER	1.3	1.0 - 2.0	RATIO
ASPARTATE AMINOTRANSFERASE(AST/SGOT)	28	0 - 32	U/L
ALANINE AMINOTRANSFERASE (ALT/SGPT)  METHOD: UV WITHOUT PYRIDOXAL-5 PHOSPHATE	37 High	0 - 31	U/L
ALKALINE PHOSPHATASE  METHOD: PNPP - AMP BUFFER	142 High	35 - 105	U/L
GAMMA GLUTAMYL TRANSFERASE (GGT)  METHOD: GAMMA GLUTAMYLCARBOXY 4NITROANILIDE	26	5 - 36	U/L
LACTATE DEHYDROGENASE  METHOD: LACTATE -PYRUVATE	166	135 - 214	U/L

METHOD: LACTATE -PYRUVATE

Chardni garg **DR.CHANDNI GARG** 

Dr.Pranjali Vasisht

Coracalit

**LAB HEAD** 





Page 8 Of 19



Agilus Diagnostics Ltd. 24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

**CONSULTANT PATHOLOGIST** 







**PATIENT NAME: PURI HEENA REF. DOCTOR: SELF** 

CODE/NAME & ADDRESS: C000138383 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

**NEW DELHI 110030** 8800465156

ACCESSION NO: 0080XC008158

PATIENT ID : PURIF17018780

CLIENT PATIENT ID: ABHA NO

AGE/SEX

RECEIVED: 23/03/2024 08:16:00 REPORTED :23/03/2024 18:35:51

:37 Years

**Test Report Status** Results **Biological Reference Interval** Units <u>Final</u>

**BLOOD UREA NITROGEN (BUN), SERUM** 

**BLOOD UREA NITROGEN** 9 6 - 20mg/dL

METHOD: UREASE - UV

**CREATININE, SERUM** 

CREATININE 0.79 0.50 - 0.90mg/dL

METHOD: ALKALINE PICRATE-KINETIC

**BUN/CREAT RATIO** 

5.00 - 15.00 BUN/CREAT RATIO 11.39

METHOD: CALCULATED PARAMETER

**URIC ACID, SERUM** 

URIC ACID 5.1 2.4 - 5.7mg/dL

METHOD: URICASE, COLORIMETRIC

**TOTAL PROTEIN, SERUM** 

TOTAL PROTEIN 7.7 6.6 - 8.7g/dL

METHOD : BIURET

**ALBUMIN, SERUM** 

4.3 3.97 - 4.94g/dL ALBUMIN

METHOD: BROMOCRESOL GREEN

**DR.CHANDNI GARG CONSULTANT PATHOLOGIST**  Dr.Pranjali Vasisht LAB HEAD

Coraculit





Page 9 Of 19



Chardni garg

Agilus Diagnostics Ltd. 24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India







CODE/NAME & ADDRESS: C000138383 ACCESSION NO: 0080XC008158 AGE/SEX :37 Years Female

: PURIF17018780

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID F-703, LADO SARAI, MEHRAULISOUTH WEST

CLIENT PATIENT ID: DELHÍ

**NEW DELHI 110030** 8800465156

ABHA NO

RECEIVED: 23/03/2024 08:16:00 REPORTED :23/03/2024 18:35:51

Test Report Status <u>Final</u>	Results	Biological Reference	Interval Units
GLOBULIN			
GLOBULIN	3.4	2.0 - 4.0 Neonates - Pre Mature: 0.29 - 1.04	g/dL
METHOD: CALCULATED PARAMETER			
ELECTROLYTES (NA/K/CL), SERUI	М		
SODIUM, SERUM	142	136 - 145	mmol/L
METHOD: ISE INDIRECT			
POTASSIUM, SERUM METHOD: ISE INDIRECT	4.38	3.5 - 5.1	mmol/L
CHLORIDE, SERUM METHOD: ISE INDIRECT	105	98 - 107	mmol/L

## Interpretation(s)

Sodium	Potassium	Chloride
Decreased in:CCF, cirrhosis, vomiting, diarrhea, excessive sweating, salt-losing nephropathy, adrenal insufficiency, nephrotic syndrome, water intoxication, SIADH. Drugs: thiazides, diuretics, ACE inhibitors, chlorpropamide, carbamazepine, anti depressants (SSRI), antipsychotics.	Decreased in: Low potassium intake,prolonged vomiting or diarrhea, RTA types I and II, hyperaldosteronism, Cushing's syndrome,osmotic diuresis (e.g., hyperglycemia),alkalosis, familial periodic paralysis,trauma (transient).Drugs: Adrenergic agents, diuretics.	Decreased in: Vomiting, diarrhea, renal failure combined with salt deprivation, over-treatment with diuretics, chronic respiratory acidosis, diabetic ketoacidosis, excessive sweating, SIADH, salt-losing nephropathy, porphyria, expansion of extracellular fluid volume, adrenalinsufficiency, hyperaldosteronism, metabolic alkalosis. Drugs: chronic laxative, corticosteroids, diuretics.
Increased in: Dehydration (excessivesweating, severe vomiting or diarrhea), diabetes mellitus, diabetesinsipidus, hyperaldosteronism, inadequate water intake. Drugs: steroids, licorice, oral contraceptives.	Increased in: Massive hemolysis, severe tissue damage, rhabdomyolysis, acidosis, dehydration, renal failure, Addison's disease, RTA type IV, hyperkalemic familial periodic paralysis. Drugs: potassium salts, potassium- sparing diuretics, NSAIDs, beta-blockers, ACE inhibitors, highdose trimethoprim-sulfamethoxazole.	Increased in: Renal failure, nephrotic syndrome, RTA, dehydration, overtreatment with saline, hyperparathyroidism, diabetes insipidus, metabolic acidosis from diarrhea (Loss of HCO3-), respiratory alkalosis, hyperadrenocorticism.  Drugs: acetazolamide, androgens, hydrochlorothiazide, salicylates.

Chardni garg

**DR.CHANDNI GARG CONSULTANT PATHOLOGIST**  Coraculit

Dr.Pranjali Vasisht **LAB HEAD** 



Page 10 Of 19







Agilus Diagnostics Ltd. 24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India







**REF. DOCTOR: SELF PATIENT NAME: PURI HEENA** 

CODE/NAME & ADDRESS: C000138383 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

**NEW DELHI 110030** 

8800465156

ACCESSION NO : 0080XC008158

PATIENT ID : PURIF17018780

CLIENT PATIENT ID: ABHA NO

AGE/SEX

RECEIVED: 23/03/2024 08:16:00

:37 Years

REPORTED: 23/03/2024 18:35:51

**Test Report Status** Results Biological Reference Interval **Final** Units

Interferences: Severe linemia or hyperproteinemi, if sodium analysis involves a dilution step can cause spurious results. The serum sodium falls about 1.6 mEq/L for each 100 mg/dL increase in blood glucose.

Interferences: Hemolysis of sample. delayed separation of serum, prolonged fist clenching during blood drawing, and prolonged tourniquet placement. Very high WBC/PLT counts may cause spurious. Plasma potassium levels are normal.

Interferences: Test is helpful in assessing normal and increased anion gap metabolic acidosis and in distinguishing hypercalcemia due to hyperparathyroidism (high serum chloride) from that due to malignancy (Normal serum chloride)

#### Interpretation(s)

GLUCOSE FASTING, FLUORIDE PLASMA-TEST DESCRIPTION

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in the

Increased in:Diabetes mellitus, Cushing's syndrome (10 – 15%), chronic pancreatitis (30%). Drugs:corticosteroids, phenytoin, estrogen, thiazides. Decreased in:Pancreatic islet cell disease with increased insulin, insulinoma, adrenocortical insufficiency, hypopituitarism, diffuse liver disease, malignancy(adrenocortical,stomach,fibrosarcoma),infant of a diabetic mother,enzyme deficiency diseases(e.g.galactosemia),Drugs-insulin,ethanol,propranolol

sulfonylureas,tolbutamide,and other oral hypoglycemic agents.

NOTE: While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values),there is wide fluctuation within individuals. Thus, glycosylated hemoglobin(HbA1c) levels are favored to monitor glycemic control.

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic

index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.

GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc. Additional test HbA1c

LIVER FUNCTION PROFILE, SERUM
Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis.

**ALP** is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Pagets disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilsons disease.

**GGT** is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive

liver disease, high alcohol consumption and use of enzyme-inducing drugs etc. **Total Protein** also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease.Lower-than-normal levels may be due to: Agammaglobulinemia,Bleeding (hemorrhage),Burns,Glomerulonephritis,Liver disease, Malabsorption,Malnutrition,Nephrotic syndrome,Protein-losing enteropathy etc.

**Albumin** is the most abundant protein in human blood plasma.It is produced in the liver.Albumin constitutes about half of the blood serum protein.Low blood albumin levels (hypoalbuminemia) can be caused by:Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular

permeability or decreased lymphatic clearance,malnutrition and wasting etc
BLOOD UREA NITROGEN (BUN), SERUM-**Causes of Increased** levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)

Causes of decreased level include Liver disease, SIADH.
CREATININE, SERUM-Higher than normal level may be due to:

• Blockage in the urinary tract, Kidney problems, such as kidney damage or failure, infection, or reduced blood flow, Loss of body fluid (dehydration), Muscle problems, such as breakdown of muscle fibers, Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:• Myasthenia Gravis, Muscuophy

URIC ACID, SERUM-Causes of Increased levels:-Dietary(High Protein Intake,Prolonged Fasting,Rapid weight loss),Gout,Lesch nyhan syndrome,Type 2 DM,Metabolic syndrome Causes of decreased levels:-Low Zinc intake,OCP,Multiple Sclerosis

TOTAL PROTEIN, SERUM-is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin.

Chardni Garg DR.CHANDNI GARG

CONSULTANT PATHOLOGIST

Dr. Pranjali Vasisht LAB HEAD

(Desaraht



Page 11 Of 19



Agilus Diagnostics Ltd. 24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India







PATIENT NAME: PURI HEENA REF. DOCTOR: SELF

CODE/NAME & ADDRESS : C000138383

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL
F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

**NEW DELHI 110030** 

8800465156

ACCESSION NO : 0080XC008158

| |PATIENT ID : PURIF17018780

CLIENT PATIENT ID: ABHA NO : AGE/SEX

RECEIVED : 23/03/2024 08:16:00

:37 Years

REPORTED :23/03/2024 18:35:51

Test Report Status <u>Final</u> Results Biological Reference Interval Units

**Higher-than-normal levels may be due to:** Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease. **Lower-than-normal levels may be due to:** Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

ALBUMIN, SERUM-Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. **Low blood albumin levels (hypoalbuminemia) can be caused by:** Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance,malnutrition and wasting etc.

Chardni Garg

DR.CHANDNI GARG
CONSULTANT PATHOLOGIST

Obrasalit.

Dr.Pranjali Vasisht LAB HEAD





Page 12 Of 19

View Details

View Report



Agilus Diagnostics Ltd. 24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India







CODE/NAME & ADDRESS: C000138383 ACCESSION NO: 0080XC008158 AGE/SEX :37 Years ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI

**NEW DELHI 110030** 

8800465156

PATIENT ID : PURIF17018780

CLIENT PATIENT ID: ABHA NO

Female

RECEIVED: 23/03/2024 08:16:00 REPORTED :23/03/2024 18:35:51

**Test Report Status Biological Reference Interval Units** Results **Final** 

### **CLINICAL PATH - URINALYSIS**

#### MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

PHYSICAL EXAMINATION, URINE

COLOR PALE YELLOW

**APPEARANCE CLEAR** 

### CHEMICAL EXAMINATION, URINE

PH 6.5 4.7 - 7.5

METHOD: REFLECTANCE SPECTROPHOTOMETRY- DOUBLE INDICATOR METHOD 1.005 1.003 - 1.035 SPECIFIC GRAVITY

METHOD: REFLECTANCE SPECTROPHOTOMETRY (PKA CHANGE OF PRETREATED POLY ELECTROLYTES)

NOT DETECTED **PROTEIN** NOT DETECTED

METHOD: REFLECTANCE SPECTROPHOTOMETRY (PROTEIN-ERROR-OF-INDICATORS PRINCIPLE) NOT DETECTED NOT DETECTED GLUCOSE

METHOD: REFLECTANCE SPECTROPHOTOMETRY(GLUCOSE OXIDAE/PEROXIDASE METHOD) NOT DETECTED KFTONES NOT DETECTED

METHOD: REFLECTANCE SPECTROPHOTOMETRY (SODIUM NITROPRUSSIDE REACTION)

NOT DETECTED BLOOD NOT DETECTED

METHOD: REFLECTANCE SPECTROPHOTOMETRY (PEROXIDASE METHOD)

**BILIRUBIN** NOT DETECTED NOT DETECTED

METHOD: REFLECTANCE SPECTROPHOTOMETRY (DIAZO REACTION)

UROBILINOGEN **NORMAL NORMAL** 

METHOD: REFLECTANCE SPECTROPHOTOMETRY - EHRLICH REACTION

NITRITE NOT DETECTED NOT DETECTED

METHOD: REFLECTANCE SPECTROPHOTOMETRY, CONVERSION OF NITRATE TO NITRITE

LEUKOCYTE ESTERASE NOT DETECTED NOT DETECTED

### MICROSCOPIC EXAMINATION, URINE

/HPF **NOT DETECTED** RED BLOOD CELLS NOT DETECTED

METHOD: MICROSCOPIC EXAMINATION

PUS CELL (WBC'S) 1-2 0 - 5/HPF

METHOD: MICROSCOPIC EXAMINATION

1-2 0-5 /HPF EPITHELIAL CELLS

METHOD: MICROSCOPIC EXAMINATION

DR.CHANDNI GARG CONSULTANT PATHOLOGIST Deraralit

Dr.Pranjali Vasisht LAB HEAD



Page 13 Of 19





Chardni garg

Agilus Diagnostics Ltd. 24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India







CODE/NAME & ADDRESS : C000138383 ACCESSION NO: 0080XC008158 AGE/SEX :37 Years Female

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID : PURIF17018780

F-703, LADO SARAI, MEHRAULISOUTH WEST

CLIENT PATIENT ID: RECEIVED: 23/03/2024 08:16:00 DELHI REPORTED :23/03/2024 18:35:51 ABHA NO **NEW DELHI 110030** 

**Test Report Status Final Results Biological Reference Interval** Units

NOT DETECTED **CASTS** NOT DETECTED **CRYSTALS** 

METHOD: MICROSCOPIC EXAMINATION

**BACTERIA** NOT DETECTED NOT DETECTED

METHOD: MICROSCOPIC EXAMINATION

YEAST NOT DETECTED NOT DETECTED

### Interpretation(s)

8800465156

The following table describes the probable conditions, in which the analytes are present in urine

Presence of	Conditions					
Proteins	Inflammation or immune illnesses					
Pus (White Blood Cells)	Urinary tract infection, urinary tract or kidney stone, tumors or any kind					
	of kidney impairment					
Glucose	Diabetes or kidney disease					
Ketones	Diabetic ketoacidosis (DKA), starvation or thirst					
Urobilinogen	Liver disease such as hepatitis or cirrhosis					
Blood	Renal or genital disorders/trauma					
Bilirubin	Liver disease					
Erythrocytes	Urological diseases (e.g. kidney and bladder cancer, urolithiasis), urinary tract infection and glomerular diseases					
Leukocytes	Urinary tract infection, glomerulonephritis, interstitial nephritis either acute or chronic, polycystic kidney disease, urolithiasis, contamination by genital secretions					
Epithelial cells	Urolithiasis, bladder carcinoma or hydronephrosis, ureteric stents or bladder catheters for prolonged periods of time					
Granular Casts	Low intratubular pH, high urine osmolality and sodium concentration, interaction with Bence-Jones protein					
Hyaline casts	Physical stress, fever, dehydration, acute congestive heart failure, renal diseases					
Calcium oxalate	Metabolic stone disease, primary or secondary hyperoxaluria, intravenous infusion of large doses of vitamin C, the use of vasodilator naftidrofuryl oxalate or the gastrointestinal lipase inhibitor orlistat, ingestion of ethylene glycol or of star fruit (Averrhoa carambola) or its juice					
Uric acid	arthritis					

Chardni garg

**DR.CHANDNI GARG** 

Dr.Pranjali Vasisht

Coracalit

**LAB HEAD** 





Page 14 Of 19





Agilus Diagnostics Ltd. 24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

**CONSULTANT PATHOLOGIST** 







PATIENT NAME: PURI HEENA REF. DOCTOR: SELF

CODE/NAME & ADDRESS: C000138383

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL
F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHÍ

NEW DELHI 110030 8800465156 ACCESSION NO: 0080XC008158

PATIENT ID : PURIF17018780

CLIENT PATIENT ID: ABHA NO : AGE/SEX :37 Years

RECEIVED : 23/03/2024 08:16:00

REPORTED : 23/03/2024 18:35:51

Test Report Status Final Results Biological Reference Interval Units

Bacteria	Urinary infectionwhen present in significant numbers & with pus cells.					
Trichomonas vaginalis	Vaginitis, cervicitis or salpingitis					

Chardni Garg

DR.CHANDNI GARG
CONSULTANT PATHOLOGIST

Coraculit

Dr.Pranjali Vasisht LAB HEAD





Page 15 Of 19

View Details





Agilus Diagnostics Ltd. 24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India







CODE/NAME & ADDRESS: C000138383 ACCESSION NO: 0080XC008158 AGE/SEX :37 Years Female ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

**NEW DELHI 110030** 8800465156

PATIENT ID : PURIF17018780

CLIENT PATIENT ID: ABHA NO

DRAWN

RECEIVED: 23/03/2024 08:16:00

REPORTED :23/03/2024 18:35:51

**Test Report Status** Results **Biological Reference Interval** Units <u>Final</u>

#### **CYTOLOGY**

### MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

#### **PAPANICOLAOU SMEAR**

TEST METHOD CONVENTIONAL GYNEC CYTOLOGY

TWO UNSTAINED CERVICAL SMEARS RECEIVED SPECIMEN TYPE

2014 BETHESDA SYSTEM FOR REPORTING CERVICAL CYTOLOGY REPORTING SYSTEM

**SATISFACTORY** SPECIMEN ADEQUACY

SMEARS SHOW ADEQUATE CELLULARITY COMPOSED PREDOMINANTLY **MICROSCOPY** 

OF INTERMEDIATE SQUAMOUS EPITHELIAL CELLS ALONG WITH FEW SUPERFICIAL SQUAMOUS EPITHELIAL CELLS IN A BACKGROUND OF POLYMORPHS AND BLOOD.ENDOCERVICAL CELLS SEEN.NO EVIDENCE

OF MALIGNANCY SEEN.

NEGATIVE FOR INTRAEPITHELIAL LESION OR MALIGNANCY INTERPRETATION / RESULT

REACTIVE CELLULAR CHANGES ASSOCIATED WITH INFLAMMATION.

Chardni Garg

DR.CHANDNI GARG CONSULTANT PATHOLOGIST (Poraralit

Dr.Pranjali Vasisht LAB HEAD





Page 16 Of 19



Agilus Diagnostics Ltd. 24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India





CODE/NAME & ADDRESS: C000138383 ACCESSION NO: 0080XC008158 AGE/SEX :37 Years Female ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

PATIENT ID : PURIF17018780

F-703, LADO SARAI, MEHRAULISOUTH WEST CLIENT PATIENT ID: DELHÍ

RECEIVED: 23/03/2024 08:16:00 REPORTED :23/03/2024 18:35:51 ABHA NO **NEW DELHI 110030** 8800465156

**Biological Reference Interval Test Report Status** Results Units <u>Final</u>

### MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

**LETTER** 

REQUEST LETTER CX/34/2024

Chardni garg

**DR.CHANDNI GARG CONSULTANT PATHOLOGIST**  Coraculit

Dr.Pranjali Vasisht **LAB HEAD** 





Page 17 Of 19





Agilus Diagnostics Ltd. 24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India







F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI
NEW DELHI 110030

CLIENT PATIENT ID:
ABHA NO :

CLIENT PATIENT ID: RECEIVED : 23/03/2024 08:16:00
ABHA NO : REPORTED : 23/03/2024 18:35:51

Test Report Status <u>Final</u> Results Biological Reference Interval Units

#### **SPECIALISED CHEMISTRY - HORMONE**

# MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

### THYROID PANEL, SERUM

8800465156

T3 137.80 80.00 - 200.00 ng/dL

METHOD : COMPETITIVE (ECLIA)

T4 7.58 5.10 - 14.10  $\mu g/dL$ 

METHOD: COMPETITIVE (ECLIA)

TSH (ULTRASENSITIVE) 4.010 Non Pregnant Women µIU/mL

0.27 - 4.20

Pregnant Women (As per American Thyroid Association) 1st Trimester 0.100 - 2.500 2nd Trimester 0.200 - 3.000 3rd Trimester 0.300 - 3.000

METHOD: SANDWICH (ECLIA)

### Interpretation(s)

Triiodothyronine T3, Thyroxine T4, and Thyroid Stimulating Hormone TSH are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hyperthyroidism, TSH levels are low. Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3.Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Sr. No.	TSH	Total T4	FT4	Total T3	Possible Conditions
1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3)
					Post Thyroidectomy (4) Post Radio-Iodine treatment
2	High	Normal	Normal	Normal	(1)Subclinical Hypothyroidism (2) Patient with insufficient thyroid
					hormone replacement therapy (3) In cases of Autoimmune/Hashimoto
					thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical
					inflammation, drugs like amphetamines, Iodine containing drug and
					dopamine antagonist e.g. domperidone and other physiological reasons.
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism

Chardni Garg

Caralit

Dr.Pranjali Vasisht LAB HEAD Page 18 Of 19

DR.CHANDNI GARG
CONSULTANT PATHOLOGIST





View Details

View Repor



Agilus Diagnostics Ltd. 24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India Tel: 9111591115, Fax:

CIN - U74899PB1995PLC045956







 CODE/NAME & ADDRESS : C000138383
 ACCESSION NO : 0080XC008158
 AGE/SEX : 37 Years
 Female

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHÍ

NEW DELHI 110030

8800465156

DATIENT ID . DUDIES 701 0700

PATIENT ID : PURIF17018780

CLIENT PATIENT ID: ABHA NO : DRAWN :

RECEIVED : 23/03/2024 08:16:00

REPORTED :23/03/2024 18:35:51

Test Report Status <u>Final</u> Results Biological Reference Interval Units

4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre
					(3)Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid
					hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4
					replacement therapy (7) First trimester of Pregnancy
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism
6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent
					treatment for Hyperthyroidism
8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies

REF: 1. TIETZ Fundamentals of Clinical chemistry 2.Guidlines of the American Thyroid association during pregnancy and Postpartum, 2011. **NOTE: It is advisable to detect Free T3,FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.**TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

\*\*End Of Report\*\*
Please visit www.agilusdiagnostics.com for related Test Information for this accession

### **CONDITIONS OF LABORATORY TESTING & REPORTING**

- 1. It is presumed that the test sample belongs to the patient named or identified in the test requisition form.
- 2. All tests are performed and reported as per the
- turnaround time stated in the AGILUS Directory of Services.
- 3. Result delays could occur due to unforeseen circumstances such as non-availability of kits / equipment breakdown / natural calamities / technical downtime or any other unforeseen event.
- 4. A requested test might not be performed if:
  - i. Specimen received is insufficient or inappropriate
  - ii. Specimen quality is unsatisfactory
  - iii. Incorrect specimen type
  - iv. Discrepancy between identification on specimen container label and test requisition form

- 5. AGILUS Diagnostics confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity.
- 6. Laboratory results should not be interpreted in isolation; it must be correlated with clinical information and be interpreted by registered medical practitioners only to determine final diagnosis.
- 7. Test results may vary based on time of collection, physiological condition of the patient, current medication or nutritional and dietary changes. Please consult your doctor or call us for any clarification.
- 8. Test results cannot be used for Medico legal purposes.
- 9. In case of queries please call customer care (91115 91115) within 48 hours of the report.

**Agilus Diagnostics Ltd** 

Fortis Hospital, Sector 62, Phase VIII, Mohali 160062

Chardni garg

Openult.

Dr.Pranjali Vasisht LAB HEAD Page 19 Of 19

DR.CHANDNI GARG
CONSULTANT PATHOLOGIST





View Details

View Report



Agilus Diagnostics Ltd. 24 Sco, Sector 11 D Chandigarh, 160011 Punjab, India

