PID No.
 : MED110827678
 Register On
 : 25/12/2021 9:31 AM

 SID No.
 : 421100127
 Collection On
 : 25/12/2021 11:41 AM

 Age / Sex
 : 43 Year(s) / Female
 Report On
 : 27/12/2021 7:02 PM

 Type
 : OP
 Printed On
 : 27/12/2021 8:10 PM

Ref. Dr : MediWheel

<u>Investigation</u>	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HAEMATOLOGY			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood'Spectrophotometry)	13.5	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood/Derived from Impedance)	42.1	%	37 - 47
RBC Count (EDTA Blood/Impedance Variation)	4.80	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (EDTA Blood/Derived from Impedance)	88.0	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood/Derived from Impedance)	28.2	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood/Derived from Impedance)	32.1	g/dL	32 - 36
RDW-CV (EDTA Blood/Derived from Impedance)	13.2	%	11.5 - 16.0
RDW-SD (EDTA Blood/Derived from Impedance)	40.66	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood/Impedance Variation)	7900	cells/cu.mm	4000 - 11000
Neutrophils (EDTA Blood/Impedance Variation & Flow Cytometry)	59.6	%	40 - 75
Lymphocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	31.3	%	20 - 45



VERIFIED BY



APPROVED BY

: MediWheel

 PID No.
 : MED110827678
 Register On
 : 25/12/2021 9:31 AM

 SID No.
 : 421100127
 Collection On
 : 25/12/2021 11:41 AM

 Age / Sex
 : 43 Year(s) / Female
 Report On
 : 27/12/2021 7:02 PM

Printed On

: 27/12/2021 8:10 PM

Type : OP

Ref. Dr

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Eosinophils (EDTA Blood/Impedance Variation & Flow Cytometry)	2.5	%	01 - 06
Monocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	6.1	%	02 - 10
Basophils (Blood/Impedance Variation & Flow Cytometry)	0.5	%	00 - 02
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	4.71	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	2.47	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.20	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.48	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.04	10^3 / μl	< 0.2
Platelet Count (EDTA Blood/Impedance Variation)	296	10^3 / μl	150 - 450
MPV (EDTA Blood/Derived from Impedance)	8.6	fL	8.0 - 13.3
PCT (EDTA Blood/Automated Blood cell Counter)	0.25	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood/Modified Westergren)	08	mm/hr	0 - 20



VERIFIED BY



APPROVED BY

 PID No.
 : MED110827678
 Register On
 : 25/12/2021 9:31 AM

 SID No.
 : 421100127
 Collection On
 : 25/12/2021 11:41 AM

 Age / Sex
 : 43 Year(s) / Female
 Report On
 : 27/12/2021 7:02 PM

 Type
 : OP
 Printed On
 : 27/12/2021 8:10 PM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulfanilic Acid)	0.4	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.2	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.20	mg/dL	0.1 - 1.0
Total Protein (Serum/Biuret)	7.8	g/dL	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.3	g/dL	3.5 - 5.0
Globulin (Serum/Derived)	3.50	g/dL	2.3 - 3.5
A : G Ratio (Serum/Derived)	1.23		1.1 - 2.4
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC without P5P)	22	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC without P5P)	25	U/L	5 - 41
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	59	U/L	42 - 98
GGT(Gamma Glutamyl Transpeptidase) (Serum/Modified IFCC)	18	U/L	< 38



Page 3 of 11

 PID No.
 : MED110827678
 Register On
 : 25/12/2021 9:31 AM

 SID No.
 : 421100127
 Collection On
 : 25/12/2021 11:41 AM

 Age / Sex
 : 43 Year(s) / Female
 Report On
 : 27/12/2021 7:02 PM

Printed On

Ref. Dr : MediWheel

: OP

Type

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/Cholesterol oxidase/Peroxidase)	194	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol phosphate oxidase / peroxidase)	178	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

: 27/12/2021 8:10 PM

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the õusualö"circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	48	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/Calculated)	110.4	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	35.6	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	146.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220



PID No. : MED110827678 Register On : 25/12/2021 9:31 AM : 421100127 SID No. Collection On : 25/12/2021 11:41 AM Age / Sex : 43 Year(s) / Female Report On : 27/12/2021 7:02 PM **Type** : OP

Printed On

Ref. Dr : MediWheel

Investigation <u>Observed</u> **Unit Biological** Reference Interval **Value**

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

: 27/12/2021 8:10 PM

Total Cholesterol/HDL Cholesterol Ratio 4 Optimal: < 3.3Low Risk: 3.4 - 4.4 (Serum/Calculated) Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio Optimal: < 2.53.7 Mild to moderate risk: 2.5 - 5.0 (TG/HDL)

High Risk: > 5.0(Serum/Calculated)

LDL/HDL Cholesterol Ratio 2.3 Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 (Serum/Calculated)

High Risk: > 6.0



Register On : 25/12/2021 9:31 AM PID No. : MED110827678 : 421100127 SID No. Collection On : 25/12/2021 11:41 AM Age / Sex : 43 Year(s) / Female Report On : 27/12/2021 7:02 PM **Type** : OP

Ref. Dr : MediWheel

Investigation <u>Observed</u> <u>Unit</u> **Biological** Reference Interval <u>Value</u> Glycosylated Haemoglobin (HbA1c) 5.6 HbA1C % Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 (Whole Blood/HPLC) Diabetic: ≥ 6.5

: 27/12/2021 8:10 PM

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Printed On

Estimated Average Glucose 114.02 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.



 PID No.
 : MED110827678
 Register On
 : 25/12/2021 9:31 AM

 SID No.
 : 421100127
 Collection On
 : 25/12/2021 11:41 AM

Age / Sex : 43 Year(s) / Female Report On : 27/12/2021 7:02 PM

Printed On

Type : OP

Ref. Dr : MediWheel

<u>Investigation</u> <u>Observed</u> <u>Unit</u> <u>Biological</u> <u>Value</u> <u>Reference Interval</u>

: 27/12/2021 8:10 PM

IMMUNOASSAY

THYROID PROFILE / TFT

Total T3 (Triiodothyronine) 0.85 ng/mL 0.8 - 2.0

(Serum/CMIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is

Metabolically active.

Total T4 (Thyroxine) 6.43 μg/dL 5.1-14.1

(Serum/CMIA)

INTERPRETATION:

Comment:

(Serum/CMIA)

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 2.76 µIU/mL Women (Non pregnant):0.27-4.2

Pregnant women 1st trimester: 0.1-2.5 2nd trimester: 0.2-3.0

2nd trimester: 0.2-3.0 3rd trimester: 0.3-3.0

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester: 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.

2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.

3. Values&lt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.

DR SRIJANA RAO S P
M D Pathology
Consultant Fathologist
KMC No :104646
APPROVED BY

PID No. : MED110827678 Register On : 25/12/2021 9:31 AM : 421100127 SID No. Collection On : 25/12/2021 11:41 AM Age / Sex : 43 Year(s) / Female Report On : 27/12/2021 7:02 PM : OP **Type Printed On** : 27/12/2021 8:10 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	Value		Reference Interval

CLINICAL PATHOLOGY

PHYSICAL EXAMINATION

Colour Pale Yellow

(Urine)

Volume 30 mL

(Urine)

Clear Clear Appearance

(Urine)

CHEMICAL EXAMINATION

6.5 4.6 - 8.0 pН

(Urine)

1.025 1.003 - 1.030 Specific Gravity

(Urine)

Negative Protein Negative

(Urine)

Glucose Negative Negative

(Urine)

Ketones Negative Negative

(Urine)

Negative Negative Leukocytes


(Urine)

Nitrite Negative Negative

(Urine)

Negative Negative Bilirubin

(Urine)



 PID No.
 : MED110827678
 Register On
 : 25/12/2021 9:31 AM

 SID No.
 : 421100127
 Collection On
 : 25/12/2021 11:41 AM

 Age / Sex
 : 43 Year(s) / Female
 Report On
 : 27/12/2021 7:02 PM

 Type
 : OP
 Printed On
 : 27/12/2021 8:10 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Blood	Negative		Negative
(Urine)			
Urobilinogen	0.1		0.1 - 1.0
(Urine)			
Urine Microscopy Pictures			
D. G. II		л с	0. 2
Pus Cells	6-8	/hpf	0 - 2
(Urine)	2.2	/I C	0. 2
Epithelial Cells (Urine)	2-3	/hpf	0 - 2
RBCs	Nil	/hpf	0 - 1
(Urine)	INII	/Hpi	0 - 1
Others	Nil		Nil
(Urine)	1111		1111
Casts	Nil	/hpf	0 - 1
(Urine)		1	
Crystals	Nil		NIL
(Urine)			
Bacteria	Nil		



 Age / Sex
 : 43 Year(s) / Female
 Report On
 : 27/12/2021 7:02 PM

 Type
 : OP
 Printed On
 : 27/12/2021 8:10 PM

Ref. Dr : MediWheel

InvestigationObservedUnitBiologicalValueReference Interval

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING 'B' 'Positive'

(EDTA Blood/Agglutination)



VERIFIED BY



APPROVED BY

 PID No.
 : MED110827678
 Register On
 : 25/12/2021 9:31 AM

 SID No.
 : 421100127
 Collection On
 : 25/12/2021 11:41 AM

 Age / Sex
 : 43 Year(s) / Female
 Report On
 : 27/12/2021 7:02 PM

 Type
 : OP
 Printed On
 : 27/12/2021 8:10 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	9.7		
Glucose Fasting (FBS) (Plasma - F/GOD- POD)	94	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Urine Glucose - Fasting (Urine - F/GOD - POD)	Sample Not Given		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD - POD)	116	mg/dL	70 - 140

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose (Postprandial) (Urine - PP/GOD - POD)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease-GLDH)	9	mg/dL	7.0 - 21
Creatinine (Serum/Modified Jaffe)	0.9	mg/dL	0.6 - 1.1

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine etc

Uric Acid **6.2** mg/dL 2.6 - 6.0

(Serum/Uricase/Peroxidase)



-- End of Report --