Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787 Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



General Physical Examination

Date of Examination: $26-11-2022$
Name: GAURAV AGARWAL Age: 31 Sex: Male
DOB: 19-04-1991
Referred By: BOB (Mediwheel)
Photo ID: AAD HAR ID#: attached ,
Ht: <u>186</u> (cm) Wt: <u>89</u> (Kg)
Chest (Expiration): 99 (cm) Abdomen Circumference: 96 (cm)
Blood Pressure: 126 / 80 mm Hg PR: 64 / min RR: 16 / min Temp: Alebail
BMI 25.7
Eye Examination: Dis. Visron 6 6 with space. Near Misron.
MG. (BIC eyes) Normal Colornision.
Other: Not svenificant
On examination he/she appears physically and mentally fit: Yes/No
Signature Of Examine :
Signature Medical Examiner: Name Medical Examiner



मेरा आधार, मेरी पहचान

po medical test puritore

Dr Piyush doyal M.B.B.S. D.M.R.D RMC Reg No-017996



DR.GOYAL PATH LAB & IMAGING CENTER, JAIPUR
2912 / MR GAURAV AGARWAL / 32 Yrs / M/ Non Smoker
Heart Rate: 68 bpm / Tested On: 26-Nov-22 14:31:12 / HF 0.05 Hz - LF 35 Hz / Notch 50 Hz / Sn 1.00 Cm/mV / Sw 25 mm/s / Refd By: MEDIWHEEL



ECG

REPRING BIO. 35703 MBBS, DIP, CARDIO (ESCORTS) Dr. Naresh Kumaradol D.E.M (RCGP-UK) P 20.00° T 49.00° Axis R 36.00° -30。 06 180° QT/QTC Int : 370/384 ms P-QRS-T axis: 20.00 36.00 49.00 Allengers ECG (Pisces)(PIS218210312) mdq 89 152 ms 90 ms QRS Duration: Vent Rate PR Interval



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date :- 26/11/2022 10:43:19

NAME :- Mr. GAURAV AGARWAL

Sex / Age :- Male 32 Yrs

Company :- MediWheel

Patient ID :-122228302

Ref. By Doctor:-Lab/Hosp :-

Final Authentication: 26/11/2022 11:17:58

BOB PACKAGE BELOW 40MALE

X RAY CHEST PA VIEW:

Both lung fields appears clear.

Bronchovascular markings appear normal.

Trachea is in midline.

Both the hilar shadows are normal.

Both the C.P.angles is clear.

Both the domes of diaphragm are normally placed.

Bony cage and soft tissue shadows are normal.

Heart shadows appear normal.

Impression: - Normal Study

(Please correlate clinically and with relevant further investigations)

DR ABHISHEK JAIN

MBBS. DNB. (RADIO DIAGNOSIS)

RMC NO. 21687

*** End of Report ***

AHSAN

Page No: 1 of 1



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Final Authentication: 26/11/2022 15:42:00

Date :- 26/11/2022 10:43:19

NAME :- Mr. GAÜRAV AGARWAL

Sex / Age :- Male 32 Yrs Company :- MediWheel Patient ID :-122228302 Ref. By Doctor:-Lab/Hosp :-

BOB PACKAGE BELOW 40MALE

USG WHOLE ABDOMEN

Liver is of normal size. Echo-texture is normal. No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is of normal size. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape. Echotexture is normal. No focal lesion is seen.

Kidneys are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. No focal lesion is seen. Collecting system does not show any dilatation or calculus.

Urinary bladder is well distended and showing smooth wall with normal thickness. Urinary bladder does not show any calculus or mass lesion.

Prostate is normal in size with normal echo-texture and outline.

No enlarged nodes are visualised. No retro-peritoneal lesion is identified No significant free fluid is seen in peritoneal cavity.

IMPRESSION:

* Normal study

Needs clinical correlation for further evaluation

*** End of Report ***

Page No: 1 of 1

P- Abhishek Jain MBBS, DNB, (Radio Diaynosis) RMC No. 21687

BILAL

Dr. Piyush Goyal M.B.B.S., D.M.R.D. RMC Reg No. 017996 Dr. Poonam Gupta MBBS, MD (Radio Diagnosis) RMC No. 32495 Dr. Ashish Choudhary
MBBS, MD (Radio Diagnosis)
Fetal Medicine Consultant
FMF ID - 260517 | RMC No 22430

Dr. Rathod Hetali Amrutlal MBBS, M.D. (Radio-Diagnosis) RMC No. 17163 Transcript by



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date :- 26/11/2022 10:43:19 NAME :- Mr. GAURAV AGARWAL

Sex / Age :- Male 32 Yrs Company :-MediWheel

Patient ID: -122228302 Ref. By Doctor:-Lab/Hosp :- '

Final Authentication: 26/11/2022 15:46:39

BOB PACKAGE BELOW 40MALE 2D ECHO OPTION TMT (ADULT/CHILD)

2D-ECHOCARDIOGRAPHY M.MODE WITH DOPPLER STUDY:

FAIR TRANSTHORACIC ECHOCARIDIOGRAPHIC WINDOW MORPHO

MITRAL VALV	/E	NOR	MAL		TRICUSPID VALVE				
AORTIC VALVE NORMAL					NORMAL				
M.MODE			ODE EXAMITATION:		ONARY VALVE	Y VALVE NO		NORMAL	
AO	24	mm	LA	32	Mm	IVS-D	17	mm	
IVS-S	. 15	mm	LVID	44	Mm	LVSD	27	mm	
LVPW-D	7-	mm	LVPW-S	15	Mm	RV	- 27	mm	
RVWT		mm	EDV		MI	LVVS	_		
LVEF	68%			DIAMAA				ml	
				RWMA		ABSENT			

		CH	AMBERS:	
LA	NORMAL	RA	NORMAL	
LV .	NORMAL	RV	NORMAL	
PERICARDIU	M	NORMAL	·	

		*		COI	OUR DOPPLER:			
	MI	TRAL VAI	LVE					
E VELOCITY	1.00	m/se	c PEAK	GRADIEN	Mm	Mm/hg		
A VELOCITY	0.60	m/se		N GRADIEN				
MVA BY PHT						Mm/	hg	
		Cm2	MVA	BY PLANIN	METRY	Cm2		
MITRAL REGURGITATION					ABSENT			
	- AO	RTIC VAL	VE					
PEAK VELOCITY	1.04	n	n/sec	PEAK G	RADIENT	lmm	mm/hg	
AR VMAX		· n	n/sec	MEAN O	GRADIENT		200,000	
AORTIC REGURGITATION			ABSENT mm/hg			/ng		
		JSPID VA	ALVE	POSEIVI				
PEAK VELOCITY .	0.68		m/sec	PEAK G	RADIENT		nm/hg	
MEAN VELOCITY			m/sec		GRADIENT			
VMax VELOCITY		714.00	1.1,000	- INCAR	UNADIENT	n	nm/hg	
RICUSPID REGURGITA	TION			ABSENT				
	PUL	MONARY	VALVE		• •			
EAK VELOCITY		0.90		M/sec.	PEAK GRADIENT		Mm/hg	
MEAN VALOCITY		1	•		MEAN GRADIENT			
PULMONARY REGURGIT	TATION						Mm/hg	
	A I I U I I				ABSENT		-	

Page No: 1 of 2

BILAL

Dr. Piyush Goyal M.B.B.S., D.M.R.D. RMC Reg No. 017996

Dr. Poonam Gupta MBBS, MD (Radio Diagnosis) RMC No. 32495

Dr. Ashish Choudhary MBBS, MD (Radio Diagnosis) Fetal Medicine Consultant FMF ID - 260517 | RMC No 22430

ABSENT

Dr. Rathod Hetali Amrutlal MBBS, M.D. (Radio-Diagnosis) RMC No. 17,163

Transcript by.



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date :- 26/11/2022 10:43:19

NAME :- Mr. GAURAV AGARWAL

Sex / Age :- Male 32 Yrs Company :- MediWheel Patient ID :-122228302 Ref. By Doctor:-Lab/Hosp :-

Final Authentication: 26/11/2022 15:46:39

Impression--

- 1. Normal LV size & contractility
- 2. No RWMA, LVEF 68 %.
- 3. Normal cardiac chamber.
- 4. Normal valve
- 5. No clot, no vegetation, no pericardial effusion.

(Cardiologist)

*** End of Report ***

Page No: 2 of 2

BILAL

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyusb@gmail.com



:- 26/11/2022* 10:43:19

NAME :- Mr. GAURAV AGARWAL

Sex / Age :- Male

Sample Type :- EDTA

32 Yrs

Company :- MediWheel

Patient ID: -122228302

Ref. By Dr:-

Lab/Hosp :-

Final Authentication: 26/11/2022 14:50:13

Sample Collected Time 26/11/2022 11:00:16 HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
BOB PACKAGE BELOW 40MALE HAEMOGARAM HAEMOGLOBIN (Hb) TOTAL LEUCOCYTE COUNT	15.6 7.45	g/dL	13.0 - 17.0
DIFFERENTIAL LEUCOCYTE COUNT	1.43	/cumm	4.00 - 10.00
NEUTROPHIL LYMPHOCYTE EOSINOPHIL MONOCYTE BASOPHIL NEUT# LYMPH# EO# MONO# BASO#	54.9 37.1 5.0 2.5 0.5 3.94 2.77 0.40 0.18 0.04	% % % % 10^3/uL 10^3/uL 10^3/uL 10^3/uL 10^3/uL	40.0 - 80.0 20.0 - 40.0 1.0 - 6.0 2.0 - 10.0 0.0 - 2.0 1.50 - 7.00 1.00 - 3.70 0.00 - 0.40 0.00 - 0.70 0.00 - 0.10
TOTAL RED BLOOD CELL COUNT (RBC) HEMATOCRIT (HCT) MEAN CORP VOLUME (MCV) MEAN CORP HB (MCH) MEAN CORP HB CONC (MCHC) PLATELET COUNT RDW-CV MENTZER INDEX	2.44 L 27.40 L 101.0 32.0 34.1 103 L 14.0 41.39	x10^6/uL % fL pg g/dL x10^3/uL	4.50 - 5.50 40.00 - 50.00 83.0 - 101.0 27.0 - 32.0 31.5 - 34.5 150 - 410 11.6 - 14.0

The Mentzer index is used to differentiate iron deficiency anemia from beta thalassemia trait. If a CBC indicates microcytic anemia, these are two of the most likely causes, making it necessary to distinguish between them.

If the quotient of the mean corpuscular volume divided by the red blood cell count is less than 13, thalassemia is more likely. If the result is greater than 13, then iron-deficiency anemia is more likely.

MUKESHSINGH **Technologist**

Page No: 1 of 11



Dr. Goya

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date :- 26/11/2022 10:43:19

NAME :- Mr. GAURAV AGARWAL

Sex / Age :- Male

Sample Type :- EDTA

32 Yrs Company :- MediWheel

Sample Collected Time 26/11/2022 11:00:16

Final Authentication: 26/11/2022 14:50:13

HAEMATOLOGY

Test Name Value Unit **Biological Ref Interval**

Ref. By Dr:-

Lab/Hosp :-

Erythrocyte Sedimentation Rate (ESR)

16 .H

mm/hr.

Patient ID :-122228302

00 - 13

(ESR) Methodology: Measurment of ESR by cells aggregation.

Instrument Name : Indepedent form Hematocrit value by Automated Analyzer (Roller-20)

: ESR test is a non-specific indicator ofinflammatory disease and abnormal protein states.

The test in used to detect, follow course of a certain disease (e.g-tuberculosis, rheumatic fever, myocardial infarction

Levels are higher in pregnency due to hyperfibrinogenaemia.

The "3-figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia of Bonnettye disease. Fluorescent Flow cytometry, HB SLS method, TRBC, PCV, PLT Hydrodynamically focused Impedance. and MCH, MCV, MCHC, MENTZER INDEX are calculated. InstrumentName: Sysmex 6 part fully automatic analyzer XN-L, Japan

MUKESHSINGH **Technologist**

Page No: 2 of 11



r. Goya

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 26/11/2022 10:43:19

Patient ID :-122228302

NAME :- Mr. GAURAV AGARWAL Sex / Age :- Male

32 Yrs

Ref. By Dr:-

Company :- MediWheel

Lab/Hosp :-

Sample Type :- EDTA, KOx/Na FLUORIDE-F, KSamhdal-Collegian ETPR-20511112022 11:00:16

Final Authentication: 26/11/2022 15:35:56

HAEMATOLOGY

Test Name

Value

Unit

Biological Ref Interval

BLOOD GROUP ABO

"B"POSITIVE .

BLOOD GROUP ABO Methodology: Haemagglutination reaction Kit Name: Monoclonal agglutinating antibodies (Span clone).

FASTING BLOOD SUGAR (Plasma)

Method:- GOD PAP

94.6

mg/dl

75.0 - 115.0

Impaired glucose tolerance (IGT)	111 - 125 mg/dL	
Diabetes Mellituş (DM)	> 126 mg/dL	

Instrument Name: Randox Rx Imola Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels(hypoglycemia) may result from excessive insulin therapy or various liver diseases .

BLOOD SUGAR PP (Plasma)

117.9

mg/dl

70.0 - 140.0

Instrument Name: Randox Rx Imola Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels(hypoglycemia) may result from excessive insulin therapy or various liver diseases . .

URINE SUGAR (FASTING)
Collected Sample Received

Nil

KAUSHAL, MUKESHSINGH, POOJABOHRA **Technologist** DR.HANSA Page No: 3 of 11

Dr. Piyush Goyal (D.M.R.D.) Dr. Rashmi Bakshi Dr. Chandrika Gupta

Path Lab & Imaging Centre

. B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787 Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 26/11/2022 10:43:19

NAME :- Mr. GAURAV AGARWAL

Patient ID: -122228302 Ref. By Dr:-

Sex / Age :- Male

32 Yrs

Lab/Hosp.:-

Company :- MediWheel Sample Type :- STOOL

Sample Collected Time 26/11/2022 11:00:16

Final Authentication: 26/11/2022 15:12:25

CLINICAL PATHOLOGY

Test Name Value Unit **Biological Ref Interval**

STOOL ANALYSIS

PHYSICAL EXAMINATION

MUCUS **BLOOD**

MICROSCOPIC EXAMINATION

RBC's

WBC/HPF

OVA

CYSTS

OTHERS Collected Sample Received

/HPF

/HPF

POOJABOHRA Technologist DR.HANSA Page No: 4 of 11




Dr. Rashmi Bakshi MBBS. MD (Path) RMC No. 17975/008828

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787.

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 26/11/2022 10:43:19 NAME :- Mr. GAURAV AGARWAL

Sex / Age :- Male 32 Yrs

Company :- MediWheel Sample Type :- PLAIN/SERUM Patient ID: -122228302

Ref. By Dr:- .

Lab/Hosp :-



Final Authentication: 26/11/2022 14:05:38

BIOCHEN	MISTRY
----------------	--------

Sample Collected Time 26/11/2022 11:00:16

Test Name	¥	Value	Unit	Biological Ref Interval
LIPID PROFILE	8			
TOTAL CHOLESTEROL Method:- Enzymatic Endpoint Method	*	130.37	mg/dl	Desirable <200 Borderline 200-239
TRIGLYCERIDES Method:- GPO-PAP		82.59	mg/dl	High> 240 Normal <150 Borderline high 150-199 High 200-499 Very high >500
DIRECT HDL CHOLESTEROL Method:- Direct clearance Method	•	41.76	mg/dl	Low < 40 High > 60
DIRECT LDL CHOLESTEROL Method:- Direct clearance Method		74.85	mg/dl	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159
				High 160-189 Very High > 190
VLDL CHOLESTEROL Method:- Calculated		16.52	mg/dl	0.00 - 80.00
T.CHOLESTEROL/HDL CHOLE Method:- Calculated	STEROL RATIO	3.12		0.00 - 4.90
LDL / HDL CHOLESTEROL RA'	TIO	1.79		0.00 - 3.50
TOTAL LIPID Method:- CALCULATED TOTAL CHOI ESTEROL I DESTRUMENT NO.		· 395.97 L	mg/dl	400.00 - 1000.00

TOTAL CHOLESTEROL InstrumentName: Randox Rx Imola Interpretation: Cholesterol measurements are used in the diagnosis and treatments of lipid lipoprotein metabolism

TRIGLYCERIDES InstrumentName: Randox Rx Imola Interpretation: Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction.

DIRECT HDLCHOLESTERO InstrumentName: Randox Rx Imola Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.

DIRECT LDL-CHOLESTEROLInstrumentName: Randox Rx Imola Interpretation: Accurate measurement of LDL-Cholesterol is of vital importance in therapies which focus on lipid reduction to prevent atherosclerosis or reduce its progress and to avoid plaque rupture.

TOTAL LIPID AND VLDL ARE CALCULATED

KAUSHAL

Page No: 5 of 11




Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgovalspathlab.com | E-mail: drgovalpivush@gmail.com



NAME :- Mr. GAURAV AGARWAL

Sex / Age :- Male 32 Yrs

Company :- MediWheel

Sample Type :- PLAIN/SERUM

Patient ID: -122228302

Ref. By Dr:-

Lab/Hosp:-



Final Authentication: 26/11/2022 14:05:38

BIOCHEMISTRY

Sample Collected Time 26/11/2022 11:00:16

		DIOCHE	IISIKI	
Test Name		Value	Unit	Biological Ref Interval
LIVER PROFILE WITH GGT ·				
SERUM BILIRUBIN (TOTAL) Method:- Colorimetric method		1.22	mg/dl	Up to - 1.0 Cord blood <2 Premature < 6 days <16
	í			Full-term < 6 days= 12 1month - <12 months <2 1-19 years <1.5 Adult - Up to - 1.2 Ref-(ACCP 2020)
SERUM BILIRUBIN (DIRECT) Method:- Colorimetric Method	•	0.44	mg/dL	Adult - Up to 0.25 Newborn - <0.6 mg/dL >- 1 month - <0.2 mg/dL
SERUM BILIRUBIN (INDIRECT) Method:- Calculated	•	0.78	mg/dl	0.30-0.70
SGOT Method:- IFCC		24.7	U/L	Men- Up to - 37.0 Women - Up to - 31.0
SGPT Method:- IFCC		18.1	U/L	Men- Up to - 40.0 Women - Up to - 31.0
SERUM ALKALINE PHOSPHATASE Method:-AMP Buffer		78.50	IU/L	30.00 - 120.00
SERUM GAMMA GT Method:- IFCC		20.40	· U/L	11.00 - 50.00
SERUM TOTAL PROTEIN Method:- Biuret Reagent		6.72	g/dl	6.40 - 8.30
SERUM ALBUMIN Method:- Bromocresol Green		4.19	g/dl	3.80 - 5.00
SERUM GLOBULIN Method:- CALCULATION	· • 27	2.53	gm/dl	2.20 - 3.50
A/G RATIO	i	1.66		1.30 - 2.50

Total BilirubinMethodology:Colorimetric method InstrumentName:Randox Rx Imola Interpretation An increase in bilirubin concentration in the serum occurs in toxic or infectious diseases of the liver e.g. hepatitis B or obstruction of the bile duct and in rhesus incompatible babies. High levels of unconjugated bilirubin indicate that too much haemoglobin is being destroyed or that the liver is not actively treating the haemoglobin it is receiving.

AST Aspartate Aminotransferase Methodology: IFCC InstrumentName:Randox Rx Imola Interpretation: Elevated levels of AST can signal myocardial infarction, hepatic disease, muscular dystrophy and organ damage. Although heart muscle is found to have the most activity of the enzyme, significant activity has also been seen in the brain, liver, gastric mucosa, adipose tissue and kidneys of humans.

ALT Alanine Aminotransferase Methodology: IFCCInstrumentName:Randox Rx Imola Interpretation: The enzyme ALT has been found to be in highest concentrations in the liver, with decreasing concentrations found in kidney, heart, skeletal muscle; pancreas, spleen and lung tissue respectively. Elevated levels of the transaminases can indicate myocardial infarction, hepatic disease, muscular dystrophy and organ damage.

Alkaline Phosphatase Methodology: AMP Buffer InstrumentName: Randox Rx Imola Interpretation: Measurements of alkaline phosphatase are of use in the diagnosis, treatment and investigation of hepatobilary disease and in bone disease associated with increased osteoblastic activity. Alkaline phosphatase is also used in the diagnosis of parathyroid and intestinal disease.

TOTAL PROTEIN Methodology: Bruret Reagent Instrument Name: Randox Rx Imola Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

ALBUMIN (ALB) Methodology: Bromocresol Green InstrumentName:Randox Rx Imola Interpretation: Albumin measurements are used in the diagnosis and treatment of numerous diseases involving primarily the liver or kidneys. Globulin & A/G ratio is calculated.

Instrument Name Randox Rx Imola Interpretation: Elevations in GGT levels are seen earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or post-hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal)

KAUSHAL

Page No: 6 of 11



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019
Tele: D141-2293346, 4049787, 9887049787
Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date :- 26/11/2022 10:43:19

NAME :- Mr. GAURAV AGÁRWAL

Ref. By Dr:-Lab/Hosp :-

Patient ID: -122228302

Sex / Age :- Male 32 Yrs Company :- MediWheel

Sample Type :- PLAIN/SERUM

Sample Collected Time 26/11/2022 11:00:16

Final Authentication: 26/11/2022 14:05:38

Women - 2.4-5.7

		BIOCHEM	IISTRY	
Test Name	Name		Unit	Biological Ref Interval
SERUM CREATININE Method:- Colorimetric Method	*	1.00	mg/dl	Men - 0.6-1.30 Women - 0.5-1.20
SERUM URIC ACID + Method:- Enzymatic colorimetric	•	4.91	mg/dl	Men - 3.4-7.0

KAUSHAL

Page No: 7 of 11 ·



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787 Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 26/11/2022 10:43:19

NAME :- Mr. GAURAV AGARWAL

Sex / Age :- Male 32 Yrs

Company :- MediWheel

Patient ID: -122228302

Ref. By Dr:-

Lab/Hosp :-

Sample Type :- PLAIN/SERUM

Sample Collected Time 26/11/2022 11:00:16

Final Authentication: 26/11/2022 14:05:38

BIOCHEMISTRY

Test Name

Value

Unit

Biological Ref Interval

BLOOD UREA NITROGEN (BUN)

13.7

mg/dl

0.0 - 23.0

KAUSHAL


Page No: 8 of 11



B-51, Ganesh Nagar, Opp. Janpath Corner. New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



NAME :- Mr. GAURAV AGARWAL

Sex / Age :- Male 32 Yrs

Company :-MediWheel

Sample Type :- EDTA

Patient ID: -122228302

Ref. By Dr:-

Lab/Hosp :-



Sample Collected Time 26/11/2022 11:00:16

Final Authentication: 26/11/2022 14:50:13

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
GLYCOSYLATED HEMOGLOBIN (HbA1C) Method:-HPLC	5.2	%	Non-diabetic: < 5.7

Pre-diabetics: 5.7-6.4 Diabetics: = 6.5 or higher ADA Target: 7.0 Action suggested: > 6.5

Instrument name: ARKRAY's ADAMS Lite HA 8380V, JAPAN.

Test Interpretation:

HbA1C is formed by the condensation of glucose with n-terminal valine residue of each beta chain of HbA to form an unstable schiff base. It is the major fraction, constituting approximately 80% of HbA1c. Formation of glycated hemoglobin (GHb) is essentially irreversible and the concentration in the blood depends on both the lifespan of the red blood cells (RBC) (120 days) and the blood glucose concentration. The GHb concentration represents the integrated values for glucose overthe period of 6 to 8 weeks. GHb values are free of day to day glucose fluctuations and are unaffected by recent exercise or food ingestion. Concentration of plasmaglucose concentration in GHb depends on the time interval, with more recent values providing a larger contribution than earlier values. The interpretation of GHbdepends on RBC having a normal life span. Patients with hemolytic disease or other conditions with shortened RBC survival exhibit a substantial reduction of GHb. High GHb have been reported in iron deficiency anemia. GHb has been firmly established as an index of long term blood glucose concentrations and as a measureof the risk for the development of complications in patients with diabetes mellitus. The absolute risk of retinopathy and nephropathy are directly proportional to themean of HbA1C.Genetic variants (e.g. HbS trait, HbC trait), elevated HbF and chemically modified derivatives of hemoglobin can affect the accuracy of HbA1cmeasurements. The effects vary depending on the specific Hb vatiant or derivative and the specific HbA1c method.

Ref by ADA 2020

MEAN PLASMA GLUCOSE Method:- Calculated Parameter

103.

mg/dL

Non Diabetic < 100 mg/dL Prediabetic 100- 125 mg/dL Diabetic 126 mg/dL or Higher

MUKESHSINGH **Technologist**


Page No: 9 of 11



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787.

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



NAME :- Mr. GAURAV AGARWAL

Sex / Age :- Male 32 Yrs

Company :- MediWheel
Sample Type :- URINE

Patient ID: -122228302

Ref. By Dr:-

Lab/Hosp :-

Final Authentication: 26/11/2022 15:12:25

Sample Collected Time 26/11/2022 11:00:16 CLINICAL PATHOLOGY

Test Name		Value	Unit		Biological	Ref Interval
Urine Routine						
PHYSICAL EXAMINA	TION ·	-,				
· COLOUR	•	PALE YEL	ĹOW	P	ALE YELLOW	
APPEARANCE		Clear			Clear	
CHEMICAL EXAMINA	ATION					
REACTION(PH)	,	5.5		5.	.0 - 7.5	
SPECIFIC GRAVITY	•	1.025		1.	.010 - 1.030	
PROTEIN	. •	. NIL		N	IIL	
SUGAR		, NIL.		N	IIL	
BILIRUBIN		NEGATIVI	3	· N	EGATIVE	
UROBILINOGEN	+	NORMAL		N	ORMAL	
KETONES		NEGATIVI	3	. N	EGATIVE	
NITRITE .		NEGATIVE	3	N	EGATIVE	*0
MICROSCOPY EXAM	INATION					
RBC/HPF		NIL	/HPF	N	IL	•
WBC/HPF		2-3	/HPF	2-	-3	
EPITHELIAL CELLS		. , 2-3	/HPF	2-	-3	
CRYSTALS/HPF	•	ABSENT		A	BSENT	
· CAST/HPF		ABSENT		A	BSENT	*
AMORPHOUS SEDIMENT	`	ABSENT		A	BSENT	
BACTERIAL FLORA		ABSENT		A	BSENT	
YEAST CELL	•	ABSENT		A	BSENT	
OTHER	•	ABSENT				

POOJABOHRA Technologist DR.HANSA Page No: 10 of 11




Dr. Rashmi Bakshi MBBS. MD (Path) RMC No. 17975/008828

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sangarier Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



NAME :- Mr. GAURAV AGARWAL

Sex / Age :- Male 32 Yrs

Company :- MediWheel
Sample Type :- PLAIN/SERUM

Patient ID: -122228302

Ref. By Dr:-

Lab/Hosp :-



Sample Collected Time 26/11/2022 11:00:16 Final Authentication: 26/11/2022 13:27:35

IMMUNOASSAY

Test Name	Value	Unit	Biological Ref Interval
TOTAL THYROID PROFILE			
SERUM TOTAL T3 Method:- Chemiluminescence(Competitive immunoassay)	1.032	ng/ml	0.970 - 1.690
SERUM TOTAL T4 Method:- Chemiluminescence(Competitive immunoassay)	7.780	ug/dl .	5.530 - 11.000
SERUM TSH ULTRA Method:- Enhanced Chemiluminescence Immunoassay	5.239 H	μIU/mL .	0.400 - 4.649

Interpretation: Triiodothyronine (T3) contributes to the maintenance of the euthyroid state. A decrease in T3 concentration of up to 50% occurs in a variety of clinical situations, including acute and chronic disease. Although T3 results alone cannot be used to diagnose hypothyroidism, T3 concentration may be more sensitive than thyroxine (T4) for hyperthyroidism. Consequently, the total T3 assay can be used in conjunction with other assays to aid in the differential diagnosis of thyroid disease. T3 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, Free T3 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake, or T4-uptake can be used with the total T3 result to calculate the free T3 index and estimate the concentration of free T3.

Interpretation: The measurement of Total T4 aids in the differential diagnosis of thyroid disease. While >99.9% of T4 is protein-bound, primarily to thyroxine-binding globulin (TBG), it is the free fraction that is biologically active. In most patients, the total T4 concentration is a good indicator of thyroid status. T4 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, free T4 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake may be used with the total T4 result to calculate the free T4 index (FT4I) and estimate the concentration of free T4. Some drugs and some nonthyroidal patient conditions are known to alter TT4 concentrations in vivo.

Interpretation: TSH stimulates the production of thyroxine (T4) and triiodothyronine (T3) by the thyroid gland. The diagnosis of overt hypothyroidism by the finding of a low total T4 or free T4 concentration is readily confirmed by a raised TSH concentration. Measurement of low or undetectable TSH concentrations may assist the diagnosis of hyperthyroidism, where concentrations of T4 and T3 are elevated and TSH secretion is suppressed. These have the advantage of discriminating between the concentrations of TSH observed in thyrotoxicosis, compared with the low, but detectable, concentrations that occur in subclinical hyperthyroidism. The performance of this assay has not been established for neonatal specimens. Some drugs and some nonthyroidal patient conditions are known to alter TSH concentrations in vivo.

INTERPRETATION

PREGNANCY	REFERENCE RANGE FOR TSH IN uIU/mL (As per American Thyroid Association)	
1st Trimester	0.10-2.50	
2nd Trimester	0.20-3.00	
3rd Trimester	0.30-3.00	

*** End of Report ***

NARENDRAKUMAR Technologist

Page No: 11 of 11



