Rate

Male

36 Years

. ST elev, probable normal early repol pattern.....ST elevation, age<55 132 PR 91 QRSD 362 QT QTc 431 --AXIS--75 75 - NORMAL ECG -QRS 53 12 Lead; Standard Placement Unconfirmed Diagnosis **V**1 1 aVR in the same of the 100B CL? Chest: 10.0 mm/mV F 60~ 0.15-100 Hz Speed: 25 mm/sec Limb: 10 mm/mV Device:

Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name : MR MANISH KUMAR Age : 36 Yr(s) Sex :Male

Referred By : HEALTH CHECK MHD Reporting Date : 13 Mar 2024 12:56

Receiving Date : 13 Mar 2024 11:15

Department of Transfusion Medicine (Blood Bank)

BLOOD GROUPING, RH TYPING & ANTIBODY SCREEN (TYPE & SCREEN) Specimen-Blood

Blood Group & Rh Typing (Agglutinaton by gel/tube technique)

Blood Group & Rh typing B Rh(D) Positive

Antibody Screening (Microtyping in gel cards using reagent red cells)

Final Antibody Screen Result Negative

Technical Note:

ABO grouping and Rh typing is done by cell and serum grouping by microplate / gel technique. Antibody screening is done using a 3 cell panel of reagent red cells coated with Rh, Kell, Duffy, Kidd, Lewis, P, MNS, Lutheran and Xg antigens using gel technique.

Page 1 of 4

-----END OF REPORT-----

Dampa

Dr Himanshu Lamba

Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name : MR MANISH KUMAR Age : 36 Yr(s) Sex : Male

Referred By: HEALTH CHECK MHD **Reporting Date**: 13 Mar 2024 13:45

Receiving Date : 13 Mar 2024 11:01

BIOCHEMISTRY

Specimen: EDTA Whole blood

As per American Diabetes Association(ADA) 2010

HbA1c (Glycosylated Hemoglobin) 5.3 % [4.0-6.5]

HbA1c in %

Non diabetic adults : < 5.7 %

Prediabetes (At Risk) : 5.7 % - 6.4 %

Diabetic Range : > 6.5 %

Methodology High-Performance Liquid Chromatography (HPLC)

Estimated Average Glucose (eAG) 105 mg/dl

Use

- 1.Monitoring compliance and long-term blood glucose level control in patients with diabetes.
- 2. Index of diabetic control (direct relationship between poor control and development of complications).
- 3. Predicting development and progression of diabetic microvascular complications.

Limitations :

- 1. A1C values may be falsely elevated or decreased in those with chronic kidney disease.
- 2.False elevations may be due in part to analytical interference from carbamylated hemoglobin formed in the presence of elevated concentrations of urea, with some assays.
- 3. False decreases in measured A1C may occur with hemodialysis and altered red cell turnover, especially in the setting of erythropoietin treatment

References: Rao.L.V., Michael snyder.L.(2021). Wallach's Interpretation of Diagnostic Tests. 11th Edition. Wolterkluwer. NaderRifai, Andrea Rita Horvath, Carl T.wittwer. (2018) Teitz Text book

of Clinical Chemistry and Molecular Diagnostics. First edition, Elsevier, South Asia.

Page 2 of 4

P 011 4967 4967 **E** info@manipalhospitals.com **Emergency** 011 4040 7070

www.hcmct.in www.manipalhospitals.com/delhi/

Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

 Name
 : MR MANISH KUMAR
 Age
 : 36 Yr(s) Sex :Male

 Registration No
 : MH011770913
 Lab No
 : 32240306656

 Patient Episode
 : H03000060971
 Collection Date : 13 Mar 2024 10:15

Referred By : HEALTH CHECK MHD

Receiving Date : 13 Mar 2024 11:01

BIOCHEMISTRY

Reporting Date:

13 Mar 2024 15:30

Lipid Profile (Serum)

| TOTAL CHOLESTEROL (CHOD/POD) | 206 # | mg/dl | [<200] Moderate risk:200-239 High risk:>240 |
|---|-------|------------|--|
| TRIGLYCERIDES (GPO/POD) | 143 | mg/dl | [<150] Borderline high:151-199 High: 200 - 499 Very high:>500 |
| HDL - CHOLESTEROL (Direct) | 67 # | mg/dl | [30-60] |
| Methodology: Homogenous Enzymatic VLDL - Cholesterol (Calculated) | 29 | mg/dl | [10-40] |
| | | | |
| (CALCULATED) LDL- CHOLEST T.Chol/HDL.Chol ratio | 3.1 | 110 #mg/dl | [<100] Near/Above optimal-100-129 Borderline High:130-159 High Risk:160-189 <4.0 Optimal 4.0-5.0 Borderline >6 High Risk |

Note:

Reference ranges based on ATP III Classifications. Recommended to do fasting Lipid Profile after a minimum of 8 hours of overnight fasting.

Technical Notes:

Lipid profile is a panel of blood tests that serves as initial broad medical screening tool for abnormalities in lipids, the results of these tests can identify certain genetic

Page 3 of 4

P 011 4967 4967 **E** info@manipalhospitals.com **Emergency** 011 4040 7070

www.hcmct.in www.manipalhospitals.com/delhi/

Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name : MR MANISH KUMAR Age : 36 Yr(s) Sex :Male

Referred By : HEALTH CHECK MHD Reporting Date : 13 Mar 2024 15:30

Receiving Date : 13 Mar 2024 11:01

BIOCHEMISTRY

diseases and determine approximate risks for cardiovascular disease, certain forms of pancreatitis and other diseases.

Page 4 of 4

----END OF REPORT----

Dr. Neelam Singal

CONSULTANT BIOCHEMISTRY

Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name : MR MANISH KUMAR Age : 36 Yr(s) Sex :Male

Referred By : HEALTH CHECK MHD
Receiving Date : 13 Mar 2024 11:01

BIOCHEMISTRY

Reporting Date:

14 Mar 2024 07:32

| THYROID PROFILE, Serum | | 2 | Specimen Type : Serum |
|-------------------------------------|-------|--------|-----------------------|
| T3 - Triiodothyronine (ECLIA) | 1.460 | ng/ml | [0.800-2.040] |
| T4 - Thyroxine (ECLIA) | 7.590 | μg/dl | [4.600-10.500] |
| Thyroid Stimulating Hormone (ECLIA) | 2.350 | μIU/mL | [0.340-4.250] |

Note: TSH levels are subject to circadian variation, reaching peak levels between 2-4.a.m.and at a minimum between 6-10 pm.Factors such as change of seasons hormonal fluctuations, Ca or Fe supplements, high fibre diet, stress and illness affect TSH results.

- * References ranges recommended by the American Thyroid Association
- 1) Thyroid. 2011 Oct; 21(10):1081-125.PMID .21787128
- 2) http://www.thyroid-info.com/articles/tsh-fluctuating.html

| Test Name | Result | Unit | Biological Ref. Interval |
|------------------------------------|--------|-------|--------------------------|
| LIVER FUNCTION TEST (Serum) | | | |
| BILIRUBIN-TOTAL (Diazonium Ion) | 0.82 | mg/dl | [0.10-1.20] |
| BILIRUBIN - DIRECT (Diazotization) | 0.27 | mg/dl | [0.00-0.30] |
| BILIRUBIN - INDIRECT (Calculated) | 0.55 | mg/dl | [0.20-1.00] |
| SGOT/ AST (UV without P5P) | 29.7 | U/L | [10.0-50.0] |
| SGPT/ ALT (UV without P5P) | 36.4 | U/L | [0.0-41.0] |
| ALP (p-NPP, kinetic) * | 89 | U/L | [45-135] |
| TOTAL PROTEIN (Biuret) | 8.0 | g/dl | [6.0-8.2] |
| SERUM ALBUMIN (BCG-dye) | 4.7 | g/dl | [3.5-5.2] |
| SERUM GLOBULIN (Calculated) | 3.3 | g/dl | [1.8-3.4] |
| ALB/GLOB (A/G) Ratio(Calculated) | 1.42 | | [1.10-1.80] |

Page 1 of 7



Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name : MR MANISH KUMAR Age : 36 Yr(s) Sex :Male

Referred By: HEALTH CHECK MHD Reporting Date: 13 Mar 2024 15:30

Receiving Date : 13 Mar 2024 11:01

BIOCHEMISTRY

Technical Notes:

Liver function test aids in diagnosis of various pre hepatic, hepatic and post hepatic causes of dysfunction like hemolytic anemia's, viral and alcoholic hepatitis and cholestasis of obstructive causes.

| Test Name | Result | Unit B | iological Ref. Interval |
|-----------------------------------|--------|---------------|-------------------------|
| KIDNEY PROFILE (Serum) | | | |
| BUN (Urease/GLDH) | 9.00 | mg/dl | [6.00-20.00] |
| SERUM CREATININE (Jaffe's method) | 0.85 | mg/dl | [0.80-1.60] |
| SERUM URIC ACID (Uricase) | 6.8 | mg/dl | [3.5-7.2] |
| SERUM CALCIUM (NM-BAPTA) | 9.58 | mg/dl | [8.00-10.50] |
| SERUM PHOSPHORUS (Molybdate, UV) | 2.5 | mg/dl | [2.5-4.5] |
| SERUM SODIUM (ISE) | 139.0 | mmol/l | [134.0-145.0] |
| SERUM POTASSIUM (ISE) | 5.07 | mmol/l | [3.50-5.20] |
| SERUM CHLORIDE (ISE Indirect) | 103.1 | mmol/L | [95.0-105.0] |
| eGFR | 112.1 | ml/min/1.73sq | .m [>60.0] |
| | | | |

Technical Note

eGFR which is primarily based on Serum Creatinine is a derivation of CKD-EPI 2009 equation normalized to1.73 sq.m BSA and is not applicable to individuals below 18 years. eGFR tends to be less accurate when Serum Creatinine estimation is indeterminate e.g. patients at extremes of muscle mass, on unusual diets etc. and samples with severe Hemolysis / Icterus / Lipemia.

Page 2 of 7

----END OF REPORT----

Dr. Neelam Singal

CONSULTANT BIOCHEMISTRY



Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name : MR MANISH KUMAR Age : 36 Yr(s) Sex : Male

Referred By: HEALTH CHECK MHD Reporting Date: 14 Mar 2024 07:42

Receiving Date : 13 Mar 2024 18:03

BIOCHEMISTRY

Specimen Type : Plasma
PLASMA GLUCOSE - PP

Plasma GLUCOSE - PP (Hexokinase) 122 mg/dl [70-140]

Note: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying,

brisk glucose absorption , post exercise

Specimen Type : Serum/Plasma

Plasma GLUCOSE-Fasting (Hexokinase) 89 mg/dl [74-106]

Page3 of 7

-----END OF REPORT-----

Dr. Neelam Singal

CONSULTANT BIOCHEMISTRY

Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name : MR MANISH KUMAR Age : 36 Yr(s) Sex :Male

Referred By: HEALTH CHECK MHD Reporting Date: 13 Mar 2024 13:02

Receiving Date : 13 Mar 2024 10:50

HAEMATOLOGY

ERYTHROCYTE SEDIMENTATION RATE (Automated) Specimen-Whole Blood

ESR 14.0 # mm/1sthour [0.0-10.0]

Interpretation :

Erythrocyte sedimentation rate (ESR) is a non-specific phenomena and is clinically useful in the diagnosis and monitoring of disorders associated with an increased production of acute phase reactants (e.g. pyogenic infections, inflammation and malignancies). The ESR is increased in pregnancy from about the 3rd month and returns to normal by the 4th week postpartum.

ESR is influenced by age, sex, menstrual cycle and drugs (eg. corticosteroids, contraceptives).

It is especially low (0 - 1mm) in polycythemia, hypofibrinogenemia or congestive cardiac failure and when there are abnormalities of the red cells such as poikilocytosis, spherocytosis or sickle cells.

| Test Name | Result | Unit Bi | ological Ref. Interval |
|------------------------------------|--------|---------------|------------------------|
| COMPLETE BLOOD COUNT (EDTA Blood) | | | |
| WBC Count (Flow cytometry) | 5900 | /cu.mm | [4000-10000] |
| RBC Count (Impedence) | 5.12 | million/cu.mm | [4.50-5.50] |
| Haemoglobin (SLS Method) | 15.2 | g/dL | [13.0-17.0] |
| Haematocrit (PCV) | 48.0 | % | [40.0-50.0] |
| (RBC Pulse Height Detector Method) | | | |
| MCV (Calculated) | 93.8 | fL | [83.0-101.0] |
| MCH (Calculated) | 29.7 | bà | [25.0-32.0] |
| MCHC (Calculated) | 31.7 | g/dL | [31.5-34.5] |
| Platelet Count (Impedence) | 200000 | /cu.mm | [150000-410000] |
| RDW-CV (Calculated) | 14.4 # | % | [11.6-14.0] |
| DIFFERENTIAL COUNT | | | |
| Neutrophils (Flowcytometry) | 67.1 | ૄ | [40.0-80.0] |
| Lymphocytes (Flowcytometry) | 25.3 | ૾ૢ | [20.0-40.0] |

Page 4 of 7



Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name : MR MANISH KUMAR Age : 36 Yr(s) Sex : Male

Referred By : HEALTH CHECK MHD Reporting Date : 13 Mar 2024 13:02

Receiving Date : 13 Mar 2024 10:50

HAEMATOLOGY

| Monocytes (Flowcytometry) | 5.6 | | 9 | [2.0-10.0] |
|--------------------------------------|----------------|-----|--------|---------------------------|
| Eosinophils (Flowcytometry) | 1.5 | | 용 | [1.0-6.0] |
| Basophils (Flowcytometry) | 0.5 # | | % | [1.0-2.0] |
| IG | 0.30 | | 용 | |
| Neutrophil Absolute (Flouroscence fi | low cytometry) | 4.0 | /cu mm | $[2.0-7.0] \times 10^{3}$ |
| Lymphocyte Absolute (Flouroscence fi | low cytometry) | 1.5 | /cu mm | $[1.0-3.0] \times 10^{3}$ |
| Monocyte Absolute (Flouroscence flow | w cytometry) | 0.3 | /cu mm | $[0.2-1.2] \times 10^{3}$ |
| Eosinophil Absolute (Flouroscence fi | low cytometry) | 0.1 | /cu mm | $[0.0-0.5] \times 10^{3}$ |
| Basophil Absolute (Flouroscence flow | w cytometry) | 0.0 | /cu mm | $[0.0-0.1] \times 10^{3}$ |

Complete Blood Count is used to evaluate wide range of health disorders, including anemia, infection, and leukemia. Abnormal increase or decrease in cell counts as revealed may indicate that an underlying medical condition that calls for further evaluation.

-----END OF REPORT-----

Dr. Shalakha Agrawal Associate Consultant,M.B.B.S,M.D. Pathology --2020

Page 5 of 7



Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name : MR MANISH KUMAR Age : 36 Yr(s) Sex :Male

Patient Episode: H03000060971Collection Date : 13 Mar 2024 10:15Referred By: HEALTH CHECK MHDReporting Date : 13 Mar 2024 16:53

Receiving Date : 13 Mar 2024 14:14

CLINICAL PATHOLOGY

| Test Name | Result | Biological Ref. Interval |
|---------------------------------------|-----------------------------|--------------------------|
| ROUTINE URINE ANALYSIS | | |
| MACROSCOPIC DESCRIPTION | | |
| Colour (Visual) | PALE YELLOW | (Pale Yellow - Yellow) |
| Appearance (Visual) | CLEAR | |
| CHEMICAL EXAMINATION | | |
| Reaction[pH] | 6.0 | (5.0-9.0) |
| (Reflectancephotometry(Indicator Meth | nod)) | |
| Specific Gravity | 1.015 | (1.003-1.035) |
| (Reflectancephotometry(Indicator Meth | nod)) | |
| Bilirubin | Negative | NEGATIVE |
| Protein/Albumin | Negative | (NEGATIVE-TRACE) |
| (Reflectance photometry(Indicator Met | hod)/Manual SSA) | |
| Glucose | NOT DETECTED | (NEGATIVE) |
| (Reflectance photometry (GOD-POD/Bene | edict Method)) | |
| Ketone Bodies | NOT DETECTED | (NEGATIVE) |
| (Reflectance photometry(Legal's Test) | /Manual Rotheras) | |
| Urobilinogen | NORMAL | (NORMAL) |
| Reflactance photometry/Diazonium salt | reaction | |
| Nitrite | NEGATIVE | NEGATIVE |
| Reflactance photometry/Griess test | | |
| Leukocytes | NIL | NEGATIVE |
| Reflactance photometry/Action of Este | erase | |
| BLOOD | NIL | NEGATIVE |
| (Reflectance photometry(peroxidase)) | | |
| MICROSCOPIC EXAMINATION (Manual) | Method: Light microscopy on | centrifuged urine |
| WBC/Pus Cells | 1-2 /hpf | (4-6) |
| Red Blood Cells | 1-2 /hpf | (1-2) |
| Epithelial Cells | 2-4 /hpf | (2-4) |
| Casts | NIL | (NIL) |
| Crystals | NIL | (NIL) |
| Bacteria | NIL | |
| Yeast cells | NIL | |

Page 6 of 7



Interpretation:

Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name : MR MANISH KUMAR Age 36 Yr(s) Sex :Male

Registration No : MH011770913 Lab No 38240301474

Patient Episode : H03000060971 **Collection Date:** 13 Mar 2024 10:15

: HEALTH CHECK MHD **Reporting Date:** 13 Mar 2024 16:53 **Referred By**

: 13 Mar 2024 14:14 **Receiving Date**

CLINICAL PATHOLOGY

URINALYSIS-Routine urine analysis assists in screening and diagnosis of various metabolic , urological, kidney and liver disorders

Protein: Elevated proteins can be an early sign of kidney disease. Urinary protein excretion can also be temporarily elevated by strenuous exercise, orthostatic proteinuria, dehydration, urina tract infections and acute illness with fever

Glucose: Uncontrolled diabetes mellitus can lead to presence of glucose in urine.

Other causes include pregnancy, hormonal disturbances, liver disease and certain medications.

Ketones: Uncontrolled diabetes mellitus can lead to presence of ketones in urine.

Ketones can also be seen in starvation, frequent vomiting, pregnancy and strenuous exercise.

Blood: Occult blood can occur in urine as intact erythrocytes or haemoglobin, which can occur in various urological, nephrological and bleeding disorders.

Leukocytes: An increase in leukocytes is an indication of inflammation in urinary tract or kidneys Most Common cause is bacterial urinary tract infection.

Nitrite: Many bacteria give positive results when their number is high. Nitrite concentration duri infection increases with length of time the urine specimen is retained in bladder prior to collection.

pH: The kidneys play an important role in maintaining acid base balance of the body. Conditions of the body producing acidosis/alkalosis or ingestion of certain type of food can affect the pH of urine.

Specific gravity: Specific gravity gives an indication of how concentrated the urine is. Increased Specific gravity is seen in conditions like dehydration, glycosuria and proteinuria while decrease Specific gravity is seen in excessive fluid intake, renal failure and diabetes insipidus.

Bilirubin: In certain liver diseases such as biliary obstruction or hepatitis, bilirubin gets excreted in urine.

Urobilinogen: Positive results are seen in liver diseases like hepatitis and cirrhosis

and in case of hemolytic anemia.

Page 7 of 7 -----END OF REPORT------

> Dr. Shalakha Agrawal Associate Consultant, M.B.B.S, M.D. Pathology --2020



Sector-6, Dwarka, New Delhi 110 075



GST: 07AAAAH3917LIZM PAN NO: AAAAH3917L

| NAME | MR Manish KUMAR | STUDY DATE | 13/03/2024 11:26AM |
|---------------|-------------------|--------------|--------------------|
| AGE / SEX | 36 y / M | HOSPITAL NO. | MH011770913 |
| ACCESSION NO. | R7045474 | MODALITY | US |
| REPORTED ON | 13/03/2024 3:10PM | REFERRED BY | Health Check MHD |

USG WHOLE ABDOMEN

Results:

Liver is normal in size (13.8 cm) and shows grade I fatty changes. No focal intra-hepatic lesion is detected. Intra-hepatic biliary radicals are not dilated. Portal vein is normal in calibre.

Gall bladder appears echofree with normal wall thickness. Common bile duct is normal in calibre.

Pancreas is normal in size and echopattern.

Spleen is normal in size (7.9 cm) and echopattern.

Both kidneys are normal in position, size and outline. Cortico-medullary differentiation of both kidneys is maintained. No focal lesion or calculus seen on either side. Bilateral pelvicalyceal systems are not dilated.

Urinary bladder is normal in wall thickness with clear contents. No significant intra or extraluminal mass is seen.

Prostate is normal in size, shape and echopattern. It measures 19.5 cc in volume.

No significant free fluid is detected.

IMPRESSION: USG findings are suggestive of grade I fatty liver.

Kindly correlate clinically

Dr. Nipun Gumber MBBS, MD DMC No.90272

ASSOCIATE CONSULTANT

*****End Of Report*****











Awarded Emergency Excellence Services MC/3228/04/09/2019-03/09/2021 E-2019-0026/27/07/2019-26/07/2021

Awarded Nursing Excellence Services N-2019-0113/27/07/2019-26/07/2021 IND18.6278/05/12/2018-04/12/2019