

(MEDIWHEEL) - BOB BELOW 40 MALE

CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST **DELHI**

NEW DELHI 110030

8800465156

ACCESSION NO: 0290WG002974 PATIENT ID

: SHRIM11068327

CHIENT BATIENT ID:

AGE/SEX :40 Years DRAWN

RECEIVED: 17/07/2023 09:07:24 REPORTED :17/07/2023 16:06:15

Test Report Status Results Biological Reference Interval Units **Final**

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

XRAY-CHEST

BOTH THE LUNG FIELDS ARE CLEAR

BOTH THE COSTOPHRENIC AND CARIOPHRENIC ANGELS ARE CLEAR

BOTH THE HILA ARE NORMAL **»**»

CARDIAC AND AORTIC SHADOWS APPEAR NORMAL BOTH THE DOMES OF THE DIAPHRAM ARE NORMAL **»**»

VISUALIZED BONY THORAX IS NORMAL >> >>

NO ABNORMALITY DETECTED. **IMPRESSION**

> Dr G.S. Saluja, (MBBS, DMRD) (Consultant Radiologist)

ECG

FCG

SUPRAVENTRICULAR ARRHYTHMIA.

OTHERWISE NORMAL ECG.

NOT SIGNIFICANT

MEDICAL HISTORY

RELEVANT PRESENT HISTORY

RELEVANT PAST HISTORY

P/H/O:-DM/HTN/ASTHMA. NOT SIGNIFICANT RELEVANT PERSONAL HISTORY RELEVANT FAMILY HISTORY NOT SIGNIFICANT NOT SIGNIFICANT OCCUPATIONAL HISTORY HISTORY OF MEDICATIONS NOT SIGNIFICANT

ANTHROPOMETRIC DATA & BMI

HEIGHT IN METERS 1.63 mts WEIGHT IN KGS. 64 Kgs

BMI 24 BMI & Weight Status as follows/sqmts

> Below 18.5: Underweight 18.5 - 24.9: Normal 25.0 - 29.9: Overweight 30.0 and Above: Obese

GENERAL EXAMINATION

Dr. Arpita Pasari, MD **Consultant Pathologist**





Page 1 Of 25

View Report





(MEDIWHEEL) - BOB BELOW 40 MALE

CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST **DELHI**

NECK

NEW DELHI 110030 8800465156

ACCESSION NO: 0290WG002974

PATIENT ID : SHRIM11068327

CHIENT BATIENT ID:

AGE/SEX :40 Years

DRAWN

RECEIVED: 17/07/2023 09:07:24 REPORTED :17/07/2023 16:06:15

Results **Biological Reference Interval Test Report Status** Units **Final**

MENTAL / EMOTIONAL STATE **NORMAL** NORMAL PHYSICAL ATTITUDE GENERAL APPEARANCE / NUTRITIONAL HEALTHY **STATUS**

BUILT / SKELETAL FRAMEWORK **AVERAGE** FACIAL APPEARANCE **NORMAL NORMAL** SKIN **NORMAL** UPPER LIMB NORMAL LOWER LIMB **NORMAL**

NOT ENLARGED OR TENDER NECK LYMPHATICS / SALIVARY GLANDS

THYROID GLAND **NOT ENLARGED**

CAROTID PULSATION **NORMAL TEMPERATURE AFEBRILE**

PULSE 65/MIN, REGULAR, ALL PERIPHERAL PULSES WELL FELT, NO CAROTID

BRUIT

RESPIRATORY RATE NORMAL

CARDIOVASCULAR SYSTEM

118/72 MM HG mm/Hg BP

> (SUPINE) **NORMAL** NORMAL **NORMAL**

> > ABSENT

RESPIRATORY SYSTEM

NORMAL SIZE AND SHAPE OF CHEST SYMMETRICAL MOVEMENTS OF CHEST BREATH SOUNDS INTENSITY **NORMAL**

BREATH SOUNDS QUALITY VESICULAR (NORMAL)

ADDED SOUNDS **ABSENT**

PER ABDOMEN

PERICARDIUM

APEX BEAT **HEART SOUNDS**

MURMURS

APPEARANCE NORMAL ABSENT VENOUS PROMINENCE

Dr. Arpita Pasari, MD **Consultant Pathologist**





Page 2 Of 25

View Report





(MEDIWHEEL) - BOB BELOW 40 MALE

CODE/NAME & ADDRESS : C000138355
ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: 0290WG002974

PATIENT ID : SHRIM11068327

CHIENT BATIENT ID:

AGE/SEX :40 Years N

DRAWN :

RECEIVED :17/07/2023 09:07:24 REPORTED :17/07/2023 16:06:15

Test Report Status Final Results Biological Reference Interval Units

LIVER NOT PALPABLE
SPLEEN NOT PALPABLE
HERNIA ABSENT

CENTRAL NERVOUS SYSTEM

HIGHER FUNCTIONS

CRANIAL NERVES

CEREBELLAR FUNCTIONS

SENSORY SYSTEM

MOTOR SYSTEM

REFLEXES

NORMAL

NORMAL

NORMAL

MUSCULOSKELETAL SYSTEM

SPINE NORMAL JOINTS NORMAL

BASIC EYE EXAMINATION

CONJUNCTIVA NORMAL
EYELIDS NORMAL
EYE MOVEMENTS NORMAL
CORNEA NORMAL

DISTANT VISION RIGHT EYE WITHOUT 6/6, WITHIN NORMAL LIMIT

GLASSES

DISTANT VISION LEFT EYE WITHOUT 6/6, WITHIN NORMAL LIMIT

GLASSES

NEAR VISION RIGHT EYE WITHOUT GLASSES

N6, WITHIN NORMAL LIMIT

N6, WITHIN NORMAL LIMIT

N6, WITHIN NORMAL LIMIT

NORMAL

COLOUR VISION
BASIC ENT EXAMINATION

EXTERNAL EAR CANAL NORMAL TYMPANIC MEMBRANE NORMAL

NOSE NO ABNORMALITY DETECTED

SINUSES NORMAL
THROAT NORMAL
TONSILS NOT ENLARGED

BASIC DENTAL EXAMINATION

Popila

Dr.Arpita Pasari, MD Consultant Pathologist





Page 3 Of 25

View Details

View Report





(MEDIWHEEL) - BOB BELOW 40 MALE

CODE/NAME & ADDRESS : C000138355

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO : 0290WG002974

PATIENT ID : SHRIM11068327

CHIENT BATIENT ID:

AGE/SEX :40 Years

DRAWN :

RECEIVED :17/07/2023 09:07:24 REPORTED :17/07/2023 16:06:15

Test Report Status Final Results Biological Reference Interval Units

TEETH NORMAL GUMS HEALTHY

SUMMARY

RELEVANT HISTORY NOT SIGNIFICANT RELEVANT GP EXAMINATION FINDINGS OVERWEIGHT

REMARKS / RECOMMENDATIONS NONE

FITNESS STATUS

FITNESS STATUS FIT (WITH MEDICAL ADVICE) (AS PER REQUESTED PANEL OF TESTS)

Comments

CLINICAL FINDINGS:-

RAISED SUGAR

RAISED HbA1C AND ESTIMATED AVERAG GLUCOSE (EAG)

DYSLIPIDEMIA.

OVER WEIGHT STATUS.

USG ABDOMEN SHOWS. :- EARLY FATTY INFILTRATION OF LIVER.

- THERE IS DEFECT 1.2 CM SEEN IN UMBLICAL REGION WITH
- REDUCIBLE CONTENT- HERNIA.

FITNESS STATUS :-

FITNESS STATUS: FIT (WITH MEDICAL ADVICE) (AS PER REQUESTED PANEL OF TESTS)

ADVICE: WEIGHT REDUCTION, LOW FAT& CARBOHYDRATE DIET AND REGULAR PHYSICAL EXERCISE FOR OVERWEIGHT STATUS AND DYSLIPIDEMIA.

NEED PHYSICIAN CONSULTATION FOR LIFE STYLE MODIFICATION.

Dr.Arpita Pasari, MD Consultant Pathologist



Page 4 Of 25

View Details







(MEDIWHEEL) - BOB BELOW 40 MALE

CODE/NAME & ADDRESS : C000138355
ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: **0290WG002974**PATIENT ID : SHRIM11068327

PATIENT ID : SHRIM1106832

AGE/SEX :40 Years
DRAWN :

RECEIVED : 17/07/2023 09:07:24

REPORTED :17/07/2023 16:06:15

Test Report Status Final Results Units

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE ULTRASOUND ABDOMEN ULTRASOUND ABDOMEN

Comments

USG-

IMPRESSION- EARLY FATTY INFILTRATION OF LIVER.

- THERE IS DEFECT 1.2 CM SEEN IN UMBLICAL REGION WITH REDUCIBLE CONTENT- HERNIA.

TMT OR ECHO

NEGATIVE

Interpretation(s)

MEDIČAL

FITNESS STATUS-Conclusion on an individual's Fitness, which is commented upon mainly for Pre employment cases, is based on multi factorial findings and does not depend on any one single parameter. The final Fitness assigned to a candidate will depend on the Physician's findings and overall judgement on a case to case basis, details of the candidate's past and personal history as well as the comprehensiveness of the diagnostic panel which has been requested for .These are then further correlated with details

of the job under consideration to eventually fit the right man to the right job.

Basis the above, Agilus diagnostic classifies a candidate's Fitness Status into one of the following categories:

- Fit (As per requested panel of tests) AGILUS Limited gives the individual a clean chit to join the organization, on the basis of the General Physical Examination and the specific test panel requested for.
- Fit (with medical advice) (As per requested panel of tests) This indicates that although the candidate can be declared as FIT to join the job, minimal problems have been detected during the Pre- employment examination. Examples of conditions which could fall in this category could be cases of mild reversible medical abnormalities such as height disproportions, borderline raised Blood Pressure readings, mildly raised Blood sugar and Blood Lipid levels, Hematuria, etc. Most of these relate to sedentary lifestyles and come under the broad category of life style disorders. The idea is to caution an individual to bring about certain lifestyle changes as well as seek a Physician"""s consultation and counseling in order to bring back to normal the mildly deranged parameters. For all purposes the individual is FIT to join the job.
- Fitness on Hold (Temporary Unfit) (As per requested panel of tests) Candidate's reports are kept on hold when either the diagnostic tests or the physical findings reveal the presence of a medical condition which warrants further tests, counseling and/or specialist opinion, on the basis of which a candidate can either be placed into Fit, Fit (With Medical Advice), or Unfit category. Conditions which may fall into this category could be high blood pressure, abnormal ECG, heart murmurs, abnormal vision, grossly elevated blood sugars, etc.
- Unfit (As per requested panel of tests) An unfit report by Agilus diagnostic Limited clearly indicates that the individual is not suitable for the respective job profile e.g. total color blindness in color related jobs.

Dr.Arpita Pasari, M

Dr.Arpita Pasari, MD Consultant Pathologist





Page 5 Of 25

/iew Details

View Report







(MEDIWHEEL) - BOB BELOW 40 MALE

CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI **NEW DELHI 110030** 8800465156

ACCESSION NO: 0290WG002974 PATIENT ID : SHRIM11068327

CHENT BATTENT ID:

AGE/SEX :40 Years

DRAWN

RECEIVED: 17/07/2023 09:07:24 REPORTED :17/07/2023 16:06:15

Test Report Status Results **Biological Reference Interval Units Final**

н	AEMATOLOGY - CBC		
MEDI WHEEL FULL BODY HEALTH CHECK UP B	ELOW 40 MALE		
BLOOD COUNTS,EDTA WHOLE BLOOD			
HEMOGLOBIN (HB) METHOD: SPECTROPHOTOMETRY	14.8	13.0 - 17.0	g/dL
RED BLOOD CELL (RBC) COUNT METHOD: ELECTRICAL IMPEDANCE	4.88	4.5 - 5.5	mil/μL
WHITE BLOOD CELL (WBC) COUNT METHOD: ELECTRICAL IMPEDANCE	8.50	4.0 - 10.0	thou/µL
PLATELET COUNT METHOD: ELECTRICAL IMPEDANCE	241	150 - 410	thou/µL
RBC AND PLATELET INDICES			
HEMATOCRIT (PCV) METHOD: CALCULATED	43.9	40 - 50	%
MEAN CORPUSCULAR VOLUME (MCV) METHOD: CALCULATED	90.0	83 - 101	fL
MEAN CORPUSCULAR HEMOGLOBIN (MCH) METHOD: CALCULATED	30.4	27.0 - 32.0	pg
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION (MCHC) METHOD: CALCULATED	33.8	31.5 - 34.5	g/dL
RED CELL DISTRIBUTION WIDTH (RDW) METHOD: CALCULATED	11.9	11.6 - 14.0	%
MENTZER INDEX	18.4		
MEAN PLATELET VOLUME (MPV)	9.3	6.8 - 10.9	fL
METHOD : CALCULATED			
WBC DIFFERENTIAL COUNT			
NEUTROPHILS METHOD: IMPEDANCE / MICROSCOPY	64	40 - 80	%
LYMPHOCYTES METHOD: IMPEDANCE / MICROSCOPY	28	20 - 40	%
MONOCYTES METHOD: IMPEDANCE / MICROSCOPY	06	2 - 10	%
EOSINOPHILS	02	1 - 6	%

Dr.Arpita Pasari, MD **Consultant Pathologist**





Page 6 Of 25

View Report





(MEDIWHEEL) - BOB BELOW 40 MALE

CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST **DELHI**

NEW DELHI 110030 8800465156

ACCESSION NO: 0290WG002974 :40 Years PATIENT ID DRAWN : SHRIM11068327

CHIENT BATIENT ID:

RECEIVED: 17/07/2023 09:07:24 REPORTED :17/07/2023 16:06:15

		i	
Test Report Status <u>Final</u>	Results	Biological Reference	e Interval Units
METHOD : IMPEDANCE / MICROSCOPY			
BASOPHILS METHOD: IMPEDANCE / MICROSCOPY	00	0 - 2	%
ABSOLUTE NEUTROPHIL COUNT METHOD: CALCULATED	5.44	2.0 - 7.0	thou/μL
ABSOLUTE LYMPHOCYTE COUNT METHOD: CALCULATED	2.38	1.0 - 3.0	thou/μL
ABSOLUTE MONOCYTE COUNT METHOD: CALCULATED	0.51	0.2 - 1.0	thou/μL
ABSOLUTE EOSINOPHIL COUNT METHOD: CALCULATED	0.17	0.02 - 0.50	thou/μL

Interpretation(s)
BLOOD COUNTS,EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading

RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients

A.-P. Yang, et al. International Immunopharmacology 84 (2020) 106504

This ratio element is a calculated parameter and out of NABL scope.

Dr. Arpita Pasari, MD **Consultant Pathologist**





Page 7 Of 25





PATIENT ID

CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0290WG002974

: SHRIM11068327

CHIENT BATIENT ID:

AGE/SEX :40 Years

DRAWN

RECEIVED: 17/07/2023 09:07:24 REPORTED :17/07/2023 16:06:15

Results Biological Reference Interval Units **Test Report Status Final**

HAEMATOLOGY

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE

BLOOD 80

E.S.R

0 - 14

mm at 1 hr

METHOD: MODIFIED WESTERGREN

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE **BLOOD**

9.7 High HBA1C

Non-diabetic: < 5.7 Pre-diabetics: 5.7 - 6.4 Diabetics: > or = 6.5Therapeutic goals: < 7.0 Action suggested: > 8.0

(ADA Guideline 2021)

METHOD: HPLC TECHNOLOGY


ESTIMATED AVERAGE GLUCOSE(EAG)

231.7 High

< 116.0

mg/dL

%



Interpretation(s)

Page 8 Of 25











(MEDIWHEEL) - BOB BELOW 40 MALE

CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0290WG002974 PATIENT ID

: SHRIM11068327

CHENT BATTENT ID:

AGE/SEX: 40 Years

DRAWN

RECEIVED: 17/07/2023 09:07:24 REPORTED: 17/07/2023 16:06:15

Test Report Status Results Biological Reference Interval **Final** Units

ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD-TEST DESCRIPTION :-

Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change TEST INTERPRETATION

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy, Estrogen medication, Aging

Earloger information, againgth and in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias, Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis).

In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum. Decreased in: Polycythermia vera, Sickle cell anemia

LIMITATIONS

False elevated ESR: Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia False Decreased: Poikilocytosis,(SickleCells,spherocytes),Microcytosis, Low fibrinogen, Very high WBC counts, Drugs(Quinine,

salicylates)

REFERENCE:

- 1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis,10th edition. GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-**Used For**:
- 1. Evaluating the long-term control of blood glucose concentrations in diabetic patients.
- 2. Diagnosing diabetes.

3. Identifying patients at increased risk for diabetes (prediabetes).
The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for well-controlled type 2 diabetic patients) to determine whether a patients metabolic control has remained continuously within the target range.

- eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.
 eAG gives an evaluation of blood glucose levels for the last couple of months.
- 3. eAG is calculated as eAG (mg/dl) = 28.7 * HbA1c 46.7

HbA1c Estimation can get affected due to :

- 1. Shortened Erythrocyte survival: Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss, hemolytic anemia) will falsely lower HbA1c test résults. Fructosamine is recommended in these patients which indicatés diabetes control over 15 days.
- 2.Vitamin C & E are reported to falsely lower test results.(possibly by inhibiting glycation of hemoglobin.
 3. Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates
- addiction are reported to interfere with some assay methods, falsely increasing results.

 4. Interference of hemoglobinopathies in HbA1c estimation is seen in
- a) Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c. b) Heterozygous state detected (D10 is corrected for HbS & HbC trait.)
- c) HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c. Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy

Page 9 Of 25





View Report





(MEDIWHEEL) - BOB BELOW 40 MALE

CODE/NAME & ADDRESS : C000138355
ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: 0290WG002974

PATIENT ID : SHRIM11068327

CHIENT BATTENT ID:

AGE/SEX :40 Years

DRAWN : RECEIVED : 17/07/2023 09:07:24

REPORTED :17/07/2023 16:06:15

Test Report Status Final Results Biological Reference Interval Units

IMMUNOHAEMATOLOGY

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

ABO GROUP TYPE AB

METHOD: TUBE AGGLUTINATION

RH TYPE POSITIVE

METHOD: TUBE AGGLUTINATION

Interpretation(s)

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same."

The test is performed by both forward as well as reverse grouping methods.

Proite

Dr.Arpita Pasari, MD Consultant Pathologist





Page 10 Of 25

View Details

View Report





(MEDIWHEEL) - BOB BELOW 40 MALE

CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0290WG002974 PATIENT ID : SHRIM11068327

CHIENT BATIENT ID:

AGE/SEX :40 Years DRAWN

RECEIVED: 17/07/2023 09:07:24 REPORTED :17/07/2023 16:06:15

Results Biological Reference Interval Units **Test Report Status Final**

BIOCHEMISTRY

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

GLUCOSE FASTING, FLUORIDE PLASMA


FBS (FASTING BLOOD SUGAR)

159 High

74 - 99

mg/dL

METHOD: HEXOKINASE



GLUCOSE, POST-PRANDIAL, PLASMA

PPBS(POST PRANDIAL BLOOD SUGAR)

222 High

Normal: < 140,

Impaired Glucose Tolerance: 140-199

Diabetic > or = 200

METHOD: HEXOKINASE

LIPID PROFILE, SERUM

CHOLESTEROL, TOTAL

212 High

Desirable: <200

mg/dL

mg/dL

BorderlineHigh: 200-239

High: > or = 240

METHOD: OXIDASE, ESTERASE, PEROXIDASE

Dr. Arpita Pasari, MD **Consultant Pathologist**





Page 11 Of 25



Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001 Madhya Pradesh, India Tel: 0731 2490008



(MEDIWHEEL) - BOB BELOW 40 MALE AGE/SEX :40 Years Male

CODE/NAME & ADDRESS: C000138355

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

PATIENT ID : SHRIM11068327

ACCESSION NO: 0290WG002974

DRAWN :

DELHI NEW DELHI 110030 8800465156 RECEIVED :17/07/2023 09:07:24 REPORTED :17/07/2023 16:06:15

	İ	I .
Test Report Status <u>Final</u>	Results	Biological Reference Interval Units
TRIGLYCERIDES	102	Desirable: < 150 mg/dL Borderline High: 150 - 199 High: 200 - 499 Very High: > or = 500
METHOD: ENZYMATIC ASSAY		, , , , , , , , , , , , , , , , , , ,
HDL CHOLESTEROL METHOD: DIRECT- NON IMMUNOLOGICAL	46	< 40 Low mg/dL > or = 60 High
CHOLESTEROL LDL	146 High	Adult levels: mg/dL Optimal < 100 Near optimal/above optimal: 100-129 Borderline high: 130-159 High: 160-189 Very high: = 190
NON HDL CHOLESTEROL METHOD: CALCULATED	166 High	Desirable: Less than 130 mg/dL Above Desirable: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very high: > or = 220
VERY LOW DENSITY LIPOPROTEIN METHOD: CALCULATED	20.4	< or = 30 mg/dL
CHOL/HDL RATIO	4.6 High	3.3 - 4.4
LDL/HDL RATIO	3.2 High	0.5 - 3.0 Desirable/Low Risk 3.1 - 6.0 Borderline/Moderate Risk >6.0 High Risk



Dr.Arpita Pasari, MD Consultant Pathologist





Page 12 Of 25

View Details







CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0290WG002974 PATIENT ID : SHRIM11068327


CPIENT BATIENT ID:


AGE/SEX :40 Years

DRAWN

RECEIVED: 17/07/2023 09:07:24 REPORTED :17/07/2023 16:06:15

Test Report Status Results Biological Reference Interval Units **Final**





Dr.Arpita Pasari, MD **Consultant Pathologist**



Page 13 Of 25

Patient Ref. No. 775000003935792



(MEDIWHEEL) - BOB BELOW 40 MALE ACCESSION NO: 0290WG002974

DRAWN

CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

PATIENT ID : SHRIM11068327 AGE/SEX :40 Years


DELHI

NEW DELHI 110030 8800465156

CHIENT BATTENT ID:

RECEIVED: 17/07/2023 09:07:24 REPORTED :17/07/2023 16:06:15

Test Report Status Results Biological Reference Interval Units **Final**



Interpretation(s)

Serum lipid profile is measured for cardiovascular risk prediction. Lipid Association of India recommends LDL-C as primary target and Non HDL-C as co-primary treatment target.

Risk Stratification for ASCVD (Atherosclerotic cardiovascular disease) by Lipid Association of India

Risk Category	,		
Extreme risk group	A.CAD with > 1 feature of high risk group		
	B. CAD with > 1 feature of Very high risk g	group or recurrent ACS (within 1 year) despite LDL-C < or =	
	50 mg/dl or polyvascular disease		
Very High Risk	1. Established ASCVD 2. Diabetes with 2 1	najor risk factors or evidence of end organ damage 3.	
	Familial Homozygous Hypercholesterolemi	a	
High Risk	1. Three major ASCVD risk factors. 2. Diabetes with 1 major risk factor or no evidence of end organ		
	damage. 3. CKD stage 3B or 4. 4. LDL >190 mg/dl 5. Extreme of a single risk factor. 6. Coronary		
	Artery Calcium - CAC >300 AU. 7. Lipoprotein a >/= 50mg/dl 8. Non stenotic carotid plaque		
Moderate Risk	2 major ASCVD risk factors		
Low Risk	0-1 major ASCVD risk factors		
Major ASCVD (Ath	Major ASCVD (Atherosclerotic cardiovascular disease) Risk Factors		
1. Age $>$ or $=$ 45 year	= 45 years in males and > or = 55 years in females 3. Current Cigarette smoking or tobacco use		
2. Family history of p	f premature ASCVD 4. High blood pressure		
5. Low HDL			
	<u> </u>	<u> </u>	

Newer treatment goals and statin initiation thresholds based on the risk categories proposed by LAI in 2020.

Dr. Arpita Pasari, MD **Consultant Pathologist**





Page 14 Of 25

View Report





CPIENT BALIENT ID:

(MEDIWHEEL) - BOB BELOW 40 MALE AGE/SEX :40 Years

CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0290WG002974 PATIENT ID

: SHRIM11068327

DRAWN

RECEIVED: 17/07/2023 09:07:24

REPORTED :17/07/2023 16:06:15

Test Report Status Final Results

Biological Reference Interval Units

Risk Group	Treatment Goals		Consider Drug T	herapy
	LDL-C (mg/dl)	Non-HDL (mg/dl)	LDL-C (mg/dl)	Non-HDL (mg/dl)
Extreme Risk Group Category A	<50 (Optional goal	< 80 (Optional goal	>OR = 50	>OR = 80
	$\langle OR = 30 \rangle$	<OR = 60)		
Extreme Risk Group Category B	<or 30<="" =="" td=""><td><OR = 60</td><td>> 30</td><td>>60</td></or>	<OR = 60	> 30	>60
Very High Risk	<50	<80	>OR= 50	>OR= 80
High Risk	<70	<100	>OR= 70	>OR= 100
Moderate Risk	<100	<130	>OR= 100	>OR= 130
Low Risk	<100	<130	>OR= 130*	>OR= 160

^{*}After an adequate non-pharmacological intervention for at least 3 months.

References: Management of Dyslipidaemia for the Prevention of Stroke: Clinical Practice Recommendations from the Lipid Association of India. Current Vascular Pharmacology, 2022, 20, 134-155.

LIVER FUNCTION PROFILE, SERUM

BILIRUBIN, TOTAL METHOD: JENDRASSIK AND GROFF	1.05	0.0 - 1.2	mg/dL
BILIRUBIN, DIRECT METHOD: DIAZOTIZATION	0.38 High	0.0 - 0.2	mg/dL
BILIRUBIN, INDIRECT METHOD: CALCULATED	0.67	0.00 - 1.00	mg/dL
TOTAL PROTEIN METHOD: BIURET	8.1	6.4 - 8.3	g/dL
ALBUMIN METHOD: BROMOCRESOL GREEN	5.1	3.50 - 5.20	g/dL
GLOBULIN METHOD: CALCULATED	3	2.0 - 4.1	g/dL
ALBUMIN/GLOBULIN RATIO METHOD: CALCULATED	1.7	1.0 - 2.0	RATIO
ASPARTATE AMINOTRANSFERASE(AST/SGOT) METHOD: UV WITH P5P	15	UPTO 40	U/L
ALANINE AMINOTRANSFERASE (ALT/SGPT) METHOD: UV WITH P5P	9	UP TO 45	U/L
ALKALINE PHOSPHATASE METHOD: PNPP	70	40 - 129	U/L
GAMMA GLUTAMYL TRANSFERASE (GGT) METHOD: G-GLUTAMYL-CARBOXY-NITROANILIDE	25	8 - 61	U/L
LACTATE DEHYDROGENASE METHOD: ENZYMATIC LACTATE - PYRUVATE(IFCC)	189	135 - 225	U/L

Dr.Arpita Pasari, MD **Consultant Pathologist**



Page 15 Of 25

View Report





CODE/NAME & ADDRESS: C000138355 ACCESSION NO: 0290WG002974 AGE/SEX :40 Years ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

PATIENT ID : SHRIM11068327

CPIENT BALIENT ID:

DRAWN


RECEIVED: 17/07/2023 09:07:24 REPORTED :17/07/2023 16:06:15

Test Report Status Results **Biological Reference Interval Units Final**

BLOOD UREA NITROGEN (BUN), SERUM

BLOOD UREA NITROGEN 7 6 - 20mg/dL

METHOD: UREASE KINETIC



CREATININE, SERUM

mg/dL **CREATININE** 0.74 0.70 - 1.20

METHOD: ALKALINE PICRATE KINETIC JAFFES

Dr.Arpita Pasari, MD **Consultant Pathologist** Page 16 Of 25













CODE/NAME & ADDRESS : C000138355

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL
F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: 0290WG002974

: SHRIM11068327

SPIENT BATIENT ID:


PATIENT ID

AGE/SEX :40 Years Mal

DRAWN :

RECEIVED : 17/07/2023 09:07:24 REPORTED :17/07/2023 16:06:15

Test Report Status <u>Final</u> Results Biological Reference Interval Units





Dr.Arpita Pasari, MD Consultant Pathologist





Page 17 Of 25

View Details







(MEDIWHEEL) - BOB BELOW 40 MALE

CODE/NAME & ADDRESS : C000138355

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL
E-703 LADO SARAI MEHRAUI ISOUTH WEST

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: **0290WG002974**PATIENT ID : SHRIM11068327

CHENT BATTENT ID:

AGE/SEX :40 Years
DRAWN :

RECEIVED : 17/07/2023 09:07:24 REPORTED :17/07/2023 16:06:15

Test Report Status <u>Final</u> Results Biological Reference Interval Units

ELECTROLYTES (NA/K/CL), SERUM

SODIUM, SERUM 143.4 136.0 - 146.0 mmol/L

METHOD: DIRECT ION SELECTIVE ELECTRODE

POTASSIUM, SERUM 4.10 3.50 - 5.10 mmol/L

METHOD: DIRECT ION SELECTIVE ELECTRODE

CHLORIDE, SERUM 103.8 98.0 - 106.0 mmol/L

METHOD: DIRECT ION SELECTIVE ELECTRODE

Interpretation(s)

Sodium	Potassium	Chloride
Decreased in:CCF, cirrhosis, vomiting, diarrhea, excessive sweating, salt-losing nephropathy, adrenal insufficiency, nephrotic syndrome, water intoxication, SIADH. Drugs: thiazides, diuretics, ACE inhibitors, chlorpropamide, carbamazepine, anti depressants (SSRI), antipsychotics.	Decreased in: Low potassium intake, prolonged vomiting or diarrhea, RTA types I and II, hyperaldosteronism, Cushing's syndrome, osmotic diuresis (e.g., hyperglycemia), alkalosis, familial periodic paralysis, trauma (transient). Drugs: Adrenergic agents, diuretics.	Decreased in: Vomiting, diarrhea, renal failure combined with salt deprivation, over-treatment with diuretics, chronic respiratory acidosis, diabetic ketoacidosis, excessive sweating, SIADH, salt-losing nephropathy, porphyria, expansion of extracellular fluid volume, adrenalinsufficiency, hyperaldosteronism, metabolic alkalosis. Drugs: chronic laxative, corticosteroids, diuretics.
Increased in: Dehydration (excessivesweating, severe vomiting or diarrhea), diabetes mellitus, diabetesinsipidus, hyperaldosteronism, inadequate water intake. Drugs: steroids, licorice, oral contraceptives.	Increased in: Massive hemolysis, severe tissue damage, rhabdomyolysis, acidosis, dehydration,renal failure, Addison's disease, RTA type IV, hyperkalemic familial periodic paralysis. Drugs: potassium salts, potassium- sparing diuretics,NSAIDs, beta-blockers, ACE inhibitors, highdose trimethoprim-sulfamethoxazole.	Increased in: Renal failure, nephrotic syndrome, RTA, dehydration, overtreatment with saline, hyperparathyroidism, diabetes insipidus, metabolic acidosis from diarrhea (Loss of HCO3-), respiratory alkalosis, hyperadrenocorticism. Drugs: acetazolamide, androgens, hydrochlorothiazide, salicylates.
Interferences: Severe lipemia or hyperproteinemi, if sodium analysis involves a dilution step can cause spurious results. The serum sodium falls about 1.6 mEq/L for each 100 mg/dL increase in blood glucose.	Interferences: Hemolysis of sample, delayed separation of serum, prolonged fist clenching during blood drawing, and prolonged tourniquet placement. Very high WBC/PLT counts may cause spurious. Plasma potassium levels are normal.	Interferences:Test is helpful in assessing normal and increased anion gap metabolic acidosis and in distinguishing hypercalcemia due to hyperparathyroidism (high serum chloride) from that due to malignancy (Normal serum chloride)

Interpretation(s)

GLUCOSE FASTING, FLUORIDE PLASMA-TEST DESCRIPTION

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in the

Increased in:Diabetes mellitus, Cushing's syndrome (10 – 15%), chronic pancreatitis (30%). Drugs:corticosteroids,phenytoin, estrogen, thiazides.

Decreased in:Pancreatic islet cell disease with increased insulin,insulinoma,adrenocortical insufficiency,hypopituitarism,diffuse liver disease, malignancy(adrenocortical,stomach,fibrosarcoma),infant of a diabetic mother,enzyme deficiency diseases(e.g.galactosemia),Drugs-insulin,ethanol,propranolol sulfonylureas,tolbutamide,and other oral hypoglycemic agents.

Proite

Dr.Arpita Pasari, MD Consultant Pathologist



Page 18 Of 25

view Details

View Report





(MEDIWHEEL) - BOB BELOW 40 MALE

CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0290WG002974

PATIENT ID : SHRIM11068327

CHIENT BATTENT ID:

:40 Years AGE/SEX

DRAWN

RECEIVED: 17/07/2023 09:07:24

REPORTED: 17/07/2023 16:06:15

Test Report Status Results Biological Reference Interval Units **Final**

NOTE: While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values), there is wide fluctuation within individuals.Thus, glycosylated hemoglobin(HbA1c) levels are favored to monitor glycemic control.

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment,Renal Glyosuria,Glycaemic

index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.

GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.Additional test HbA1c LIVER FUNCTION PROFILE, SERUM-

Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. **Elevated levels** results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health.AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Pagets disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilsons disease.

GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive

liver disease,high alcohol consumption and use of enzyme-inducing drugs etc.

Total Protein also known as total protein,is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin.Higher-than-normal levels may be due to:Chronic inflammation or infection,including HIV and hepatitis B or C,Multiple myeloma,Waldenstroms disease.Lower-than-normal levels may be due to: Agammaglobulinemia,Bleeding (hemorrhage),Burns,Glomerulonephritis,Liver disease, Malabsorption,Malnutrition,Nephrotic syndrome, Protein-losing enteropathy etc.

Albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc

BLOOD UREA NITROGEN (BUN), SERUM-Causes of Increased levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)

Causes of decreased level include Liver disease, SIADH.

CREATININE, SERUM-Higher than normal level may be due to:

• Blockage in the urinary tract, Kidney problems, such as kidney damage or failure, infection, or reduced blood flow, Loss of body fluid (dehydration), Muscle problems, such as breakdown of muscle fibers, Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:

• Myasthenia Gravis, Muscuophy

LINE ACID SERUM Contract of Tearner and Involve Distance (high Problems David Mathelia)

URIC ACID, SERUM-Causes of Increased levels:-Dietary(High Protein Intake,Prolonged Fasting,Rapid weight loss),Gout,Lesch nyhan syndrome,Type 2 DM,Metabolic syndrome Causes of decreased levels-Low Zinc intake,OCP,Multiple Sclerosis

TOTAL PROTEIN, SERUM-is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin.

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease.

Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

ALBUMIN, SERUM-Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.

Dr. Arpita Pasari, MD

Page 19 Of 25





View Report

Consultant Pathologist





(MEDIWHEEL) - BOB BELOW 40 MALE

CODE/NAME & ADDRESS : C000138355
ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: 0290WG002974

PATIENT ID : SHRIM11068327

CHIENT BATIENT ID:

AGE/SEX :40 Years

DRAWN : RECEIVED : 17/07/2023 09:07:24

REPORTED :17/07/2023 16:06:15

Test Report Status Final Results Biological Reference Interval Units

CLINICAL PATH - URINALYSIS

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

PHYSICAL EXAMINATION, URINE

COLOR PALE YELLOW APPEARANCE CLEAR

CHEMICAL EXAMINATION, URINE

PH	5.0	4.7 - 7.5
SPECIFIC GRAVITY	<=1.005	1.003 - 1.035
PROTEIN	NOT DETECTED	NOT DETECTED
GLUCOSE	NOT DETECTED	NOT DETECTED
KETONES	NOT DETECTED	NOT DETECTED
BLOOD	NOT DETECTED	NOT DETECTED
BILIRUBIN	NOT DETECTED	NOT DETECTED
UROBILINOGEN	NORMAL	NORMAL
NITRITE	NOT DETECTED	NOT DETECTED
LEUKOCYTE ESTERASE	NOT DETECTED	NOT DETECTED

MICROSCOPIC EXAMINATION, URINE

RED BLOOD CELLS	NOT DETECTED	NOT DETECTED	/HPF
PUS CELL (WBC'S)	2-3	0-5	/HPF
EPITHELIAL CELLS	2-3	0-5	/HPF

CASTS NOT DETECTED
CRYSTALS NOT DETECTED

BACTERIA NOT DETECTED NOT DETECTED
YEAST NOT DETECTED NOT DETECTED

REMARKS Please note that all the urinary findings are confirmed manually as well.

Interpretation(s)

The following table describes the probable conditions, in which the analytes are present in urine

Presence of	Conditions
Proteins	Inflammation or immune illnesses

Dr.Arpita Pasari, MD

Consultant Pathologist

Page 20 Of 25





View Details

View Repor





(MEDIWHEEL) - BOB BELOW 40 MALE

CODE/NAME & ADDRESS : C000138355

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: 0290WG002974

PATIENT ID : SHRIM11068327

CHENT BATTENT ID:

AGE/SEX :40 Years

Mal

DRAWN :

RECEIVED : 17/07/2023 09:07:24 REPORTED :17/07/2023 16:06:15

Test Report Status Final Results Biological Reference Interval Units

Pus (White Blood Cells)	Urinary tract infection, urinary tract or kidney stone, tumors or any kind of kidney impairment
Glucose	Diabetes or kidney disease
Ketones	Diabetic ketoacidosis (DKA), starvation or thirst
Urobilinogen	Liver disease such as hepatitis or cirrhosis
Blood	Renal or genital disorders/trauma
Bilirubin	Liver disease
Erythrocytes	Urological diseases (e.g. kidney and bladder cancer, urolithiasis), urinary tract infection and glomerular diseases
Leukocytes	Urinary tract infection, glomerulonephritis, interstitial nephritis either acute or chronic, polycystic kidney disease, urolithiasis, contamination by genital secretions
Epithelial cells	Urolithiasis, bladder carcinoma or hydronephrosis, ureteric stents or bladder catheters for prolonged periods of time
Granular Casts	Low intratubular pH, high urine osmolality and sodium concentration, interaction with Bence-Jones protein
Hyaline casts	Physical stress, fever, dehydration, acute congestive heart failure, renal diseases
Calcium oxalate	Metabolic stone disease, primary or secondary hyperoxaluria, intravenous infusion of large doses of vitamin C, the use of vasodilator naftidrofuryl oxalate or the gastrointestinal lipase inhibitor orlistat, ingestion of ethylene glycol or of star fruit (Averrhoa carambola) or its juice
Uric acid	arthritis
Bacteria	Urinary infectionwhen present in significant numbers & with pus cells.
Trichomonas vaginalis	Vaginitis, cervicitis or salpingitis

Dr.Arpita Pasari, MD



Page 21 Of 25

View Details

View Report

Consultant Pathologist





(MEDIWHEEL) - BOB BELOW 40 MALE

CODE/NAME & ADDRESS : C000138355
ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: 0290WG002974

PATIENT ID : SHRIM11068327

CHIENT BATIENT ID:

AGE/SEX :40 Years
DRAWN :

RECEIVED : 17/07/2023 09:07:24 REPORTED :17/07/2023 16:06:15

Test Report Status <u>Final</u> Results Biological Reference Interval Units

CLINICAL PATH - STOOL ANALYSIS

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

PHYSICAL EXAMINATION, STOOL

COLOUR BROWN

CONSISTENCY WELL FORMED

MUCUS NOT DETECTED NOT DETECTED

VISIBLE BLOOD ABSENT ABSENT

ADULT PARASITE NOT DETECTED

CHEMICAL EXAMINATION, STOOL

STOOL PH ALKALINE

OCCULT BLOOD NOT DETECTED NOT DETECTED

MICROSCOPIC EXAMINATION, STOOL

PUS CELLS 2-3 /hpf

RED BLOOD CELLS NOT DETECTED NOT DETECTED /HPF

CYSTS NOT DETECTED NOT DETECTED

OVA NOT DETECTED

LARVAE NOT DETECTED NOT DETECTED

TROPHOZOITES NOT DETECTED NOT DETECTED

FAT ABSENT
VEGETABLE CELLS ABSENT
CHARCOT LEYDEN CRYSTALS ABSENT

Interpretation(s)

Stool routine analysis is only a screening test for disorders of gastrointentestinal tract like infection, malabsorption, etc. The following table describes the probable conditions, in which the analytes are present in stool.

PRESENCE OF	CONDITION
Pus cells	Pus in the stool is an indication of infection
Red Blood cells	Parasitic or bacterial infection or an inflammatory bowel condition such as ulcerative colitis

- wint -

Dr.Meena Jinwah ,MBBS . MD Consultant Microbiologist



Dr.Arpita Pasari, MD Consultant Pathologist





Page 22 Of 25

View Details

View Report







(TIEDIWILLE) BOD BLEOW 101

CODE/NAME & ADDRESS: C000138355

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL
F-703, LADO SARAI, MEHRAULISOUTH WEST

PATIENT ID : SHRIM11068327

AGE/SEX :40 Years

DRAWN

DELHI

CHENT BATIENT ID:

ACCESSION NO: 0290WG002974

RECEIVED : 17/07/2023 09:07:24

NEW DELHI 110030 8800465156 REPORTED :17/07/2023 16:06:15

Test Report Status Final Results Biological Reference Interval Units

D '4	Tu Coution a Called Historia and a second Called American Course and American Course	
Parasites	Infection of the digestive system. Stool examination for ova and parasite detects	
	presence of parasitic infestation of gastrointestinal tract. Various forms of	
	parasite that can be detected include cyst, trophozoite and larvae. One negative	
	result does not rule out the possibility of parasitic infestation. Intermittent	
	shedding of parasites warrants examinations of multiple specimens tested on	
	consecutive days. Stool specimens for parasitic examination should be collected	
	before initiation of antidiarrheal therapy or antiparasitic therapy. This test does	
	not detect presence of opportunistic parasites like Cyclospora, Cryptosporidia	
	and Isospora species. Examination of Ova and Parasite has been carried out by	
	direct and concentration techniques.	
Mucus	Mucus is a protective layer that lubricates, protects& reduces damage due to	
	bacteria or viruses.	
Charcot-Leyden crystal	Parasitic diseases.	
Ova & cyst	Ova & cyst indicate parasitic infestation of intestine.	
Frank blood	Bleeding in the rectum or colon.	
Occult blood	Occult blood indicates upper GI bleeding.	
Macrophages	Macrophages in stool are an indication of infection as they are protective cells.	
Epithelial cells	Epithelial cells that normally line the body surface and internal organs show u	
	in stool when there is inflammation or infection.	
Fat	Increased fat in stool maybe seen in conditions like diarrhoea or malabsorption.	
рН	Normal stool pH is slightly acidic to neutral. Breast-fed babies generally have an acidic stool.	

ADDITIONAL STOOL TESTS:

- Stool Culture: This test is done to find cause of GI infection, make decision about best treatment for GI infection & to find out if treatment for GI infection worked.
- 2. <u>Fecal Calprotectin</u>: It is a marker of intestinal inflammation. This test is done to differentiate Inflammatory Bowel Disease (IBD) from Irritable Bowel Syndrome (IBS).
- 3. Fecal Occult Blood Test(FOBT): This test is done to screen for colon cancer & to evaluate possible cause of unexplained anaemia.
- 4. Clostridium Difficile Toxin Assay: This test is strongly recommended in healthcare associated bloody or waterydiarrhoea, due to overuse of broad spectrum antibiotics which alter the normal GI flora.
- 5. <u>Biofire (Film Array) GI PANEL</u>: In patients of Diarrhoea, Dysentry, Rice watery Stool, FDA approved, Biofire Film Array Test,(Real Time Multiplex PCR) is strongly recommended as it identifies organisms, bacteria, fungi, virus ,parasite and other opportunistic pathogens, Vibrio cholera infections only in 3 hours. Sensitivity 96% & Specificity 99%.
- 6. Rota Virus Immunoassay: This test is recommended in severe gastroenteritis in infants & children associated with watery diarrhoea, vomitting& abdominal cramps. Adults are also affected. It is highly contagious in nature.

- wind -

Dr.Meena Jinwah ,MBBS . MD Consultant Microbiologist Pepita

Dr.Arpita Pasari, MD Consultant Pathologist





Page 23 Of 25

View Details

View Report





(MEDIWHEEL) - BOB BELOW 40 MALE

CODE/NAME & ADDRESS : C000138355 ACCESSION NO: 0290WG002974 AGE/SEX :40 Years ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

PATIENT ID : SHRIM11068327

DRAWN

CHIENT BATIENT ID:

RECEIVED: 17/07/2023 09:07:24 REPORTED: 17/07/2023 16:06:15

Results **Biological Reference Interval Units Test Report Status Final**

SPECIALISED CHEMISTRY - HORMONE

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

THYROID PANEL, SERUM

ТЗ	95.68	80.0 - 200.0	ng/dL
METHOD: CHEMILUMINESCENCE TECHNOLOGY			
T4	7.66	5.10 - 14.10	μg/dL
METHOD: CHEMILUMINESCENCE TECHNOLOGY			
TSH (ULTRASENSITIVE)	1.690	0.270 - 4.200	μIU/mL

METHOD: CHEMILUMINESCENCE TECHNOLOGY

Interpretation(s)

Triiodothyronine T3, Thyroxine T4, and Thyroid Stimulating Hormone TSH are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low. Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3. Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, Free T4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Sr. No.	TSH	Total T4	FT4	Total T3	Possible Conditions
1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3)
					Post Thyroidectomy (4) Post Radio-Iodine treatment
2	High	Normal	Normal	Normal	(1)Subclinical Hypothyroidism (2) Patient with insufficient thyroid
					hormone replacement therapy (3) In cases of Autoimmune/Hashimoto
					thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical
					inflammation, drugs like amphetamines, Iodine containing drug and
					dopamine antagonist e.g. domperidone and other physiological reasons.
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre
					(3)Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid
					hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4
					replacement therapy (7) First trimester of Pregnancy
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism
6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor

Dr. Arpita Pasari, MD **Consultant Pathologist**



Page 24 Of 25

View Report



Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001 Madhya Pradesh, India





(MEDIWHEEL) - BOB BELOW 40 MALE AGE/SEX :40 Years

CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0290WG002974 PATIENT ID

: SHRIM11068327

CHENT BATTENT ID:

DRAWN RECEIVED: 17/07/2023 09:07:24

REPORTED :17/07/2023 16:06:15

Results **Test Report Status** Biological Reference Interval Units **Final**

7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent treatment for Hyperthyroidism
8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies

REF: 1. TIETZ Fundamentals of Clinical chemistry 2. Guidlines of the American Thyroid association during pregnancy and Postpartum, 2011. NOTE: It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

> **End Of Report** Please visit www.agilusdiagnostics.com for related Test Information for this accession

CONDITIONS OF LABORATORY TESTING & REPORTING

- 1. It is presumed that the test sample belongs to the patient named or identified in the test requisition form.
- 2. All tests are performed and reported as per the turnaround time stated in the AGILUS Directory of Services.
- 3. Result delays could occur due to unforeseen circumstances such as non-availability of kits / equipment breakdown / natural calamities / technical downtime or any other unforeseen event.
- 4. A requested test might not be performed if:
 - i. Specimen received is insufficient or inappropriate
 - ii. Specimen quality is unsatisfactory
 - iii. Incorrect specimen type
 - iv. Discrepancy between identification on specimen container label and test requisition form

- 5. AGILUS Diagnostics confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity.
- 6. Laboratory results should not be interpreted in isolation; it must be correlated with clinical information and be interpreted by registered medical practitioners only to determine final diagnosis.
- Test results may vary based on time of collection, physiological condition of the patient, current medication or nutritional and dietary changes. Please consult your doctor or call us for any clarification.
- Test results cannot be used for Medico legal purposes.
- 9. In case of queries please call customer care (91115 91115) within 48 hours of the report.

Agilus Diagnostics Ltd

Fortis Hospital, Sector 62, Phase VIII, Mohali 160062

Dr. Arpita Pasari, MD **Consultant Pathologist**





Page 25 Of 25



