: MediWheel

 PID No.
 : MED122500589
 Register On
 : 02/03/2024 10:16 AM

 SID No.
 : 522403552
 Collection On
 : 02/03/2024 1:40 PM

 Age / Sex
 : 37 Year(s) / Male
 Report On
 : 02/03/2024 6:48 PM

Printed On



Type : OP

Ref. Dr

RDW-SD

(EDTA Blood)

Neutrophils

Lymphocytes

Eosinophils

Monocytes

(Blood)

(Blood)

(Blood)

(Blood)

Total Leukocyte Count (TC)

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BLOOD GROUPING AND Rh TYPING (EDTA Blood/Agglutination)	'O' 'Positive'		
INTERPRETATION: Note: Slide method is scre	ening method. Kind	lly confirm with Tube method	od for transfusion.
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood/Spectrophotometry)	13.5	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	40.3	%	42 - 52
RBC Count (EDTA Blood)	4.54	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (EDTA Blood)	88.7	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood)	29.8	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood)	33.6	g/dL	32 - 36
RDW-CV	13.7	%	11.5 - 16.0

42.53

9100

61.1

31.6

0.7

6.0

fL

%

%

%

%

cells/cu.mm

: 03/03/2024 3:35 PM







39 - 46

4000 - 11000

40 - 75

20 - 45

01 - 06

01 - 10

APPROVED BY

: MediWheel

 PID No.
 : MED122500589
 Register On
 : 02/03/2024 10:16 AM

 SID No.
 : 522403552
 Collection On
 : 02/03/2024 1:40 PM

 Age / Sex
 : 37 Year(s) / Male
 Report On
 : 02/03/2024 6:48 PM

Printed On



Type : OP

Ref. Dr

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Basophils (Blood)	0.6	%	00 - 02
INTERPRETATION: Tests done on Automated	Five Part cell count	er. All abnormal results a	are reviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood)	5.56	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood)	2.88	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood)	0.06	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood)	0.55	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood)	0.05	10^3 / μl	< 0.2
Platelet Count (EDTA Blood)	400	10^3 / μl	150 - 450
MPV (Blood)	8.4	fL	7.9 - 13.7
PCT (Automated Blood cell Counter)	0.34	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood)	4	mm/hr	< 15
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	85.43	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125

: 03/03/2024 3:35 PM

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine) (Urine - F/GOD - POD)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	9.1	mg/dL	7.0 - 21
Creatinine (Serum/Modified Jaffe)	0.92	mg/dL	0.9 - 1.3







Diabetic: >= 126

The results pertain to sample tested.

 PID No.
 : MED122500589
 Register On
 : 02/03/2024 10:16 AM

 SID No.
 : 522403552
 Collection On
 : 02/03/2024 1:40 PM

 Age / Sex
 : 37 Year(s) / Male
 Report On
 : 02/03/2024 6:48 PM

Printed On



Type : OP

Ref. Dr : MediWheel

<u>Investigation</u>	Observed Unit	<u>Biological</u>
_	<u>Value</u>	Reference Interval

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine etc.

: 03/03/2024 3:35 PM

such as cefoxitin ,cefazolin, ACE inhibitors ,angiote etc.	nsin II receptor	antagonists,N-acetylcyteine,	, chemotherapeutic agent such as
Uric Acid (Serum/Enzymatic)	5.77	mg/dL	3.5 - 7.2
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	0.30	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.08	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.22	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	15.06	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	15.98	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	31.42	U/L	< 55
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	94.0	U/L	53 - 128
Total Protein (Serum/Biuret)	7.25	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.62	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	2.63	gm/dL	2.3 - 3.6
A: GRATIO (Serum/Derived)	1.76		1.1 - 2.2
<u>Lipid Profile</u>			







 PID No.
 : MED122500589
 Register On
 : 02/03/2024 10:16 AM

 SID No.
 : 522403552
 Collection On
 : 02/03/2024 1:40 PM

Ref. Dr : MediWheel



<u>Investigation</u>	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Cholesterol Total (Serum/CHOD-PAP with ATCS)	180.40	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	188.51	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual_circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	28.98	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	113.7	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	37.7	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	151.4	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2. It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.







The results pertain to sample tested.

 PID No.
 : MED122500589
 Register On
 : 02/03/2024 10:16 AM

 SID No.
 : 522403552
 Collection On
 : 02/03/2024 1:40 PM

Age / Sex : 37 Year(s) / Male Report On : 02/03/2024 6:48 PM

Ref. Dr : MediWheel



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Total Cholesterol/HDL Cholesterol Ratio (Serum/Calculated)	6.2		Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
Triglyceride/HDL Cholesterol Ratio (TG/HDL) (Serum/Calculated)	6.5		Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
LDL/HDL Cholesterol Ratio (Serum/Calculated)	3.9		Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/HPLC)	5.3	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 105.41 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 0.787 ng/ml 0.7 - 2.04 (Serum/*ECLIA*)







The results pertain to sample tested.

Page 5 of 10

 PID No.
 : MED122500589
 Register On
 : 02/03/2024 10:16 AM

 SID No.
 : 522403552
 Collection On
 : 02/03/2024 1:40 PM

Ref. Dr : MediWheel

MEDALL

<u>Investigation</u>	Observed Unit	<u>Biological</u>
	<u>Value</u>	Reference Interval

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Tyroxine) - Total 5.32 μg/dl 4.2 - 12.0

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 2.51 µIU/mL 0.35 - 5.50

(Serum/ECLIA)

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.

2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.

3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.

PHYSICAL EXAMINATION (URINE

<u>COMPLETE)</u>

Colour (Urine)	Pale yellow	Yellow to Amber
Appearance (Urine)	Clear	Clear
Volume(CLU) (Urine)	20	
CHEMICAL EXAMINATION	ON (URINE	

<u>CHEMICAL EXAMINATION (URINE</u> <u>COMPLETE)</u>

pH 6.5 4.5 - 8.0 (Urine)







APPROVED BY

The results pertain to sample tested.

Page 6 of 10

PID No. : MED122500589 : 522403552 SID No. Collection On : 02/03/2024 1:40 PM

Age / Sex : 37 Year(s) / Male Report On : 02/03/2024 6:48 PM

Type : OP

Ref. Dr

: MediWheel

Register On : 02/03/2024 10:16 AM

Printed On : 03/03/2024 3:35 PM



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Specific Gravity (Urine)	1.012		1.002 - 1.035
Ketone (Urine)	Negative		Negative
Urobilinogen (Urine)	Normal		Normal
Blood (Urine)	Negative		Negative
Nitrite (Urine)	Negative		Negative
Bilirubin (Urine)	Negative		Negative
Protein (Urine)	Negative		Negative
Glucose (Urine/GOD - POD)	Negative		Negative
Leukocytes(CP) (Urine)	Negative		
MICROSCOPIC EXAMINATION (URINE COMPLETE)			
Pus Cells (Urine)	0-1	/hpf	NIL
Epithelial Cells (Urine)	0-1	/hpf	NIL
RBCs (Urine)	NIL	/HPF	NIL
Others (Urine)	NIL		
INTERPRETATION: Note: Done with Autoreviewed and confirmed microscopically.	omated Urine Analyser	& Automated urine sedi	mentation analyser. All abnormal reports



NIL



/hpf



NIL

APPROVED BY

Casts (Urine)

PID No. : MED122500589

: 522403552 SID No.

Age / Sex : 37 Year(s) / Male

Type : OP

Register On : 02/03/2024 10:16 AM

Collection On : 02/03/2024 1:40 PM

: 02/03/2024 6:48 PM

Printed On : 03/03/2024 3:35 PM



Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Crystals	NIL	/hpf	NIL
(Urine)			



Report On





: MediWheel

PID No. : MED122500589

SID No. : 522403552

Age / Sex : 37 Year(s) / Male

BUN / Creatinine Ratio

Type : OP

Ref. Dr

ype . Or

Register On : 02/03/2024 10:16 AM

Collection On : 02/03/2024 1:40 PM

Report On : 02/03/2024 6:48 PM

Printed On : 03/03/2024 3:35 PM

9.8

MEI

Investigation Observed Unit Value

Biological Reference Interval

6.0 - 22.0





PID No. : MED122500589

: 522403552 SID No.

Age / Sex : 37 Year(s) / Male

Type : OP

Ref. Dr : MediWheel Register On : 02/03/2024 10:16 AM

Collection On : 02/03/2024 1:40 PM

Report On : 02/03/2024 6:48 PM

Printed On

: 03/03/2024 3:35 PM

<u>Observed</u> **Investigation** Value

<u>Unit</u>

Biological Reference Interval

URINE ROUTINE





-- End of Report --



Name	Mr.ARUN KUMAR P L	ID	MED122500589
Age & Gender	37/MALE	Visit Date	02/03/2024
Ref Doctor Name	MediWheel		

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in shape, size (12.1 cm) and has uniform echopattern. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER is partially distended. CBD is not dilated.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN shows normal shape, size and echopattern.

KIDNEYS

Right kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

Left kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney	11.4	1.4
Left Kidney	11.6	1.9

URINARY BLADDER is moderately distended. No evidence of diverticula.

PROSTATE: Visualized prostate appears normal.

No evidence of ascites.

IMPRESSION:

• No significant abnormality detected.

REPORT DISCLAIMER

- 1. This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- $11. Disputes, if any \ , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only. \\$



Name	Mr.ARUN KUMAR P L	ID	MED122500589
Age & Gender	37/MALE	Visit Date	02/03/2024
Ref Doctor Name	MediWheel		

DR. HEMANANDINI V.N CONSULTANT RADIOLOGIST Hn/Mi

REPORT DISCLAIMER

- 1.This is only a radiologincal imperssion.Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11. Disputes, if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



Name	Mr. ARUN KUMAR P L	ID	MED122500589
Age & Gender	37Y/M	Visit Date	Mar 2 2024 10:15AM
Ref Doctor	MediWheel		

X - RAY CHEST PA VIEW

Bilateral lung	fields a	ppear	normal.
----------------	----------	-------	---------

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

Impression:

• No significant abnormality detected.

DR.S.SHWETHA.,MDRD, CONSULTANT RADIOLOGIST