Name	MR.LOKESHA	ID	MED112152377
Age & Gender	30Y/MALE	Visit Date	20/04/2024
Ref Doctor Name	MediWheel		



ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in size and shows slightly increased echotexture.

No evidence of focal lesion or intrahepatic biliary ductal dilatation.

Hepatic and portal vein radicals are normal.

GALL BLADDER show normal shape and has clear contents.

Gall bladder wall is of normal thickness. CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern.

No evidence of ductal dilatation or calcification.

SPLEEN show normal shape, size and echopattern.

No demonstrable Para-aortic lymphadenopathy.

KIDNEYS move well with respiration and have normal shape, size and echopattern.

Cortico- medullary differentiations are well madeout.

No evidence of calculus or hydronephrosis.

	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney	11.2	1.4
Left Kidney	11.9	1.3

URINARY BLADDER is partially distended.

PROSTATE shows normal shape, size and echopattern.

No evidence of ascites.

IMPRESSION:

> GRADE I FATTY CHANGES IN LIVER.

CONSULTANT RADIOLOGISTS

DR. ANITHA ADARSH

DR. MOHAN B

> : MED112152377 Register On : 20/04/2024 9:53 AM

: 712412066 SID No.

Age / Sex : 30 Year(s) / Male Report On : 20/04/2024 3:14 PM

Type : OP

Investigation

PID No.

Ref. Dr : MediWheel Collection On : 20/04/2024 10:59 AM

Printed On : 21/04/2024 1:49 PM

> <u>Observed</u> <u>Unit</u> **Biological** Reference Interval <u>Value</u>

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING

 $({\rm EDTA~Blood} Agglutination)$

Remark: Test to be confirmed by Gel method.

'AB' 'Negative'







PID No. : MED112152377

SID No. : 712412066

Age / Sex : 30 Year(s) / Male

: MediWheel

Type : OP

Ref. Dr

: 0P

Register On : 20/04/2024 9:53 AM

Collection On : 20/04/2024 10:59 AM

Report On : 20/04/2024 3:14 PM

Printed On : 21/04/2024 1:49 PM



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	<u>Value</u>		Reference Interval

HAEMATOLOGY

Complete Blood Count With - ESR

Haemoglobin	14.7	g/dL	13.5 - 18.0
-------------	------	------	-------------

(EDTA Blood/Spectrophotometry)

INTERPRETATION: Haemoglobin values vary in Men, Women & Children. Low haemoglobin values may be due to nutritional deficiency, blood loss, renal failure etc. Higher values are often due to dehydration, smoking, high altitudes, hypoxia etc.

cross remaining etc. ringing variety are crear due	to dell' didition, sin	,, p.	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
PCV (Packed Cell Volume) / Haematocrit (EDTA Blood/Derived)	44.3	%	42 - 52
RBC Count (EDTA Blood/Automated Blood cell Counter)	5.65	mill/cu.mm	4.7 - 6.0
MCV (Mean Corpuscular Volume) (EDTA Blood/Derived from Impedance)	78.0	fL	78 - 100
MCH (Mean Corpuscular Haemoglobin) (EDTA Blood/Derived)	26.0	pg	27 - 32
MCHC (Mean Corpuscular Haemoglobin concentration) (EDTA Blood/Derived)	32.3	g/dL	32 - 36
RDW-CV (Derived)	15.1	%	11.5 - 16.0
RDW-SD (Derived)	41.22	fL	39 - 46
Total WBC Count (TC) (EDTA Blood/Derived from Impedance)	11310	cells/cu.mm	4000 - 11000
Neutrophils (Blood/Impedance Variation & Flow Cytometry)	56	%	40 - 75
Lymphocytes (Blood/Impedance Variation & Flow Cytometry)	37	%	20 - 45







PID No. : MED112152377

: 712412066

: MediWheel

SID No. Age / Sex : 30 Year(s) / Male

Type : OP

Ref. Dr

Register On : 20/04/2024 9:53 AM

Collection On : 20/04/2024 10:59 AM

Report On : 20/04/2024 3:14 PM

: 21/04/2024 1:49 PM **Printed On**



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Eosinophils (Blood/Impedance Variation & Flow Cytometry)	03	%	01 - 06
Monocytes (Blood/Impedance Variation & Flow Cytometry)	04	%	01 - 10
Basophils (Blood/Impedance Variation & Flow Cytometry)	00	%	00 - 02
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	6.33	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	4.18	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.34	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.45	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.00	10^3 / μl	< 0.2
Platelet Count (EDTA Blood/Derived from Impedance)	233	10^3 / μl	150 - 450
MPV (Blood/Derived)	12.4	fL	7.9 - 13.7
PCT	0.29	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate)	07	mm/hr	< 15



(Citrated Blood/Automated ESR analyser)





PID No. : MED112152377

SID No. : 712412066

Age / Sex : 30 Year(s) / Male

Type : OP

Ref. Dr : MediWheel

Register On : 20/04/2024 9:53 AM

Collection On : 20/04/2024 10:59 AM

Report On : 20/04/2024 3:14 PM

Printed On : 21/04/2024 1:49 PM



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulfanilic Acid)	0.7	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.3	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.40	mg/dL	0.1 - 1.0
Total Protein (Serum/Biuret)	8.1	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	5.9	gm/dl	3.5 - 5.2
Globulin (Serum/ <i>Derived</i>)	2.20	gm/dL	2.3 - 3.6
A : G Ratio (Serum/ <i>Derived</i>)	2.68		1.1 - 2.2
INTERPRETATION: Remark : Electrophoresis is the	preferred method		
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC / Kinetic)	44	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC / Kinetic)	48	U/L	5 - 41
Alkaline Phosphatase (SAP) (Serum/PNPP / Kinetic)	137	U/L	53 - 128
GGT(Gamma Glutamyl Transpeptidase)	64	U/L	< 55



(Serum/IFCC / Kinetic)





 PID No.
 : MED112152377
 Register On
 : 20/04/2024 9:53 AM

 SID No.
 : 712412066
 Collection On
 : 20/04/2024 10:59 AM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/Oxidase / Peroxidase method)	288	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Remark: Kindly correlate clinically			
Triglycerides (Serum/Glycerol phosphate oxidase / peroxidase)	417	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

: 21/04/2024 1:49 PM

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the õusualö circulating level of triglycerides during most part of the day.

Remark: Kindly correlate clinically

HDL Cholesterol (Serum/Immunoinhibition)	33	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	171.6	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	83.4	mg/dL	< 30







PID No. : MED112152377

: 712412066

Age / Sex : 30 Year(s) / Male

Type : OP

SID No.

Ref. Dr : MediWheel Register On : 20/04/2024 9:53 AM

Collection On : 20/04/2024 10:59 AM

Report On 20/04/2024 3:14 PM

: 21/04/2024 1:49 PM **Printed On**

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Non HDL Cholesterol (Serum/Calculated)	255.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio (Serum/Calculated)	8.7	Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
Triglyceride/HDL Cholesterol Ratio (TG/HDL) (Serum/Calculated)	12.6	Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
LDL/HDL Cholesterol Ratio (Serum/Calculated)	5.2	Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0







Very High: >= 220

: MED112152377 Register On : 20/04/2024 9:53 AM

Printed On

Type : OP

PID No.

Ref. Dr : MediWheel



Investigation Glycosylated Haemoglobin (HbA1c)	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HbA1C (Whole Blood/HPLC)	6.3	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

: 21/04/2024 1:49 PM

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 134.11 mg/dl

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.







: MED112152377

SID No. : 712412066

Age / Sex : 30 Year(s) / Male

Type : OP

PID No.

Ref. Dr : MediWheel

Register On : 20/04/2024 9:53 AM

Collection On : 20/04/2024 10:59 AM

Report On : 20/04/2024 3:14 PM

Printed On : 21/04/2024 1:49 PM



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	7.1		
Glucose Fasting (FBS) (Plasma - F/GOD- POD)	78	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Urine sugar, Fasting	Nil		Nil
(Urine - F)			
Glucose Postprandial (PPBS)	131	mg/dL	70 - 140
(Plasma - PP/GOD - POD)			

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Sugar (PP-2 hours) (Urine - PP)	Nil		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	8.6	mg/dL	7.0 - 21
Creatinine (Serum/Jaffe Kinetic)	1.2	mg/dL	0.9 - 1.3

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine etc.

Uric Acid **8.6** mg/dL 3.5 - 7.2

(Serum/Uricase/Peroxidase)







PID No.

Printed On

SID No. : 712412066 Collection On : 20/04/2024 10:59 AM

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u> <u>U</u>	<u>nit</u> <u>Biological</u>
	Value	Reference Interval

: 21/04/2024 1:49 PM

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.35 ng/ml 0.7 - 2.04

(Serum/Chemiluminescent Immunometric Assay (CLIA))

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Thyroxine) - Total 7.93 Microg/dl 4.2 - 12.0

(Serum/Chemiluminescent Immunometric Assay

(CLIA))

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 3.614 µIU/mL 0.35 - 5.50

(Serum/Chemiluminescent Immunometric Assay (CLIA))

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.







PID No. : MED112152377

SID No. : 712412066

. / 12412000

Age / Sex : 30 Year(s) / Male

Type : OP

Register On : 20/04/2024 9:53 AM

Collection On : 20/04/2024 10:59 AM

Report On : 20/04/2024 3:14 PM

Printed On : 21/04/2024 1:49 PM

Ref. Dr : MediWheel

medall

Investigation	<u>Observed</u> <u>Unit</u>	<u>Biological</u>
	<u>Value</u>	Reference Interval

CLINICAL PATHOLOGY

PHYSICAL EXAMINATION

Colour Pale Yellow Yellow to Amber

(Urine/Physical examination)

Volume 30 ml

(Urine/Physical examination)

Appearance Slightly Turbid

(Urine)

CHEMICAL EXAMINATION

pH 6.0 4.5 - 8.0

(Urine)

Specific Gravity 1.005 1.002 - 1.035

(Urine/Dip Stick oʻ'Reagent strip method)

Protein Negative Negative

(Urine/Dip Stick oʻ'Reagent strip method)

Glucose Nil Nil

(Urine)

Ketone Nil Nil

(Urine/Dip Stick ó"Reagent strip method)

Leukocytes Negative leuco/uL Negative

(Urine)

Nitrite Nil Nil

(Urine/Dip Stick ó"Reagent strip method)

Bilirubin Negative mg/dL Negative

(Urine)

(Urine)

Blood Nil Nil









PID No. : MED112152377

SID No. : 712412066

Age / Sex : 30 Year(s) / Male

Type : OP

Ref. Dr : MediWheel

Register On : 20/04/2024 9:53 AM

Collection On : 20/04/2024 10:59 AM

Report On : 20/04/2024 3:14 PM

Printed On : 21/04/2024 1:49 PM



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Urobilinogen (Urine/Dip Stick ó"Reagent strip method)	Normal		Within normal limits
<u>Urine Microscopy Pictures</u>			
RBCs (Urine/Microscopy)	Nil	/hpf	NIL
Pus Cells (Urine/Microscopy)	2-3	/hpf	< 5
Epithelial Cells (Urine/Microscopy)	1-2	/hpf	No ranges
Others (Urine)	Nil		Nil







APPROVED BY

-- End of Report --

Name	MR.LOKESHA	ID	MED112152377
Age & Gender	30Y/MALE	Visit Date	20/04/2024
Ref Doctor Name	MediWheel		



2 D ECHOCARDIOGRAPHIC STUDY

M mode measurement:

AORTA : 2.6cms

LEFT ATRIUM : 2.6cms

LEFT VENTRICLE (DIASTOLE) : 4.5cms

(SYSTOLE) : 2.5cms

VENTRICULAR SEPTUM (DIASTOLE) : 0.8cms

(SYSTOLE) : 1.0cms

POSTERIOR WALL (DIASTOLE) : 0.8cms

(SYSTOLE) : 1.0cms

EDV: 71ml

ESV : 27ml

FRACTIONAL SHORTENING : 33%

EJECTION FRACTION : 62%

RVID : 1.5cms

DOPPLER MEASUREMENTS:

MITRAL VALVE : E' - 0.77m/s A' - 0.29m/s NO MR

AORTIC VALVE : 0.98m/s NO AR

TRICUSPID VALVE : E' - 0.65 m/s A' - 0.25 m/s NO TR

PULMONARY VALVE : 0.70m/s NO PR

2D ECHOCARDIOGRAPHY FINDINGS:

Name	MR.LOKESHA	ID	MED112152377
Age & Gender	30Y/MALE	Visit Date	20/04/2024
Ref Doctor Name	MediWheel		



Left ventricle : Normal size, Normal systolic function.

No regional wall motion abnormalities.

Left Atrium : Normal.

Right Ventricle : Normal.

Right Atrium : Normal.

Mitral valve : Normal, No mitral valve prolapse.

Aortic valve : Normal, Trileaflet.

Tricuspid valve : Normal.

Pulmonary valve : Normal.

IAS : Intact.

IVS : Intact.

Pericardium : No pericardial effusion.

IMPRESSION:

- > NORMAL SIZED CARDIAC CHAMBERS.
- > NORMAL LV SYSTOLIC FUNCTION. EF: 62%.
- > NO REGIONAL WALL MOTION ABNORMALITIES.
- > NORMAL VALVES.
- > NO CLOTS/ PERICARDIAL EFFUSION VEGETATION.



DR. NIKHIL B
INTERVENTIONAL CARDIOLOGIST
NB/sv



Name	Mr. LOKESHA	ID	MED112152377
Age & Gender	30Y/M	Visit Date	Apr 20 2024 9:50AM
Ref Doctor	MediWheel		

X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

Impression: No significant abnormality detected.

Dr. Anitha Adarsh Consultant Radiologist



Name	Mr. LOKESHA	ID	MED112152377
Age & Gender	30Y/M	Visit Date	Apr 20 2024 9:50AM
Ref Doctor	MediWheel		

X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

Impression: No significant abnormality detected.

Dr. Anitha Adarsh Consultant Radiologist



Name	Mr. LOKESHA	ID	MED112152377
Age & Gender	30Y/M	Visit Date	Apr 20 2024 9:50AM
Ref Doctor	MediWheel		

X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

Impression: No significant abnormality detected.

Dr. Anitha Adarsh Consultant Radiologist