



Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 QN: U85110UP2003PLC193493

Patient Name	: Mr.ANKUSH	Registered On	: 24/Nov/2024 09:56:39
Age/Gender	: 32 Y 0 M 0 D / M	Collected	: 24/Nov/2024 10:07:08
UHID/MR NO	: CALI.0000061192	Received	: 24/Nov/2024 11:13:42
Visit ID	: CALI0181522425	Reported	: 24/Nov/2024 13:41:39
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

DEPARTMENT OF HAEM ATOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
Blood Group (ABO & Rh typing)** , Blood			
Blood Group	0			ERYTHROCYTE MAGNETIZED TECHNOLOGY/ TUBE AGGLUTINA
Rh (Anti-D)	POSITIVE			ERYTHROCYTE MAGNETIZED TECHNOLOGY/ TUBE AGGLUTINA
Complete Blood Count (CBC) **	, EDTA Whole Blood			
Haemoglobin	12.80	g/ dl	1 Day- 14.5-22.5 g/dl 1 Wk- 13.5-19.5 g/dl 1 Mo- 10.0-18.0 g/dl 3-6 Mo- 9.5-13.5 g/dl 0.5-2 Yr- 10.5-13.5 g/dl 2-6 Yr- 11.5-15.5 g/dl 6-12 Yr- 11.5-15.5 g/dl 12-18 Yr 13.0-16.0 g/dl Male- 13.5-17.5 g/dl Female- 12.0-15.5 g/dl	COLORIMETRIC METHOD (CYANIDE-FREE REAGENT)
TLC (WBC) DLC	8,600.00	/Qumm	4000-10000	IMPEDANCE METHOD
Polymorphs (Neutrophils)	50.00	%	40-80	FLOW CYTOMETRY
Lymphocytes	38.00	%	20-40	FLOW CYTOMETRY
Monocytes	6.00	%	2-10	FLOW CYTOMETRY
Eosinophils	6.00	%	1-6	FLOW CYTOMETRY
Basophils ESR	0.00	%	<1-2	FLOW CYTOMETRY
Observed	16.00	MM/1H	10-19 Yr 8.0 20-29 Yr 10.8 30-39 Yr 10.4 40-49 Yr 13.6 50-59 Yr 14.2 60-69 Yr 16.0 70-79 Yr 16.5 80-91 Yr 15.8	



Page 1 of 13







Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 GN: U85110UP2003PLC193493

Patient Name	: Mr.ANKUSH	Registered On	: 24/Nov/2024 09:56:39
Age/Gender	: 32 Y 0 M 0 D / M	Collected	: 24/Nov/2024 10:07:08
UHID/MR NO	: CALI.0000061192	Received	: 24/Nov/2024 11:13:42
Visit ID	: CALI0181522425	Reported	: 24/Nov/2024 13:41:39
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

DEPARTMENT OF HAEM ATOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
			Pregnancy Early gestation - 48 (62 if anaemic) Leter gestation - 70 (95 if anaemic)	
Corrected	10.00	Mm for 1st hr.	. <9	
PCV (HCT)	38.00	%	40-54	
Platelet count				
Platelet Count	1.95	LACS' cu mm	1.5-4.0	ELECTRONIC IMPEDANCE/MICROSCOPIC
PDW (Platelet Distribution width)	16.00	fL	9-17	ELECTRONIC IMPEDANCE
P-LOR (Platelet Large Cell Patio)	47.00	%	35-60	ELECTRONIC IMPEDANCE
PCT (Platelet Hematocrit)	0.24	%	0.108-0.282	ELECTRONIC IMPEDANCE
MPV (Mean Platelet Volume)	12.20	fL	6.5-12.0	ELECTRONIC IMPEDANCE
RBCCount				
RBC Count	6.43	Mill./cumm	4.2-5.5	ELECTRONIC IMPEDANCE
Blood Indices (MCV, MCH, MCHC)				
MCV	63.30	fl	80-100	CALCULATED PARAMETER
МСН	19.80	pg	27-32	CALCULATED PARAMETER
MCHC	31.30	%	30-38	CALCULATED PARAMETER
RDW-CV	17.30	%	11-16	ELECTRONIC IMPEDANCE
RDW-SD	39.00	fL	35-60	ELECTRONIC IMPEDANCE
Absolute Neutrophils Count	4,300.00	/cumm	3000-7000	
Absolute Eosinophils Count (AEC)	516.00	/cu mm	40-440	

Linth

DR.KIRITI KANAUJIA MBBS MD(PATH)



Chandan 24x7 App





Home Sample Collection 08069366666





Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 QN: U85110UP2003PLC193493

Patient Name	: Mr.ANKUSH	Registered On	: 24/Nov/2024 09:56:40
Age/Gender	: 32 Y 0 M 0 D / M	Collected	: 24/Nov/2024 10:07:07
UHID/MR NO	: CALI.0000061192	Received	: 24/Nov/2024 11:43:21
Visit ID	: CALI0181522425	Reported	: 24/Nov/2024 13:52:36
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd. -	Status	: Final Report

DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Un	it Bio. Ref. Interv	al Method
GLUCOSE FASTING ** , Plasma Glucose Fasting	87.10	mg/dl	<100 Normal 100-125 Pre-diabetes ≥ 126 Diabetes	GOD POD

Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions.b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential.c) I.G.T = Impaired Glucose Tolerance.

CLINICAL SIGNIFICANCE:- Glucose is the major source of energy in the body. Lack of insulin or resistance to it section at the cellular level causes diabetes. Therefore, the blood glucose levels are very high. Elevated serum glucose levels are observed in diabetes mellitus and may be associated with pancreatitis, pituitary or thyroid dysfunction and liver disease. Hypoglycaemia occurs most frequently due to over dosage of insulin.

Gucose PP**	135.50	mg/ dl	<140 Normal	GOD POD
Sample:Plasma After Meal			140-199 Pre-diabetes	
			>200 Diabetes	

Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions.b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential.c) I.G.T = Impaired Glucose Tolerance.

GLYCOSYLATED HAEMOGLOBIN (HBA1C) ** , EDTA Whole Blood

Glycosylated Haemoglobin (HbA1c)	5.60	%NGSP	HPLC (NGSP)
Glycosylated Haemoglobin (HbA1c)	38.00	mmol/mol/IFCC	
Estimated Average Glucose (eAG)	114	mg/dl	

Interpretation:

<u>NOTE</u>:-

• eAG is directly related to A1c.



View Reports on

Chandan 24x7 App







Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 QN: U85110UP2003PLC193493

Patient Name	: Mr.ANKUSH	Registered On	: 24/Nov/2024 09:56:40
Age/Gender	: 32 Y 0 M 0 D / M	Collected	: 24/Nov/2024 10:07:07
UHID/MR NO	: CALI.0000061192	Received	: 24/Nov/2024 11:43:21
Visit ID	: CALI0181522425	Reported	: 24/Nov/2024 13:52:36
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd. -	Status	: Final Report

DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

est Name	Result	Unit	Bio. Ref. Interval	Method
----------	--------	------	--------------------	--------

- An A1c of 7% -the goal for most people with diabetes-is the equivalent of an eAG of 154 mg/dl.
- eAG may help facilitate a better understanding of actual daily control helping you and your health care provider to make necessary changes to your diet and physical activity to improve overall diabetes mnagement.

The following ranges may be used for interpretation of results. However, factors such as duration of diabetes, adherence to therapy and the age of the patient should also be considered in assessing the degree of blood glucose control.

Haemoglobin A1C (%)NGSP	mmol/mol / IFCC Unit	eAG (mg/dl)	Degree of Glucose Control Unit
> 8	>63.9	>183	Action Suggested*
7-8	53.0 -63.9	154-183	Fair Control
< 7	<63.9	<154	Goal**
6-7	42.1 -63.9	126-154	Near-normal glycemia
< 6%	<42.1	<126	Non-diabetic level

*High risk of developing long term complications such as Retinopathy, Nephropathy, Neuropathy, Cardiopathy, etc. **Some danger of hypoglycemic reaction in Type 1diabetics. Some glucose intolerant individuals and "subclinical" diabetics may demonstrate HbA1C levels in this area.

N.B.: Test carried out on Automated VARIANT II TURBO HPLC Analyser.

<u>Clinical Implications:</u>

*Values are frequently increased in persons with poorly controlled or newly diagnosed diabetes.

*With optimal control, the HbA 1c moves toward normal levels.

*A diabetic patient who recently comes under good control may still show higher concentrations of glycosylated hemoglobin. This level declines gradually over several months as nearly normal glycosylated *Increases in glycosylated hemoglobin occur in the following non-diabetic conditions: a. Iron-deficiency anemia b. Splenectomy

c. Alcohol toxicity d. Lead toxicity

*Decreases in A 1c occur in the following non-diabetic conditions: a. Hemolytic anemia b. chronic blood loss

*Pregnancy d. chronic renal failure. Interfering Factors:

*Presence of Hb F and H causes falsely elevated values. 2. Presence of Hb S, C, E, D, G, and Lepore (autosomal recessive mutation resulting in a hemoglobinopathy) causes falsely decreased values.

BUN (Blood	Urea	Nitroger	1)**
Sampl	e:Serum		-	

12.30

mg/dL 7.0-23.0

CALCULATED





Page 4 of 13





Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 GN: U85110UP2003PLC193493

Patient Name	: Mr.ANKUSH	Registered On	: 24/Nov/2024 09:56:40
Age/Gender	: 32 Y 0 M 0 D / M	Collected	: 24/Nov/2024 10:07:07
UHID/MR NO	: CALI.0000061192	Received	: 24/Nov/2024 11:43:21
Visit ID	: CALI0181522425	Reported	: 24/Nov/2024 13:52:36
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd. -	Status	: Final Report

DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	U	nit Bio. Ref. Int	terval Method
nterpretation:				
Note: Elevated BUN levels can be seen in the	he following:			
Iigh-protein diet, Dehydration, Aging, Certain n	nedications, Burns	s, Gastrointestin	nal (GI) bleeding.	
Low BUN levels can be seen in the following	g:			
ow-protein diet, overhydration, Liver disease.				
reatinine * * ample:Serum	1.18	mg/dl	Male 0.7-1.3 Newborn 0.3-1.0 Infent 0.2-0.4 Child 0.3-0.7	MODIFIED JAFFES
Interpretation:	a interpreted in 1:	abt of the peti-	Adolescent 0.5- 1.0	
Interpretation: The significance of single creatinine value must be mass will have a higher creatinine concentration. absolute creatinine concentration. Serum creating could be affected mildly and may result in anoma- lipemic.	The trend of seru ine concentrations	m creatinine co s may increase v	ts muscle mass. A pation ncentrations over time when an ACE inhibitor	ent with a greater muscle is more important than · (ACE) is taken. The assa
The significance of single creatinine value must be mass will have a higher creatinine concentration. absolute creatinine concentration. Serum creating could be affected mildly and may result in anoma- lipemic. ric Acid ** <i>ample:Serum</i> Interpretation: Note:-	The trend of seru ine concentrations alous values if seru 7.92	m creatinine co s may increase v um samples hav	ts muscle mass. A pation ncentrations over time when an ACE inhibitor e heterophilic antibodio	ent with a greater muscle is more important than (ACE) is taken. The assa es, hemolyzed, icteric or
The significance of single creatinine value must be mass will have a higher creatinine concentration. absolute creatinine concentration. Serum creatini could be affected mildly and may result in anoma lipemic. ric Acid ** ample:Serum Interpretation: Note:- Elevated uric acid levels can be seen in the f	The trend of seru ine concentrations alous values if seru 7.92 following:	m creatinine co s may increase v um samples hav mg/ dl	ts muscle mass. A patiencentrations over time when an ACE inhibitor e heterophilic antibodie 3.5-7.2	ent with a greater muscle is more important than (ACE) is taken. The assa es, hemolyzed, icteric or
The significance of single creatinine value must be mass will have a higher creatinine concentration. absolute creatinine concentration. Serum creating could be affected mildly and may result in anoma- lipemic.	The trend of seru ine concentrations alous values if seru 7.92 following:	m creatinine co s may increase v um samples hav mg/ dl	ts muscle mass. A patiencentrations over time when an ACE inhibitor e heterophilic antibodie 3.5-7.2	ent with a greater muscle is more important than (ACE) is taken. The assa es, hemolyzed, icteric or
The significance of single creatinine value must be mass will have a higher creatinine concentration. absolute creatinine concentration. Serum creating could be affected mildly and may result in anoma- lipemic. Fric Acid ** <i>ample:Serum</i> Interpretation: Note:- Elevated uric acid levels can be seen in the f Drugs, Diet (high-protein diet, alcohol), Chronic	The trend of seru ine concentrations alous values if seru 7.92 following:	m creatinine co s may increase v um samples hav mg/ dl	ts muscle mass. A patiencentrations over time when an ACE inhibitor e heterophilic antibodie 3.5-7.2	ent with a greater muscle is more important than (ACE) is taken. The assa es, hemolyzed, icteric or
The significance of single creatinine value must be mass will have a higher creatinine concentration. absolute creatinine concentration. Serum creating could be affected mildly and may result in anoma- lipemic. Fric Acid ** Elevated uric acid levels can be seen in the f Drugs, Diet (high-protein diet, alcohol), Chronic FT (WITH GAMMA GT) ** , <i>Serum</i> SGOT / Aspartate Aminotransferase (AST) SGPT / Alanine Aminotransferase (ALT)	The trend of seru ine concentrations alous values if seru 7.92 following: kidney disease, F 31.40 46.30	m creatinine co s may increase v um samples hav mg/dl Hypertension, C U/L U/L	ts muscle mass. A patiencentrations over time when an ACE inhibitor e heterophilic antibodie 3.5-7.2	ent with a greater muscle is more important than (ACE) is taken. The assa es, hemolyzed, icteric or URICASE
The significance of single creatinine value must be mass will have a higher creatinine concentration. absolute creatinine concentration. Serum creating could be affected mildly and may result in anoma- lipemic. Fric Acid ** Elevated uric acid levels can be seen in the f Drugs, Diet (high-protein diet, alcohol), Chronic FT (WITH GAMMA GT) ** , <i>Serum</i> SGOT / Aspartate Aminotransferase (AST)	The trend of seru ine concentrations alous values if seru 7.92 following: kidney disease, F 31.40	m creatinine co s may increase v um samples hav mg/dl Hypertension, C	ts muscle mass. A patiencentrations over time when an ACE inhibitor e heterophilic antibodie 3.5-7.2 besity.	ent with a greater muscle is more important than (ACE) is taken. The assa es, hemolyzed, icteric or URICASE









Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 GN: U85110UP2003PLC193493

Patient Name	: Mr.ANKUSH	Registered On	: 24/Nov/2024 09:56:40
Age/Gender	: 32 Y 0 M 0 D / M	Collected	: 24/Nov/2024 10:07:07
UHID/MR NO	: CALI.0000061192	Received	: 24/Nov/2024 11:43:21
Visit ID	: CALI0181522425	Reported	: 24/Nov/2024 13:52:36
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	U	nit Bio. Ref. Interv	al Method
Albumin	4.06	gm/ dl	3.4-5.4	B.C.G.
Globulin	2.51	gm/ dl	1.8-3.6	CALCULATED
A:G Patio	1.62		1.1-2.0	CALCULATED
Alkaline Phosphatase (Total)	101.92	U/L	53-128	IFCC AMP KINETIC
Bilirubin (Total)	1.08	mg/ dl	Adult 0-2.0	DIAZO
Bilirubin (Direct)	0.31	mg/ dl	< 0.20	DIAZO
Bilirubin (Indirect)	0.77	mg/ dl	<1.8	CALCULATED
LIPID PROFILE (MINI) ** , Serum				
Cholesterol (Total)	176.00	mg/ dl	<200 Desirable 200-239 Borderline Higl >240 High	OHOD-PAP 1
HDL Cholesterol (Good Cholesterol)	39.50	mg/ dl	35.0-79.5	DIRECT ENZYMATIC
LDL Cholesterol (Bad Cholesterol)	55	mg/dl	< 100 Optimal 100-129 Nr. Optimal/ Above Optima 130-159 Borderline High 160-189 High > 190 Very High	
VLDL	81.38	mg/dl	10-33	CALCULATED
Triglycerides	406.90	mg/ dl	<150 Normal 150-199 Borderline Higl 200-499 High >500 Very High	GPO-PAP

Dr. Anupam Singh (MBBS MD Pathology)











Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 QN: U85110UP2003PLC193493

Patient Name	: Mr.ANKUSH	Registered On	: 24/Nov/2024 09:56:39
Age/Gender	: 32 Y 0 M 0 D / M	Collected	: 24/Nov/2024 11:58:10
UHID/MR NO	: CALI.0000061192	Received	: 24/Nov/2024 14:13:29
Visit ID	: CALI0181522425	Reported	: 24/Nov/2024 17:43:45
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

DEPARTMENT OF CLINICAL PATHOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
URINE EXAMINATION, ROUTIN	IE** , Urine			
Color	PALEYELLOW			
Specific Gravity	1.010			
Reaction PH	Acidic (6.0)			DIPSTICK
Appearance	CLEAR			
Protein	ABSENT	mg %	<10 Absent 10-40 (+) 40-200 (++) 200-500 (+++) >500 (++++)	DIPSTICK
Sugar	ABSENT	gms%	<0.5 (+) 0.5-1.0 (++) 1-2 (+++) >2 (++++)	DIPSTICK
Ketone	ABSENT	mg/dl	Serum-0.1-3.0 Urine-0.0-14.0	BICCHEMISTRY
Bile Salts	ABSENT			
Bile Pigments	ABSENT			
Bilirubin	ABSENT			DIPSTICK
Leucocyte Esterase	ABSENT			DIPSTICK
Urobilinogen(1:20 dilution)	ABSENT			
Nitrite	ABSENT			DIPSTICK
Blood	ABSENT			DIPSTICK
Microscopic Examination:				
Epithelial œlls	1-3/h.p.f			MICROSCOPIC EXAMINATION
Puscells	0-2/h.p.f			
RBC:	ABSENT			MICROSCOPIC EXAMINATION
Cast	ABSENT			
Crystals	ABSENT			MICROSCOPIC EXAMINATION
Others	ABSENT			
SUGAR, FASTING STAGE** , Ur	ine			
Sugar, Fasting stage	ABSENT	gms%		



Chandan 24x7 App





33)

CHANDAN DIAGNOSTIC CENTRE

Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 CIN: U85110UP2003PLC193493

Patient Name	: Mr.ANKUSH	Registered On	: 24/Nov/2024 09:56:39
Age/Gender	: 32 Y 0 M 0 D / M	Collected	: 24/Nov/2024 11:58:10
UHID/MR NO	: CALI.0000061192	Received	: 24/Nov/2024 14:13:29
Visit ID	: CALI0181522425	Reported	: 24/Nov/2024 17:43:45
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

DEPARTMENT OF CLINICAL PATHOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method

Interpretation:

(+) < 0.5 (++) 0.5-1.0 (+++) 1-2 (++++) > 2

SUGAR, PP STAGE** , Urine

Sugar, PP Stage

ABSENT

Interpretation:

(+) < 0.5 gms% (++) 0.5-1.0 gms% (+++) 1-2 gms% (++++) > 2 gms%



Page 8 of 13

View Reports on

Chandan 24x7 App

Dr. Mamta Barthwal MD(Micro-Biology)





 \odot





Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 QN: U85110UP2003PLC193493

Patient Name	: Mr.ANKUSH	Registered On	: 24/Nov/2024 09:56:40
Age/Gender	: 32 Y 0 M 0 D / M	Collected	: 24/Nov/2024 10:07:07
UHID/MR NO	: CALI.0000061192	Received	: 24/Nov/2024 10:57:36
Visit ID	: CALI0181522425	Reported	: 24/Nov/2024 13:07:46
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd. -	Status	: Final Report

DEPARTMENT OF IMMUNOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
PSA (Prostate Specific Antigen), Total ** Sample:Serum	0.16	ng/mL	<4.1	CLIA

Interpretation:

- 1. PSA is detected in the serum of males with normal, benign hypertrophic, and malignant prostate tissue.
- 2. Measurement of serum PSA levels is not recommended as a screening procedure for the diagnosis of cancer because elevated PSA levels also are observed in patients with benign prostatic hypertrophy. However, studies suggest that the measurement of PSA in conjunction with digital rectal examination (DRE) and ultrasound provide a better method of detecting prostate cancer than DRE alone[.]
- 3. PSA levels increase in men with cancer of the prostate, and after radical prostatectomy PSA levels routinely fall to the undetectable range.
- 4. If prostatic tissue remains after surgery or metastasis has occurred, PSA appears to be useful in detecting residual and early recurrence of tumor.
- 5. Therefore, serial PSA levels can help determine the success of prostatectomy, and the need for further treatment, such as radiation, endocrine or chemotherapy, and in the monitoring of the effectiveness of therapy.

THYROID PROFILE - TOTAL**, Serum

T3, Total (tri-iodothyronine)	86.15	ng/ dl	84.61–201.7	CLIA
T4, Total (Thyroxine)	4.70	ug/dl	3.2-12.6	CLIA
TSH (Thyroid Stimulating Hormone)	5.680	µIU/mL	0.27 - 5.5	CLIA

Interpretation:

0.3-4.5	µIU/mL	First Trimest	er	
0.5-4.6	µIU/mL	Second Trimester		
0.8-5.2	µIU/mL	Third Trimes	ter	
0.5-8.9	µIU/mL	Adults	55-87 Years	
0.7-27	µIU/mL	Premature	28-36 Week	
2.3-13.2	µIU/mL	Cord Blood	> 37Week	
0.7-64	µIU/mL	Child(21 wk	- 20 Yrs.)	
1-39	µIU/mL	Child	0-4 Days	
1.7-9.1	µIU/mL	Child	2-20 Week	

08069366666

1) Patients having low T3 and T4 levels but high TSH levels suffer from primary hypothyroidism, cretinism, juvenile myxedema or









Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 QN: U85110UP2003PLC193493

Patient Name	: Mr.ANKUSH	Registered On	: 24/Nov/2024 09:56:40
Age/Gender	: 32 Y 0 M 0 D / M	Collected	: 24/Nov/2024 10:07:07
UHID/MR NO	: CALI.0000061192	Received	: 24/Nov/2024 10:57:36
Visit ID	: CALI0181522425		: 24/Nov/2024 13:07:46
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

DEPARTMENT OF IMMUNOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method

autoimmune disorders.

2) Patients having high T3 and T4 levels but low TSH levels suffer from Grave's disease, toxic adenoma or sub-acute thyroiditis.

3) Patients having either low or normal T3 and T4 levels but low TSH values suffer from iodine deficiency or secondary hypothyroidism.

4) Patients having high T3 and T4 levels but normal TSH levels may suffer from toxic multinodular goiter. This condition is mostly a symptomatic and may cause transient hyperthyroidism but no persistent symptoms.

5) Patients with high or normal T3 and T4 levels and low or normal TSH levels suffer either from T3 toxicosis or T4 toxicosis respectively.

6) In patients with non thyroidal illness abnormal test results are not necessarily indicative of thyroidism but may be due to adaptation to the catabolic state and may revert to normal when the patient recovers.

7) There are many drugs for eg. Glucocorticoids, Dopamine, Lithium, Iodides, Oral radiographic dyes, etc. which may affect the thyroid function tests.

8) Generally when total T3 and total T4 results are indecisive then Free T3 and Free T4 tests are recommended for further confirmation along with TSH levels.

Dr. Anupam Singh (MBBS MD Pathology)











Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 QN: U85110UP2003PLC193493

Patient Name	: Mr.ANKUSH	Registered On	: 24/Nov/2024 09:56:40
Age/Gender	: 32 Y 0 M 0 D / M	Collected	: 2024-11-24 13:10:48
UHID/MR NO	: CALI.0000061192	Received	: 2024-11-24 13:10:48
Visit ID	: CALI0181522425	Reported	: 24/Nov/2024 13:11:45
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

DEPARTMENT OF X-RAY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

X-RAY DIGITAL CHEST PA** (500 mA COMPUTERISED UNIT SPOT FILM DEVICE)

DIGITAL CHEST P-A VIEW

- Small left cervical rib seen.
- Costo-phrenic angles are bilaterally clear.
- Trachea is central in position.
- Cardiac size & contours are normal.
- Hilar shadows are normal.
- Pulmonary vascularity & distribution are normal.
- Pulmonary parenchyma did not reveal any significant lesion.

IMPRESSION:

• NO SIGNIFICANT DIAGNOSTIC ABNORMALITY SEEN.

Dr. Pankaj Kumar Gupta (M.B.B.S D.M.R.D)



Page 11 of 13







Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 QN: U85110UP2003PLC193493

Patient Name	: Mr.ANKUSH	Registered On	: 24/Nov/2024 09:56:40
Age/Gender	: 32 Y 0 M 0 D / M	Collected	: 2024-11-24 10:42:23
UHID/MR NO	: CALI.0000061192	Received	: 2024-11-24 10:42:23
Visit ID	: CALI0181522425	Reported	: 24/Nov/2024 10:44:41
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd	[.] Status	: Final Report

DEPARTMENT OF ULTRASOUND

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

ULTRASOUND WHOLE ABDOM EN (UPPER & LOWER) **

WHOLE ABDOMEN ULTRASONOGRAPHY REPORT

LIVER

 Liver is normal in size ~ 13.1 cm in longitudinal span and shows diffused raised echogenicity of hepatic parenchyma S/O grade I fatty liver. No focal lesion is seen.

PORTAL SYSTEM

- The intra hepatic portal channels are normal.
- The portal vein is not dilated.
- Porta hepatis is normal.

BILIARY SYSTEM

- The intra-hepatic biliary radicles are normal.
- Common duct is not dilated.
- The gall bladder is normal in size and has regular walls. Lumen of the gall bladder is anechoic.

PANCREAS

• The pancreas is normal in size and shape and has a normal homogenous echotexture. Pancreatic duct is not dilated.

KIDNEYS

- Right kidney is normal in size ~ 9.5 x 4.3 cm position and cortical echotexture. Corticomedullary demarcation is maintained.
- Left kidney is normal in size ~ 9.7 x 4.6 cm position and cortical echotexture. Corticomedullary demarcation is maintained.
- The collecting system of both the kidneys are not dilated.

SPLEEN

• The spleen is normal in size ~ 9.8 cm and has a normal homogenous echo-texture.

ILIAC FOSSAE & PERITONEUM

- Scan over the iliac fossae does not reveal any fluid collection or mass.
- No free fluid is noted in peritoneal cavity.
- Visualized bowel loops are gaseous and grossly appear normal in caliber, peristalsis and wall thickness.



Page 12 of 13







Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 QN: U85110UP2003PLC193493

Patient Name	: Mr.ANKUSH	Registered On	: 24/Nov/2024 09:56:40
Age/Gender	: 32 Y 0 M 0 D / M	Collected	: 2024-11-24 10:42:23
UHID/MR NO	: CALI.0000061192	Received	: 2024-11-24 10:42:23
Visit ID	: CALI0181522425	Reported	: 24/Nov/2024 10:44:41
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

DEPARTMENT OF ULTRASOUND

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

URINARY BLADDER

• The urinary bladder is normal. Bladder wall is normal in thickness and is regular. No calculus is seen.

PROSTATE

The prostate gland is normal in size with smooth outline (volume ~ 18.8 cc).

FINAL IMPRESSION

- GRADE I FATTY INFILTRATION OF LIVER.
- NO OTHER SIGNIFICANT ABNORMALITY SEEN.

Adv: Clinico-pathological correlation and follow-up.

*** End Of Report ***

(**) Test Performed at Chandan Speciality Lab.

Result/s to Follow:

STOOL, ROUTINE EXAMINATION, ECG / EKG, Tread Mill Test (TMT)





This report is not for medico legal purpose. If clinical correlation is not established, kindly repeat the test at no additional cost within seven days

Facilities: MRI, CT scan, DR X-ray, Ultrasound, Sonomammography, Digital Mammography, ECG (Bedside also), 2D Echo, TMT, Holter, OPG, EEG, NCV, EMG & BERA, Audiometry, BMD, PFT, Fibroscan, Bronchoscopy, Colonoscopy and Endoscopy, Allergy Testing, Biochemistry & Immunoassay, Hematology, Microbiology & Serology, Histopathology & Immunohistochemistry, Cytogenetics and Molecular Diagnostics and Health Checkups 365 Days Open

*Facilities Available at Select Location





View Reports on Chandan 24x7 App

