

Dr. Goyal's

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sangar Road, Jaipur-302019

Tele : 0141-2293346, 4049787, 9887048787

Website : www.dr.goyalspathlab.com | E-mail :

General Physical Examination

Date of Examination: 23-11-2023

Name: ANJANA JAIN Age: 39 Sex: Female

DOB: 01-08-1984

Referred By: MedP WOHLL

Photo ID: Adhnaa ID #: attached

Ht: 159 (cm)

Wt: 57 (Kg)

Chest (Expiration): 83 (cm)

Abdomen Circumference: 73 (cm)

Blood Pressure: 129/80 mm Hg PR: 80 / min

BMI 18.5 kg/m²

Eye Examination: D/A vision G/B with spec. ²⁵/₄ Near.
vision N/G. No Glauk blindness.

Other: Not significant.

On examination he/she appears physically and mentally fit: Yes/No

Signature Of Examinee : _____

Name of Examinee: Anjana Jain

Signature Medical Examiner _____

Name Medical Examiner _____

Dr. Piyush Goyal
M.B.B.S., D.M.R.D.
RMC Reg. No.-017996

CENTRAL MOTOR VEHICLES

RULES 1989

FORM 7(See Rule 16(2))

DRIVING LICENCE



D/L NO : RJ-14/DLC/12/ 991186 Date : 08/11/2012

Name : ANJANA JAIN

Daughter of : AAKASH JAIN

Address : 482 BARKAT NAGAR

TONK PHATAK JAIPUR

is licenced to drive throughout India a vehicle of the following description.

MCY WITH GEAR,LIGHT MOTOR VEH,

The licence to drive other than transport vehicle is valid

From: 08/11/2012 To: 07/11/2032

Holder's Sig./thumb impression

Licencing Authority,JAIPUR

Dr. Piyush Goyal
M.B.B.S., D.M.R.D.
RMC Reg. No.-017996

Health checkup at tie-up Ctr

HealthChkUp Authorisatn letter



Union Bank of India

RO - JAIPUR
101 A 101 B & 102 KISAN BHAWAN, LAL
KOTHI, TONK ROAD, Jaipur- 141

To,

The Chief Medical Officer

M/S Mediwheel
<https://med/wheel.in/signup011-41195959>(A brand name of
Arofemi Healthcare Ltd),
Mumbai400021

Dear Sir,

Tie-up arrangement for Health Checkup under Health Checkup 35-40 Female

Shri/Smt./Kum. ANJANA JAIN,

P.F. No. 652697

Designation : Senior Manager

Checkup for Financial Year

2023-
2024

Approved Charges Rs.

3000.00

The above mentioned staff member of our Branch/Office desires to undergo Health Checkup at your Hospital/Centre/Clinic, under the tie-up arrangement entered into with you, by our bank.

Please send the receipt of the above payment and the relevant reports to our above address.

Thanking you,

(Signature of the Employee)

Yours Faithfully,

BRANCH MANAGER/SENIOR MANAGER

PS : Status of the application- **Sanctioned**




DR. GOYAL PATH LAB

2765 / MRS ANJANA JAIN / 39 Yrs / F / Non Smoker

Heart Rate : 67 bpm / Tested On : 23-Nov-23 09:51:51 / HF 0.05 Hz - LF 35 Hz / Notch 50 Hz / Sn 1.00 Cm/mV / Sw 25 mm/s / Refd By: MEDIWHEEL

ECG



Vent Rate : 67 bpm

PR Interval : 120 ms

QRS Duration : 74 ms

QT/QTc Int : 394/406 ms

P-QRS-T axis : 42.00° • 70.00° • 19.00°



Sinus bradycardia with + invagination in lead I

Reported By:

Dr. Naresh Kumar Mohanka
RMC No. 25703
SAS, DIP, CARDIO (ESCORTS)
D.E.M. (RCGP-UK)

Dr. Goyal's

Path Lab & Imaging Centre



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganeer Road, Jaipur-302018
Tale : 0141-2293346, 4049787, 9887048787
Website : www.drgoyalpathlab.com | E-mail : drgoyalpiyush@gmail.com

Date :- 23/11/2023 09:29:31
NAME :- Mrs. ANJANA JAIN
Sex / Age :- Female 39 Yrs 3 Mon 23 Days
Company :- MediWheel

Patient ID :- 12234347
Ref. By Dr:-
Lab/Hosp :-



Sample Type :- EDTA

Sample Collected Time 23/11/2023 10:20:29

Final Authentication : 23/11/2023 12:28:02

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
HAEMOGARAM			
HAEMOGLOBIN (Hb)	13.3	g/dL	12.0 - 15.0
TOTAL LEUCOCYTE COUNT	6.99	/cumm	4.00 - 10.00
DIFFERENTIAL LEUCOCYTE COUNT			
NEUTROPHIL	74.7	%	40.0 - 80.0
LYMPHOCYTE	21.5	%	20.0 - 40.0
EOSINOPHIL	1.0	%	1.0 - 6.0
MONOCYTE	2.4	%	2.0 - 10.0
BASOPHIL	0.4	%	0.0 - 2.0
NEUT#	5.23	$10^3/uL$	1.50 - 7.00
LYMPH#	1.51	$10^3/uL$	1.00 - 3.70
EO#	0.06	$10^3/uL$	0.00 - 0.40
MONO#	0.16	$10^3/uL$	0.00 - 0.70
BASO#	0.03	$10^3/uL$	0.00 - 0.10
TOTAL RED BLOOD CELL COUNT (RBC)	4.79	$\times 10^6/uL$	3.80 - 4.80
HEMATOCRIT (HCT)	41.60	%	36.00 - 46.00
MEAN CORP VOLUME (MCV)	87.0	fL	83.0 - 101.0
MEAN CORP HB (MCH)	27.7	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	31.9	g/dL	31.5 - 34.5
PLATELET COUNT	262	$\times 10^3/uL$	150 - 410
RDW-CV	13.7	%	11.6 - 14.0
MENTZER INDEX	18.16		

The Mentzer index is used to differentiate iron deficiency anemia from beta thalassemia trait. If a CBC indicates microcytic anemia, these are two of the most likely causes, making it necessary to distinguish between them.

If the quotient of the mean corpuscular volume divided by the red blood cell count is less than 13, thalassemia is more likely. If the result is greater than 13, then iron-deficiency anemia is more likely.

MUKESH SINGH
Technologist

Page No: 2 of 12



Dr. Chandrika Gupta
MBBS.MD (Path)
RMC NO. 21021/008037

Dr. Goyal's

Path Lab & Imaging Centre



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sangar Road, Jaipur-302019
Tale : 0141-2293346, 4049787, 9887049787
Website : www.dr.goyalpathlab.com | E-mail : dr.goyalpiyush@gmail.com

Date :- 23/11/2023 09:29:31 Patient ID :-12234347
NAME :- Mrs. ANJANA JAIN Ref. By Dr:-
Sex / Age :- Female 39 Yrs 3 Mon 23 Days Lab/Hosp :-
Company :- MediWheel



Sample Type :- EDTA Sample Collected Time 23/11/2023 10:20:29 Final Authentication : 23/11/2023 12:28:02

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

BOB PACKAGE FEMALE BELOW 40

GLYCOSYLATED HEMOGLOBIN (HbA1C)

6.0 %

Method:- HPLC

Non-diabetic: < 5.7
Pre-diabetics: 5.7-6.4
Diabetics: = 6.5 or higher
ADA Target: 7.0
Action suggested: > 6.5

Instrument name: ARKRAY's ADAMS Lite HA 8380V, JAPAN.

Test Interpretation:

HbA1C is formed by the condensation of glucose with n-terminal valine residue of each beta chain of HbA to form an unstable schiff base. It is the major fraction, constituting approximately 80% of HbA1c. Formation of glycated hemoglobin (GHb) is essentially irreversible and the concentration in the blood depends on both the lifespan of the red blood cells (RBC) (120 days) and the blood glucose concentration. The GHb concentration represents the integrated values for glucose over the period of 6 to 8 weeks. GHb values are free of day to day glucose fluctuations and are unaffected by recent exercise or food ingestion. Concentration of plasma glucose concentration in GHb depends on the time interval, with more recent values providing a larger contribution than earlier values. The interpretation of GHb depends on RBC having a normal life span. Patients with hemolytic disease or other conditions with shortened RBC survival exhibit a substantial reduction of GHb. High GHb have been reported in iron deficiency anemia. GHb has been firmly established as an index of long term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. The absolute risk of retinopathy and nephropathy are directly proportional to the mean of HbA1C. Genetic variants (e.g. HbS trait, HbC trait), elevated HbF and chemically modified derivatives of hemoglobin can affect the accuracy of HbA1c measurements. The effects vary depending on the specific Hb variant or derivative and the specific HbA1c method.

Ref by ADA 2020

MEAN PLASMA GLUCOSE

126 mg/dL

Method:- Calculated Parameter

Non Diabetic < 100 mg/dL
Prediabetic 100- 125 mg/dL
Diabetic 126 mg/dL or Higher

MUKESH SINGH
Technologist

Page No: 1 of 12



Dr. Chandrika Gupta
MBBS, MD (Path)
RMC NO. 21021/008037

Dr. Goyal's

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanapaner Road, Jaipur-302019
Tele : 0141-2293346, 4049787, 9887049787
Website : www.dr.goyalpathlab.com | E-mail : dr.goyalpiyush@gmail.com

Date :- 23/11/2023 09:29:31 Patient ID :-12234347
NAME :- Mrs. ANJANA JAIN Ref. By Dr:-
Sex / Age :- Female 39 Yrs 3 Mon 23 Days Lab/Hosp :-
Company :- MediWheel



Sample Type > EDTA

Sample Collected Time 23/11/2023 10:20:29

Final Authentication : 23/11/2023 12:28:02

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
Erythrocyte Sedimentation Rate (ESR)	10	mm/hr.	00 - 20

(ESR) Methodology : Measurement of ESR by cells aggregation.

Instrument Name : Independent form Hematocrit value by Automated Analyzer (Roller-20)

Interpretation : ESR test is a non-specific indicator of inflammatory disease and abnormal protein states.

The test is used to detect, follow course of a certain disease (e.g-tuberculosis, rheumatic fever, myocardial infarction)

Levels are higher in pregnancy due to hyperfibrinogenaemia.

The "3-figure ESR" >100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia

(CBC) Methodology : TLC, DLC Fluorescent Flow cytometry, HB SLS method, TRBC, PCV, PLT Hydrodynamically focused Impedance. and

MCH, MCV, MCHC, MENTZER INDEX are calculated. Instrument Name: Sysmex 6 part fully automatic analyzer XN-L, Japan.

MUKESH SINGH
Technologist

Page No: 3 of 12



Dr. Chandrika Gupta
MBBS, MD (Path)
RMC NO. 21021/008037



Date :- 23/11/2023 09:29:31

Patient ID :-12234347

NAME :- Mrs. ANJANA JAIN

Ref. By Dr:-

Sex / Age :- Female 39 Yrs 3 Mon 23 Days

Lab/Hosp :-

Company :- MediWheel



Sample Type > PLAIN/SERUM

Sample Collected Time 23/11/2023 10:20:29

Final Authentication : 23/11/2023 12:14:40

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
LIPID PROFILE			
TOTAL CHOLESTEROL Method:- Enzymatic Endpoint Method	108.37	mg/dl	Desirable <200 Borderline 200-239 High > 240
TRIGLYCERIDES Method:- GPO-PAP	45.56	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500
DIRECT HDL CHOLESTEROL Method:- Direct clearance Method	47.53	mg/dl	Low < 40 High > 60
DIRECT LDL CHOLESTEROL Method:- Direct clearance Method	53.25	mg/dl	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190
VLDL CHOLESTEROL Method:- Calculated	9.11	mg/dl	0.00 - 80.00
T.CHOLESTEROL/HDL CHOLESTEROL RATIO Method:- Calculated	2.28		0.00 - 4.90
LDL / HDL CHOLESTEROL RATIO Method:- Calculated	1.12		0.00 - 3.50
TOTAL LIPID Method:- CALCULATED	309.00 L	mg/dl	400.00 - 1000.00
<p>TOTAL CHOLESTEROL InstrumentName:Randox Rx Imola Interpretation: Cholesterol measurements are used in the diagnosis and treatment of lipid lipoprotein metabolism disorders.</p> <p>TRIGLYCERIDES InstrumentName:Randox Rx Imola Interpretation: Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various metabolic disorders e.g. diabetes mellitus, nephrosis and liver obstruction.</p> <p>DIRECT HDL CHOLESTEROL InstrumentName:Randox Rx Imola Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.</p> <p>DIRECT LDL CHOLESTEROL InstrumentName:Randox Rx Imola Interpretation: Accurate measurement of LDL-Cholesterol is of vital importance in therapies which focus on lipid reduction to prevent atherosclerosis or reduce its progress and to avoid plaque rupture.</p> <p>TOTAL LIPID AND VLDL ARE CALCULATED</p>			

SURENDRAKHANGA

Page No: 4 of 12



Dr. Chandrika Gupta
 MBBS,MD (Path)
 RMC NO. 21021/008037

Dr. Goyal's

Path Lab & Imaging Centre



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019
 Tele : 0141-2293346, 4048787, 9887049787
 Website : www.drgoyalpathlab.com | E-mail : drgoyalpiyush@gmail.com

Date :- 23/11/2023 09:29:31 Patient ID :-12234347
NAME :- Mrs. ANJANA JAIN Ref. By Dr:-
 Sex / Age :- Female 39 Yrs 3 Mon 23 Days Lab/Hosp :-
 Company :- Med/Wheel



Sample Type :- PLAIN/SERUM Sample Collected Time 23/11/2023 10:20:29 Final Authentication : 23/11/2023 12:14:40

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
LIVER PROFILE WITH GGT			
SERUM BILIRUBIN (TOTAL) Method:- Colorimetric method	0.33	mg/dl	Up to - 1.0 Cord blood <2 Premature < 6 days <16 Full-term < 6 days= 12 1month - <12 months <2 1-19 years <1.5 Adult - Up to - 1.2 Ref-(ACCP 2020)
SERUM BILIRUBIN (DIRECT) Method:- Colorimetric Method	0.11	mg/dL	Adult - Up to 0.25 Newborn - <0.6 >- 1 month - <0.2
SERUM BILIRUBIN (INDIRECT) Method:- Calculated	0.22	mg/dl	0.30-0.70
SGOT Method:- IFCC	10.9	U/L	Men- Up to - 37.0 Women - Up to - 31.0
SGPT Method:- IFCC	14.4	U/L	Men- Up to - 40.0 Women - Up to - 31.0
SERUM ALKALINE PHOSPHATASE Method:- AMP Buffer	55.50	IU/L	30.00 - 120.00
SERUM GAMMA GT Method:- IFCC	8.00	U/L	7.00 - 32.00
SERUM TOTAL PROTEIN Method:- Biuret Reagent	7.20	g/dl	6.40 - 8.30
SERUM ALBUMIN Method:- Bromocresol Green	4.15	g/dl	3.80 - 5.00
SERUM GLOBULIN Method:- CALCULATION	3.05	gm/dl	2.20 - 3.50
A/G RATIO	1.36		1.30 - 2.50

Total Bilirubin Methodology: Colorimetric method Instrument/Name: Randox Rx Inova Interpretation: An increase in bilirubin concentration in the serum occurs in toxic or infectious diseases of the liver e.g. hepatitis B or obstruction of the bile duct and in those incompatible babies. High levels of unconjugated bilirubin indicate that too much haemoglobin is being destroyed or that the liver is not actively treating the haemoglobin it is receiving.

AST Aspartate Aminotransferase Methodology: IFCC Instrument/Name: Randox Rx Inova Interpretation: Elevated levels of AST can signal myocardial infarction, hepatic disease, muscular dystrophy and organ damage. Although heart muscle is found to have the most activity of the enzyme, significant activity has also been seen in the brain, liver, gastric mucosa, adipose tissue and kidneys of humans.

ALT Alanine Aminotransferase Methodology: IFCC Instrument/Name: Randox Rx Inova Interpretation: The enzyme ALT has been found to be in highest concentrations in the liver, with decreasing concentrations found in kidney, heart, skeletal muscle, pancreas, spleen and lung tissue respectively. Elevated levels of the transaminase can indicate myocardial infarction, hepatic disease, muscular dystrophy and organ damage.

Alkaline Phosphatase Methodology: AMP Buffer Instrument/Name: Randox Rx Inova Interpretation: Measurements of alkaline phosphatase are of use in the diagnosis, treatment and investigation of hepatobiliary disease and in bone disease associated with increased osteoblastic activity. Alkaline phosphatase is also used in the diagnosis of parathyroid and intestinal disease.

TOTAL PROTEIN Methodology: Biuret Reagent Instrument/Name: Randox Rx Inova Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

ALBUMIN (ALB) Methodology: Bromocresol Green Instrument/Name: Randox Rx Inova Interpretation: Albumin measurements are used in the diagnosis and treatment of numerous diseases involving primarily the liver or kidneys. Globulin & A/G ratio is calculated.

Instrument Name Randox Rx Inova Interpretation: Elevations in GGT levels occur earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and metastatic neoplasms. It may reach 5 to 30 times normal levels in intra- or post-hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal).

SURENDRAKHANGA

Page No: 5 of 12



Dr. Chandrika Gupta
 MBBS,MD (Path)
 RMC NO. 21021/008037



Date :- 23/11/2023 09:29:31
NAME :- Mrs. ANJANA JAIN
 Sex / Age :- Female 39 Yrs 3 Mon 23 Days
 Company :- MediWheel

Patient ID :- 12234347
 Ref. By Dr:-
 Lab/Hosp :-



Sample Type :- PLAIN/SERUM

Sample Collected Time 23/11/2023 10:20:29

Final Authentication : 23/11/2023 11:39:19

IMMUNOASSAY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

TOTAL THYROID PROFILE

SERUM TOTAL T3 Method:- Chemiluminescence(Competitive immunoassay)	1.430	ng/ml	0.970 - 1.690
SERUM TOTAL T4 Method:- Chemiluminescence(Competitive immunoassay)	7.960	ug/dl	5.500 - 11.000
SERUM TSH ULTRA Method:- Enhanced Chemiluminescence Immunoassay	1.930	µIU/mL	0.350 - 5.500

Interpretation: Triiodothyronine (T3) contributes to the maintenance of the euthyroid state. A decrease in T3 concentration of up to 50% occurs in a variety of clinical situations, including acute and chronic disease. Although T3 results alone cannot be used to diagnose hypothyroidism, T3 concentration may be more sensitive than thyroxine (T4) for hyperthyroidism. Consequently, the total T3 assay can be used in conjunction with other assays to aid in the differential diagnosis of thyroid disease. T3 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, Free T3 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake, or T4 uptake can be used with the total T3 result to calculate the free T3 index and estimate the concentration of free T3.

Interpretation: The measurement of Total T4 aids in the differential diagnosis of thyroid disease. While >99.9% of T4 is protein-bound, primarily to thyroxine-binding globulin (TBG), it is the free fraction that is biologically active. In most patients, the total T4 concentration is a good indicator of thyroid status. T4 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, free T4 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake may be used with the total T4 result to calculate the free T4 index (FT4I) and estimate the concentration of free T4. Some drugs and some nonthyroidal patient conditions are known to alter TT4 concentrations in vivo.

Interpretation: TSH stimulates the production of thyroxine (T4) and triiodothyronine (T3) by the thyroid gland. The diagnosis of overt hypothyroidism by the finding of a low total T4 or free T4 concentration is readily confirmed by a raised TSH concentration. Measurement of low or undetectable TSH concentrations may assist the diagnosis of hyperthyroidism, where concentrations of T4 and T3 are elevated and TSH secretion is suppressed. These have the advantage of discriminating between the concentrations of TSH observed in thyrotoxicosis, compared with the low, but detectable, concentrations that occur in subclinical hyperthyroidism. The performance of this assay has not been established for neonatal specimens. Some drugs and some nonthyroidal patient conditions are known to alter TSH concentrations in vivo.

INTERPRETATION

PREGNANCY	REFERENCE RANGE FOR TSH IN uIU/mL (As per American Thyroid Association)
1st Trimester	0.10-2.50
2nd Trimester	0.20-3.00
3rd Trimester	0.30-3.00

AJAYKUMAR
Technologist

Page No: 6 of 12



Dr. Chandrika Gupta
 MBBS,MD (Path)
 RMC NO. 21021/008037

Dr. Goyal's

Path Lab & Imaging Centre



B-51, Ganesh Nagar, Opp. Jangpath Corner, New Sanganeer Road, Jaipur-302019
Tele : 0141-2293346, 4049787, 9887049787
Website : www.dr.goyalspathlab.com | E-mail : dr.goyalpiyush@gmail.com

Date :- 23/11/2023 09:29:31
NAME :- Mrs. ANJANA JAIN
Sex / Age :- Female 39 Yrs 3 Mon 23 Days
Company :- MediWheel

Patient ID :-12234347
Ref. By Dr:-
Lab/Hosp :-



Sample Type :- URINE

Sample Collected Time 23/11/2023 10:20:29

Final Authentication : 23/11/2023 11:34:07

CLINICAL PATHOLOGY

Test Name	Value	Unit	Biological Ref Interval
Urine Routine			
PHYSICAL EXAMINATION			
COLOUR	PALE YELLOW		PALE YELLOW
APPEARANCE	Clear		Clear
CHEMICAL EXAMINATION			
REACTION(PH) Method:- Reagent Strip(Double indicator blue reaction)	5.5		5.0 - 7.5
SPECIFIC GRAVITY Method:- Reagent Strip(bromthymol blue)	1.025		1.010 - 1.030
PROTEIN Method:- Reagent Strip (Sulphosalicylic acid test)	NIL		NIL
GLUCOSE Method:- Reagent Strip (Glu.Oxidase Peroxidase Benedict)	NIL		NIL
BILIRUBIN Method:- Reagent Strip (Azo-coupling reaction)	NEGATIVE		NEGATIVE
UROBILINOGEN Method:- Reagent Strip (Modified ehrlich reaction)	NORMAL		NORMAL
KETONES Method:- Reagent Strip (Sodium Nitroprusside) Rothera's	NEGATIVE		NEGATIVE
NITRITE Method:- Reagent Strip (Diazotization reaction)	NEGATIVE		NEGATIVE
MICROSCOPY EXAMINATION			
RBC/HPF	NIL	/HPF	NIL
WBC/HPF	2-3	/HPF	2-3
EPITHELIAL CELLS	2-3	/HPF	2-3
CRYSTALS/HPF	ABSENT		ABSENT
CAST/HPF	ABSENT		ABSENT
AMORPHOUS SEDIMENT	ABSENT		ABSENT
BACTERIAL FLORA	ABSENT		ABSENT
YEAST CELL	ABSENT		ABSENT
OTHER	ABSENT		ABSENT

VIJENDRAMEENA
Technologist

Page No: 7 of 12



Dr. Chandrika Gupta
MBBS.MD (Path)
RMC NO. 21021/008037

Dr. Goyal's

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganeer Road, Jaipur-302019
Tele : 0141-2293346, 4049787, 9887049787
Website : www.drgoyalpathlab.com | E-mail : drgoyalpiyush@gmail.com

Date :- 23/11/2023 09:29:31

Patient ID :-12234347



NAME :- Mrs. ANJANA JAIN

Ref. By Dr:-

Sex / Age :- Female 39 Yrs 3 Mon 23 Days

Lab/Hosp :-

Company :- MediWheel

Sample Type :- EDTA, URINE

Sample Collected Time 23/11/2023 10:20:29

Final Authentication : 23/11/2023 13:26:13

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
BLOOD GROUP ABO	"B" POSITIVE		
BLOOD GROUP ABO Methodology : Haemagglutination reaction. Kit Name : Monoclonal agglutinating antibodies (Span close).			
URINE SUGAR (FASTING) Collected Sample Received	Nil		Nil

MUKESH SINGH, VIJENDRAMEENA
Technologist

Page No: 11 of 12



Dr. Chandrika Gupta
MBBS, MD (Path)
RMC NO. 21021/008037

Dr. Goyal's

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele : 0141-2293346, 4049787, 9887049787

Website : www.drgoyalpathlab.com | E-mail : drgoyalpiyush@gmail.com

Date :- 23/11/2023 09:29:31

Patient ID :-12234347



NAME :- Mrs. ANJANA JAIN

Ref. By Dr:-

Sex / Age :- Female 39 Yrs 3 Mon 23 Days

Lab/Hosp :-

Company :- MediWheel

Sample Type :- PLAIN/SERUM

Sample Collected Time 23/11/2023 10:20:29

Final Authentication : 23/11/2023 12:14:40

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
BLOOD UREA NITROGEN (BUN)	9.1	mg/dl	0.0 - 23.0

*** End of Report ***

SURENDRAKHANGA

Page No: 12 of 12



Dr. Chandrika Gupta
MBBS.MD (Path)
RMC NO. 21021/008037



Date :- 23/11/2023 09:29:31
NAME :- Mrs. ANJANA JAIN
Sex / Age :- Female 39 Yrs 3 Mon 23 Days
Company :- MediWheel

Patient ID :- 12234347
Ref. By Doctor :-
Lab/Hosp :-

Final Authentication : 23/11/2023 11:50:28

BOB PACKAGEFEMALE BELOW 40

X RAY CHEST PA VIEW:

Both lung fields appears clear.

Bronchovascular markings appear normal.

Trachea is in midline.

Both the hilar shadows are normal.

Both the C.P.angles is clear.

Both the domes of diaphragm are normally placed.

Bony cage and soft tissue shadows are normal.

Heart shadows appear normal.

Impression :- Normal Study

(Please correlate clinically and with relevant further investigations)

*** End of Report ***



DR ABHISHEK JAIN
MBBS. DNB. (RADIO DIAGNOSIS)
RMC NO. 21687

Dr. Piyush Goyal
(D.M.R.D.) BILAL

Transcript by.



Date :- 23/11/2023 09:29:31
NAME :- Mrs. ANJANA JAIN
Sex / Age :- Female 39 Yrs 3 Mon 23 Days
Company :- MediWheel

Patient ID :- 12234347
Ref. By Doctor :-
Lab/Hosp :-

Final Authentication : 23/11/2023 11:10:48

BOB PACKAGEFEMALE BELOW 40

ULTRA SOUND SCAN OF ABDOMEN

Liver is of normal size. Echo-texture is normal. No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is of normal size. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape. Echotexture is normal. No focal lesion is seen.

Kidneys are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. No focal lesion is seen. Collecting system does not show any dilatation or calculus.

Urinary Bladder: is well distended and showing smooth wall with normal thickness. Urinary bladder does not show any calculus or mass lesion.

Uterus is anteverted and normal in size and measures 73 x 56 x 42mm.
Myometrium shows normal echo - pattern.

Two small hypoechoic lesions are seen measuring ~ 18x10mm on posterior wall & ~11.6x10.6mm at anterior wall of uterus.

Endometrial echo is normal. Endometrial thickness is 8 mm.

Both ovaries are visualised and are normal. No adnexal mass is seen.
No enlarged nodes are visualised. No retro-peritoneal lesion is identified.
No significant free fluid is seen in pouch of douglas.

IMPRESSION:

- Small intramural uterine fibroids.

Needs clinical correlation.

*** End of Report ***

Page No: 1 of 1

BILAL

Dr. Piyush Goyal
M.B.B.S., D.M.R.D.
RMC Reg No. 017996

Dr. Ashish Choudhary
MBBS, MD (Radio Diagnosis)
Fetal Medicine Consultant
FMF ID - 200517 | RMC No 22430

Dr. Abhishek Jain
MBBS, DNB, (Radio-Diagnosis)
RMC No. 21687

Dr. Navneet Agarwal
MD, DNB (Radio Diagnosis)
RMC No. 33613/14911

Dr. Poorvi Malik
MBBS, MD, DNB (Radio Diagnosis)
RMC No. 21505

Transcript by.

Dr. Goyal's

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele : 0141-2293346, 4049787, 9887049787

Website : www.dr.goyalspathlab.com | E-mail : dr.goyalpiyush@gmail.com

Date :- 23/11/2023 09:29:31

Patient ID :-12234347

NAME :- Mrs. ANJANA JAIN

Ref. By Dr:-

Sex / Age :- Female 39 Yrs 3 Mon 23 Days

Lab/Hosp :-

Company :- MediWheel



Sample Type :-

Sample Collected Time

Final Authentication : 23/11/2023 12:30:48

BOB PACKAGEFEMALE BELOW 40

2D ECHO OPTION TMT (ADULT/CHILD)

2D-ECHOCARDIOGRAPHY M.MODE WITH DOPPLER STUDY:

FAIR TRANSTHORACIC ECHOCARDIOGRAPHIC WINDOW MORPHOLOGY:

MITRAL VALVE	NORMAL	TRICUSPID VALVE	NORMAL
AORTIC VALVE	NORMAL	PULMONARY VALVE	NORMAL

M.MODE EXAMINATION:

AO	252	mm	LA	26	Mm	IVS-D	10	mm
IVS-S	13	mm	LVID	29	Mm	LVSD	27	mm
LVPW-D	9	mm	LVPW-S		Mm	RV		mm
RVWT		mm	EDV		ml	LVVS		ml
LVEF	60%		RWMA		ABSENT			

CHAMBERS:

LA	NORMAL	RA	NORMAL
LV	NORMAL	RV	NORMAL
PERICARDIUM		NORMAL	

COLOUR DOPPLER:

MITRAL VALVE					
E VELOCITY	0.82	m/sec	PEAK GRADIENT		Mm/hg
A VELOCITY	1.00	m/sec	MEAN GRADIENT		Mm/hg
MVA BY PHT		Cm2	MVA BY PLANIMETRY		Cm2
MITRAL REGURGITATION			ABSENT		
AORTIC VALVE					
PEAK VELOCITY	1.18	m/sec	PEAK GRADIENT		mm/hg
AR VMAX		m/sec	MEAN GRADIENT		mm/hg
AORTIC REGURGITATION			ABSENT		
TRICUSPID VALVE					
PEAK VELOCITY	0.55	m/sec	PEAK GRADIENT		mm/hg
MEAN VELOCITY		m/sec	MEAN GRADIENT		mm/hg
VMax VELOCITY					
TRICUSPID REGURGITATION			ABSENT		
PULMONARY VALVE					
PEAK VELOCITY	1.2	M/sec.	PEAK GRADIENT		Mm/hg
MEAN VELOCITY			MEAN GRADIENT		Mm/hg
PULMONARY REGURGITATION			ABSENT		

LAXMI

Page No: 1 of 2



Dr. Goyal's

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Jarpath Corner, New Sangar Road, Jaipur-302019
Tele : 0141-2293346, 4049787, 9887049787
Website : www.drgoyalpathlab.com | E-mail : drgoyalpiyush@gmail.com

Date :- 23/11/2023 09:29:31
NAME :- Mrs. ANJANA JAIN
Sex / Age :- Female 39 Yrs 3 Mon 23 Days
Company :- MediWheel

Patient ID :-12234347
Ref. By Dr:-
Lab/Hosp :-



Sample Type :-

Sample Collected Time

Final Authentication : 23/11/2023 12:30:46

Impression--

1. Normal LV size & contractility.
2. Trace MR ,Trace TR, PASP25mmHg.
3. Normal Diastolic Function.
4. No RWMA, LVEF 60 %.
5. Normal cardiac chamber.
6. Normal valve.
7. No clot, no vegetation, no pericardial effusion.

(Cardiologist)

*** End of Report ***

LAXMI

Page No: 2 of 2



Dr. Goyal's Path Lab

Name Anjana jain 39 yrs
Patient Id ANJAN71_71829

Date 11/23/2023
Diagnosis Dr.

