

CERTIFICATE OF MEDICAL FITNESS

NAME: Mr. Ram Nageppa Mages

AGE/ GENDER: 59/m

HEIGHT: 174 cm

WEIGHT: 79 kg

IDENTIFICATION MARK: -

BLOOD PRESSURE: 130/80 mm/Hg

PULSE: 69/min

CVS:

RS:P } Normal

ANY OTHER DISEASE DIAGNOSED IN THE PAST: Diabetes :- Mixtard 30/70inj
Glucobay m 25mg
Sarloc 40mg Tab

ALLERGIES, IF ANY: Nil

LIST OF PRESCRIBED MEDICINES: Nil

ANY OTHER REMARKS: NO

I Certify that I have carefully examined Mr/Mrs. Rama Mages son/daughter of Mr. Nageppa who has signed in my presence. He/ she has no physical disease and is fit for employment.


Signature of candidate

Dr. SATHI KINI
M.D. (MEDICINE)
Consultant Physician
REG. No. 24012 (K.M.C.)
Signature of Medical Officer

Place: Spectrum diagnostic & health care

Date: 09/08/23

Disclaimer: The patient has not been checked for COVID. This certificate does not relate to the covid status of the patient examined



Dr. Ashok S
Bsc., MBBS., D.O.M.S
Consultant Ophthalmologist
KMC No: 31827

DATE: 09.08.23

EYE EXAMINATION

NAME: *Ms. Ram Narayan* AGE: *59y* GENDER : F / M
mesel.

	RIGHT EYE	LEFT EYE
Vision	<i>6/18:ND</i>	<i>6/18:ND</i>
Vision With glass	<i>6/12:NB</i>	<i>6/12:NB</i>
Color Vision	Normal	Normal
Anterior segment examination	Normal	Normal
Fundus Examination	Normal	Normal
Any other abnormality	Nil	Nil
Diagnosis/ impression	Normal	Normal <i>e</i>

ms. ram narayan
Dr. ASHOK SARODHE
B.Sc., M.B.B.S., D.O.M.S.
Eye Consultant & Surgeon
KMC 31827
Consultant (Ophthalmologist)




MR RAM NAGAPPA MOGER
Male 59years

HR	: 62	bpm
P	: 129	ms
PR	: 180	ms
QRS	: 103	ms
QT/QTc	: 400/409	ms
P/QRS/T	: 40/-12/25	°
RV5/SVI	: 0.970/0.527	mV

Diagnosis Information:

Sinus Rhythm
 OS Wave in lead V1
 Left Atrial Enlargement

Report Confirmed by:



NAME : MR.RAM NAGAPPA MOGER	DATE :09/08/2023
AGE/SEX : 59 YEARS/MALE	REG NO: 0019
REF BY : C/O APOLO CLINIC	

CHEST PA VIEW

Lung fields are clear.

Cardiovascular shadows are within normal limits.

Both CP angles are free.

Domes of diaphragm and bony thoracic cage are normal.

IMPRESSION: NORMAL CHEST RADIOGRAPH.



**DR.RAM PRAKASH G MDRD
CONSULTANT RADIOLOGIST**

Your suggestion / feedback is a valuable input for improving our services



Name	: MR. RAM NAGAPPA MOGER	UHID	: 0908230019	Bill Date	: 09-Aug-2023 08:10 AM
Age / Gender	: 59 years / Male			Sample Col. Date	: 09-Aug-2023 08:10 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 09-Aug-2023 02:27 PM
Reg. No.	: 0908230019			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Complete Haemogram-Whole Blood EDTA				
Haemoglobin (HB)	15.9	g/dL	Male:14.0-17.0	Spectrophotometer
Red Blood Cell (RBC)	5.71	million/cumm	3.50 - 5.50	Volumetric
Packed Cell Volume (PCV)	48.2	%	Male: 42.0-51.0	Impedance
Mean corpuscular volume (MCV)	84.3	fL	78.0- 94.0	Electronic Pulse
Mean corpuscular hemoglobin (MCH)	27.8	pg	27.50-32.20	Calculated
Mean corpuscular hemoglobin concentration (MCHC)	33.0	%	33.00-35.50	Calculated
Red Blood Cell Distribution Width SD (RDW-SD)	35.7	fL	40.0-55.0	Volumetric
Red Blood Cell Distribution CV (RDW-CV)	14.2	%	Male: 11.80-14.50	Impedance
Mean Platelet Volume (MPV)	7.9	fL	8.0-15.0	Volumetric
Platelet	2.0	lakh/cumm	1.50-4.50	Impedance
Platelet Distribution Width (PDW)	14.4	%	8.30 - 56.60	Volumetric
White Blood cell Count (WBC)	6970.0	cells/cumm	Male: 4000.0-11000.0	Impedance
Neutrophils	61.0	%	40.0-75.0	Volumetric
Lymphocytes	30.0	%	20.0-40.0	Impedance
Eosinophils	3.0	%	0.0-6.0	Light scattering/Manual
Monocytes	5.0	%	0.0-8.0	Light scattering/Manual
Basophils	1.0	%	0.0-1.0	Light scattering/Manual
Absolute Neutrophil Count	4.15	10 ³ /uL	2.0- 7.0	Light scattering/Manual
				Calculated



Name : MR. RAM NAGAPPA MOGER	UHID : 0908230019	Bill Date : 09-Aug-2023 08:10 AM
Age / Gender : 59 years / Male	 0908230019	Sample Col. Date : 09-Aug-2023 08:10 AM
Ref. By Dr. : Dr. APOLO CLINIC		Result Date : 09-Aug-2023 02:27 PM
Reg. No. : 0908230019		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Absolute Lymphocyte Count	2.04	10 ³ /uL	1.0-3.0	Calculated
Absolute Monocyte Count	0.51	10 ³ /uL	0.20-1.00	Calculated
Absolute Eosinophil Count	230	cells/cumm	40-440	Calculated
Absolute Basophil Count	0.02	10 ³ /uL	0.0-0.10	Calculated
Erythrocyte Sedimentation Rate (ESR)	10	mm/hr	Male: 0.0-10.0	Westergren

Peripheral Smear Examination-Whole Blood EDTA

Method: (Microscopy-Manual)

RBC'S : Normocytic Normochromic.
WBC'S : Are normal in total number, morphology and distribution.
Platelets : Adequate in number and normal in morphology.
No abnormal cells or hemoparasites are present.
Impression : Normocytic Normochromic Blood picture.



Printed By : spectrum
Printed On : 09 Aug, 2023 04:19 pm



Name	: MR. RAM NAGAPPA MOGER	UHID	: 0908230019	Bill Date	: 09-Aug-2023 08:10 AM
Age / Gender	: 59 years / Male			Sample Col. Date	: 09-Aug-2023 08:10 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 09-Aug-2023 02:27 PM
Reg. No.	: 0908230019			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
RFT (Urea, Creatinine, BUN, Na+, K+, Cl-, RBS Uric acid,HB)				
RFT (Renal Function Test)-				
Serum				
Urea-Serum	25.00	mg/dL	Male: 06 - 40	Urease
Creatinine-Serum	0.97	mg/dL	Male: 0.6 - 1.5	Modified kinetic Jaffe
Blood Urea Nitrogen (BUN)-Serum	11.7	mg/dL	Male: 6 - 20	:GLDH,Kinetic Assay
Sodium (Na+)-Serum	139.4	mmol/L	Male: 135 - 145	ISE-Direct
Potassium (K+)-Serum	4.21	mmol/L	Male: 3.5 - 5.5	ISE-Direct
Chloride (Cl-)-Serum	100.20	mmol/L	94.0 - 110.0	ISE-Direct
Uric Acid-Serum	5.12	mg/dL	Male: 3.50 - 7.20	Uricase PAP



Printed By : spectrum
Printed On : 09 Aug, 2023 04:19 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajaji Nagar, Opp. St. Theresa Hospital, Bangalore - 10

+91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostic Page 3 of 3

SCAN FOR LOCATION



Name	: MR. RAM NAGAPPA MOGER	UHID	: 0908230019	Bill Date	: 09-Aug-2023 08:10 AM
Age / Gender	: 59 years / Male			Sample Col. Date	: 09-Aug-2023 08:10 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 09-Aug-2023 02:27 PM
Reg. No.	: 0908230019			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Lipid Profile-Serum				
Cholesterol Total-Serum	102.00	mg/dL	0.0-200	Cholesterol Oxidase/Peroxidase
Triglycerides-Serum	41.00	mg/dL	0.0-150	Lipase/Glycerol Dehydrogenase
High-density lipoprotein (HDL) Cholesterol-Serum	43.00	mg/dL	40.0-60.0	Accelerator/Selective Detergent
Non-HDL cholesterol-Serum	59	mg/dL	0.0-130	Calculated
Low-density lipoprotein (LDL) Cholesterol-Serum	61.00	mg/dL	0.0-100.0	Cholesterol esterase and cholesterol oxidase
Very-low-density lipoprotein (VLDL) cholesterol-Serum	8	mg/dL	0.0-40	Calculated
Cholesterol/HDL Ratio-Serum	2.37	Ratio	0.0-5.0	Calculated

Interpretation:

Parameter	Desirable	Borderline High	High	Very High
Total Cholesterol	<200	200-239	>240	
Triglycerides	<150	150-199	200-499	>500
Non-HDL cholesterol	<130	160-189	190-219	>220
Low-density lipoprotein (LDL) Cholesterol	<100	100-129	160-189	>190

Comments: As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atherosclerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus, sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.



Printed By : spectrum
Printed On : 09 Aug, 2023 04:19 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

SCAN FOR LOCATION



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajaji Nagar, Opp. St. Theresa Hospital, Bangalore - 10

+91 77604 97644 | 080 2337 1555 | info@spectrumdiagnostics.org | www.spectrumdiagnostics.org Page 4 of 4

Name : MR. RAM NAGAPPA MOGER	UHID : 0908230019	Bill Date : 09-Aug-2023 08:10 AM
Age / Gender : 59 years / Male	 0908230019	Sample Col. Date : 09-Aug-2023 08:10 AM
Ref. By Dr. : Dr. APOLO CLINIC		Result Date : 09-Aug-2023 02:27 PM
Reg. No. : 0908230019		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Fasting Blood Sugar (FBS)- Plasma	129	mg/dL	60.0-110.0	Hexo Kinase

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula $C_6H_{12}O_6$. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high. Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes : Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol ,Dietary – Intake of excessive carbohydrates and foods with high glycemic index ? Exercise in between samples ? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.

Post prandial Blood Glucose (PPBS)-Plasma	190	mg/dL	80.0-150.0	Hexo Kinase
--	-----	-------	------------	-------------

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula $C_6H_{12}O_6$. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high. Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes : Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol ,Dietary – Intake of excessive carbohydrates and foods with high glycemic index ? Exercise in between samples ? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.



Printed By : spectrum
Printed On : 09 Aug, 2023 04:19 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

SCAN FOR LOCATION



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajaji Nagar, Opp. St. Theresa Hospital, Bangalore - 10

+91 77604 97644 | 080 2337 1555 | info@spectrumdiagnostics.org | www.spectrumdiagnostics.org Page 5 of 5

Name : MR. RAM NAGAPPA MOGER	UHD : 0908230019	Bill Date : 09-Aug-2023 08:10 AM
Age / Gender : 59 years / Male	 0908230019	Sample Col. Date : 09-Aug-2023 08:10 AM
Ref. By Dr. : Dr. APOLO CLINIC		Result Date : 09-Aug-2023 02:27 PM
Reg. No. : 0908230019		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Thyroid function tests (TFT)-Serum				
Tri-Iodo Thyronine (T3)-Serum	0.71	ng/mL	0.60-1.81	Chemiluminescence Immunoassay (CLIA)
Thyroxine (T4)-Serum	8.50	µg/dL	5.50-12.10	Chemiluminescence Immunoassay (CLIA)
Thyroid Stimulating Hormone (TSH)-Serum	1.74	µIU/mL	0.35-5.50	Chemiluminescence Immunoassay (CLIA)

Comments: Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population without a clinical suspicion of hyperthyroidism.

Reference range: Cord: (37 Weeks): 0.5-1.41, Children: 1-3 Days: 1.0-7.40, 1-11 Months: 1.05-2.45, 1-5 Years: 1.05-2.69, 6-10 Years: 0.94-2.41, 11-15 Years: 0.82-2.13, Adolescents (16-20 Years): 0.80-2.10

Reference range: Adults: 20-50 Years: 0.70-2.04, 50-90 Years: 0.40-1.81, Reference range in Pregnancy: First Trimester: 0.81-1.90, Second Trimester: 1.0-2.60

Increased Levels: Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG).
Decreased Levels: Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

Comments: Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4). It also helps to monitor treatment of Hyperthyroidism with Thiouracil or other anti-thyroid drugs.

Reference Range: Males: 4.6-10.5, Females: 5.5-11.0, > 60 Years: 5.0-10.70, Cord: 7.40-13.10, Children: 1-3 Days: 11.80-22.60, 1-2 Weeks: 9.90-16.60, 1-4 Months: 7.20-14.40, 1-5 Years: 7.30-15.0, 5-10 Years: 6.4-13.3, 1-15 Years: 5.60-11.70, Newborn Screen: 1-5 Days: >7.5, 6 Days: >6.5

Increased Levels: Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia, Increased transthyretin, estrogen therapy, pregnancy.
Decreased Levels: Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.

Comments: TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, caloric intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after ~ 3 weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality.

Reference range in Pregnancy: I- trimester: 0.1-2.5; II -trimester: 0.2-3.0; III- trimester: 0.3-3.0
Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks: 1.7-9.1

Increased Levels: Primary hypothyroidism, Subclinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance.
Decreased Levels: Graves disease, Autonomous thyroid hormone secretion, TSH deficiency.



Printed By : spectrum
Printed On : 09 Aug, 2023 04:19 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist

SCAN FOR LOCATION



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajaji Nagar, Opp. St. Theresa Hospital, Bangalore - 10

+91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostics.org Page 6 of 6

Name : MR. RAM NAGAPPA MOGER	UHID : 0908230019	Bill Date : 09-Aug-2023 08:10 AM
Age / Gender : 59 years / Male		Sample Col. Date : 09-Aug-2023 08:10 AM
Ref. By Dr. : Dr. APOLO CLINIC	0908230019	Result Date : 09-Aug-2023 02:27 PM
Reg. No. : 0908230019		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
LFT-Liver Function Test -Serum				
Bilirubin Total-Serum	1.00	mg/dL	0.2-1.0	Caffeine Benzoate
Bilirubin Direct-Serum	0.20	mg/dL	0.0-0.2	Diazotised Sulphanilic Acid
Bilirubin Indirect-Serum	0.80	mg/dL	0.0-1.10	Direct Measure
Aspartate Aminotransferase (AST/SGOT)-Serum	20.00	U/L	15.0-37.0	UV with Pyridoxal - 5 - Phosphate
Alanine Aminotransferase (ALT/SGPT)-Serum	21.00	U/L	16.0-63.0	UV with Pyridoxal - 5 - Phosphate
Alkaline Phosphatase (ALP)-Serum	86.00	U/L	45.0-117.0	PNPP,AMP-Buffer
Protein, Total-Serum	6.57	g/dL	6.40-8.20	Biuret/Endpoint-With Blank
Albumin-Serum	3.92	g/dL	3.40-5.00	Bromocresol Purple
Globulin-Serum	2.65	g/dL	2.0-3.50	Calculated
Albumin/Globulin Ratio-Serum	1.48	Ratio	0.80-1.20	Calculated



Printed By : spectrum
Printed On : 09 Aug, 2023 04:19 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

SCAN FOR LOCATION



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajaji Nagar, Opp. St. Theresa Hospital, Bangalore - 10

+91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostic Page 7 of 15

Name : MR. RAM NAGAPPA MOGER	UHD : 0908230019	Bill Date : 09-Aug-2023 08:10 AM
Age / Gender : 59 years / Male	 0908230019	Sample Col. Date : 09-Aug-2023 08:10 AM
Ref. By Dr. : Dr. APOLO CLINIC		Result Date : 09-Aug-2023 02:27 PM
Reg. No. : 0908230019		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Glycosylated Haemoglobin (HbA1c)-Whole Blood EDTA				
Glycosylated Haemoglobin (HbA1c)	6.80	%	Non diabetic adults : <5.7 At risk (Prediabetes) : 5.7 - 6.4 Diagnosing Diabetes : >= 6.5 Diabetes Excellent Control : 6-7 Fair to good Control : 7-8 Unsatisfactory Control : 8-10 Poor Control : >10	HPLC
Estimated Average Glucose(eAG)	148.46	mg/dL		Calculated

Note: 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

Comments: HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.



Printed By : spectrum
Printed On : 09 Aug, 2023 04:19 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist

SCAN FOR LOCATION



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajaji Nagar, Opp. St. Theresa Hospital, Bangalore - 10

+91 77604 97644 | 080 2337 1555 | info@spectrumdiagnostics.org | www.spectrumdiagnostics.org Page 8 of 8

Name : MR. RAM NAGAPPA MOGER	Bill Date : 09-Aug-2023 08:10 AM
Age / Gender : 59 years / Male	Sample Col. Date : 09-Aug-2023 08:10 AM
Ref. By Dr. : Dr. APOLO CLINIC	Result Date : 09-Aug-2023 02:27 PM
Reg. No. : 0908230019	Report Status : Final
C/o : Apollo Clinic	

Test Name	Result	Unit	Reference Value	Method
Gamma-Glutamyl Transferase (GGT)-Serum	13.00	U/L	Male: 15.0-85.0	Other g-Glut-3-carboxy-4 nitro

Comments: Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system, and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of serum GGT levels in detecting alcohol-induced liver disease. Elevated serum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.



Printed By : spectrum
Printed On : 09 Aug, 2023 04:19 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajaji Nagar, Opp. St. Theresa Hospital, Bangalore - 10



Name	: MR. RAM NAGAPPA MOGER	UHID	: 0908230019	Bill Date	: 09-Aug-2023 08:10 AM
Age / Gender	: 59 years / Male			Sample Col. Date	: 09-Aug-2023 08:10 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 09-Aug-2023 02:27 PM
Reg. No.	: 0908230019			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Urine Routine Examination-Urine				
Physical Examination				
Colour	Pale Yellow		Pale Yellow	Visual
Appearance	Clear		Clear	Visual
Reaction (pH)	5.5		5.0 - 7.5	Dipstick
Specific Gravity	1.020		1.000 - 1.030	Dipstick
Biochemical Examination				
Albumin	Negative		Negative	Dipstick/Precipitation
Glucose	Positive (+++)		Negative	Dipstick/Benedicts
Bilirubin	Negative		Negative	Dipstick/Fouchets
Ketone Bodies	Negative		Negative	Dipstick/Rotheras
Urobilinogen	Normal		Normal	Dipstick/Ehrlichs
Nitrite	Negative		Negative	Dipstick
Microscopic Examination				
Pus Cells	2-3	hpf	0.0 - 5.0	Microscopy
Epithelial Cells	1-2	hpf	0.0 - 10.0	Microscopy
RBCs	Absent	hpf	Absent	Microscopy
Casts	Absent		Absent	Microscopy
Crystals	Absent		Absent	Microscopy
Others	Absent		Absent	Microscopy

Comments: The kidneys help infiltration of the blood by eliminating waste out of the body through urine. They also regulate water in the body by conserving electrolytes, proteins, and other compounds. But due to some conditions and abnormalities in kidney function, the urine may encompass some abnormal constituents, which are not normally present. A complete urine examination helps in detecting such abnormal constituents in urine. Several disorders can be detected by identifying and measuring the levels of such substances. Blood cells, bilirubin, bacteria, pus cells, epithelial cells may be present in urine due to kidney disease or infection. Routine urine examination helps to diagnose kidney diseases, urinary tract infections, diabetes and other metabolic disorders.



Printed By : spectrum
Printed On : 09 Aug, 2023 04:19 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

SCAN FOR LOCATION



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajaji Nagar, Opp. St. Theresa Hospital, Bangalore - 10

+91 77604 97644 | 080 2337 1555 | info@spectrumdiagnostics.org | www.spectrumdiagnostics.org Page 0 of 1

Name : MR. RAM NAGAPPA MOGER	UHID : 0908230019	Bill Date : 09-Aug-2023 08:10 AM
Age / Gender : 59 years / Male	 0908230019	Sample Col. Date : 09-Aug-2023 08:10 AM
Ref. By Dr. : Dr. APOLO CLINIC		Result Date : 09-Aug-2023 02:27 PM
Reg. No. : 0908230019		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Blood Group & Rh Typing-Whole Blood EDTA				
Blood Group	O			Slide/Tube agglutination
Rh Type	Positive			Slide/Tube agglutination

Note: Confirm by tube or gel method.

Comments: ABO blood group system, the classification of human blood based on the inherited properties of red blood cells (erythrocytes) as determined by the presence or absence of the antigens A and B, which are carried on the surface of the red cells. Persons may thus have type A, type B, type O, or type AB blood.



Printed By : spectrum
Printed On : 09 Aug, 2023 04:19 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajaji Nagar, Opp. St. Theresa Hospital, Bangalore - 10

+91 77604 97644 | 080 2337 1555  info@spectrumdiagnostics.org  www.spectrumdiagnostics.org Page 1 of 1

SCAN FOR LOCATION



CERTIFICATE OF MEDICAL FITNESS

NAME: Mrs. Leela

AGE/ GENDER: 50/F

HEIGHT: 154 cm

WEIGHT: 80 kg

IDENTIFICATION MARK: -

BLOOD PRESSURE: 120/80 mm Hg

PULSE: 72/min

CVS: Normal

RS:P

ANY OTHER DISEASE DIAGNOSED IN THE PAST: Diabetes, Cyber m 60xR Tub

ALLERGIES, IF ANY:

Hyperkalemia, Glucobay, 40mg Tub
Tazloc, 40mg Tub
eyemal Tub.

LIST OF PRESCRIBED MEDICINES:

ANY OTHER REMARKS:

I Certify that I have carefully examined Mr/Mrs. Leela Rama Mager son/daughter of Ms Naggappa who has signed in my presence. He/ she has no physical disease and is fit for employment.

Dr. SATISH KINI
(M.D. MEDICINE)
Consultant Physician
REG. No. 24012 (K.M.C.)

Leela Rama
Signature of candidate

Signature of Medical Officer

Place: Spectrum diagnostic & health care

Date: 09/08/23

Disclaimer: The patient has not been checked for COVID. This certificate does not relate to the covid status of the patient examined



Dr. Ashok S
Bsc., MBBS., D.O.M.S
Consultant Ophthalmologist
KMC No: 31827

DATE: 09-08-23

EYE EXAMINATION

NAME: *Ms. Leela* AGE: *50y* GENDER: F / M

	RIGHT EYE	LEFT EYE
Vision	<i>6/18: N10</i>	<i>6/18: N10</i>
Vision With glass	<i>6/6: N</i>	<i>6/6: N</i>
Color Vision	Normal	Normal
Anterior segment examination	Normal	Normal
Fundus Examination	Normal	Normal
Any other abnormality	Nil	Nil
Diagnosis/ impression	Normal	Normal

DR. ASHOK SARODHE
B.Sc., M.B.B.S., D.O.M.S.
Eye Consultant & Surgeon
KMC 31827
Consultant (Ophthalmologist)



NAME	AGE	GENDER
Mrs. Leela	50 yrs	female.

DENTAL EXAMINATION REPORT:

8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8
8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8

C: CAVITY → None

M: MISSING → None

O: OTHERS

ADVISED:

CLEANING / SCALING / ROOTS PLANNING / FLOSSING & POLISHING / OTHERS

REMARKS:

SIGNATURE OF THE DENTAL SURGEON

SEAL

Dr. SACHDEV NAGARKAR
B.D.S., F.A.G.E., F.P.F.A. (USA)
Reg. No : 2247/A

DATE

09/08/23



HR : 67 bpm

P : 106 ms

PR : 143 ms

QRS : 95 ms

QT/QTc : 386/408 ms

P/QRS/T : -1/46/61 °


RV5/SV1 : 1.514/1.139 mV

Diagnosis Information:

Sinus Rhythm

Normal ECG

Report Confirmed by:



NAME AND LAB NO	MRS LEELA	Reg : 30021
AGE & SEX	50 YRS	FEMALE
DATE AND AREA OF INTEREST	09.08.2023	ABDOMEN & PELVIS
REF BY	C/O APOLO CLINIC	

USG ABDOMEN AND PELVIS

LIVER: Measures 15.1cm. Normal in size with echotexture. No e/o IHBR dilatation. No evidence of SOL. Portal vein appears normal. CBD appears normal. . No e/o calculus / SOL

GALL BLADDER: Well distended. Wall appears normal.No e/o calculus/ neoplasm.

SPLEEN: Measures 8.8 cm. Normal in size and echotexture. No e/o SOL/ calcification.

PANCREAS: Normal in size and echotexture. Pancreatic duct appears normal. No e/o calculus / calcifications.

RETROPERITONEUM: Poor window.

RIGHT KIDNEY: Measures 9.8 X4.4 cm. Right kidney is normal in size & echotexture No evidence of calculus/ hydronephrosis.

LEFT KIDNEY: Measures 9.5 X 5.5 cm .Left kidney is normal in size & echotexture No evidence of calculus/ hydronephrosis.

URETERS: Bilateral ureters are not dilated.

URINARY BLADDER: Well distended. No wall thickening/ calculi.

UTERUS: Anteverted,Normal in size and echotexture Endometrium is normal.ET -6 mm.

OVARIES: Right ovary - normal in size and echotexture. Left ovary – Obscured by bowel gas shadows.

- No evidence of ascites/pleural effusion.

IMPRESSION:

- No significant sonological abnormality detected in the abdomen and pelvis.


DR AKSHATHA R BHAT
MDRD DNB FRCR



Name	: MRS. LEELA	UHID	: 0908230021	Bill Date	: 09-Aug-2023 08:16 AM
Age / Gender	: 50 years / Female			Sample Col. Date	: 09-Aug-2023 08:16 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 09-Aug-2023 02:28 PM
Reg. No.	: 0908230021			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
RFT (Urea, Creatinine, BUN, Na+, K+, Cl-, RBS Uric acid,HB)				
RFT (Renal Function Test)- Serum				
Urea-Serum	13.00	mg/dL	Female: 06 - 40	Urease
Creatinine-Serum	0.70	mg/dL	Female: 0.5 - 1.1	Modified kinetic Jaffe
Blood Urea Nitrogen (BUN)-Serum	6.1	mg/dL	Female: 6 - 20	:GLDH,Kinetic Assay
Sodium (Na+)-Serum	139.5	mmol/L	Female: 135 - 145	ISE-Direct
Potassium (K+)-Serum	4.03	mmol/L	Female: 3.5 - 5.5	ISE-Direct
Chloride (Cl-)-Serum	100.70	mmol/L	94.0 - 110.0	ISE-Direct
Uric Acid-Serum	3.31	mg/dL	Female: 2.60 - 6.00	Uricase PAP



Printed By : spectrum
Printed On : 09 Aug, 2023 04:24 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajaji Nagar, Opp. St. Theresa Hospital, Bangalore - 10

+91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostics. Page 1 of 1

SCAN FOR LOCATION



Name : MRS. LEELA	Uhid : 0908230021	Bill Date : 09-Aug-2023 08:16 AM
Age / Gender : 50 years / Female	 0908230021	Sample Col. Date : 09-Aug-2023 08:16 AM
Ref. By Dr. : Dr. APOLO CLINIC		Result Date : 09-Aug-2023 02:28 PM
Reg. No. : 0908230021		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Lipid Profile-Serum				
Cholesterol Total-Serum	132.00	mg/dL	0.0-200	Cholesterol Oxidase/Peroxidase
Triglycerides-Serum	108.00	mg/dL	0.0-150	Lipase/Glycerol Dehydrogenase
High-density lipoprotein (HDL) Cholesterol-Serum	41.00	mg/dL	40.0-60.0	Accelerator/Selective Detergent
Non-HDL cholesterol-Serum	91	mg/dL	0.0-130	Calculated
Low-density lipoprotein (LDL) Cholesterol-Serum	70.00	mg/dL	0.0-100.0	Cholesterol esterase and cholesterol oxidase
Very-low-density lipoprotein (VLDL) cholesterol-Serum	22	mg/dL	0.0-40	Calculated
Cholesterol/HDL Ratio-Serum	3.22	Ratio	0.0-5.0	Calculated

Interpretation:

Parameter	Desirable	Borderline High	High	Very High
Total Cholesterol	<200	200-239	>240	
Triglycerides	<150	150-199	200-499	>500
Non-HDL cholesterol	<130	160-189	190-219	>220
Low-density lipoprotein (LDL) Cholesterol	<100	100-129	160-189	>190

Comments: As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atherosclerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus, sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.



Printed By : spectrum
Printed On : 09 Aug, 2023 04:24 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

SCAN FOR LOCATION



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajaji Nagar, Opp. St. Theresa Hospital, Bangalore - 10

+91 77604 97644 | 080 2337 1555 | info@spectrumdiagnostics.org | www.spectrumdiagnostics.org Page 2 of 2

Name : MRS. LEELA	UHID : 0908230021	Bill Date : 09-Aug-2023 08:16 AM
Age / Gender : 50 years / Female		Sample Col. Date : 09-Aug-2023 08:16 AM
Ref. By Dr. : Dr. APOLO CLINIC	0908230021	Result Date : 09-Aug-2023 02:28 PM
Reg. No. : 0908230021		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Thyroid function tests (TFT)-Serum				
Tri-Iodo Thyronine (T3)-Serum	1.02	ng/mL	0.60-1.81	Chemiluminescence Immunoassay (CLIA)
Thyroxine (T4)-Serum	10.5	µg/dL	5.50-12.10	Chemiluminescence Immunoassay (CLIA)
Thyroid Stimulating Hormone (TSH)-Serum	2.03	µIU/mL	0.35-5.50	Chemiluminescence Immunoassay (CLIA)

Comments: Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population without a clinical suspicion of hyperthyroidism.

Reference range: Cord: (37 Weeks): 0.5-1.41, Children: 1-3 Days: 1.0-7.40, 1-11 Months: 1.05-2.45, 1-5 Years: 1.05-2.69, 6-10 Years: 0.94-2.41, 11-15 Years: 0.82-2.13, Adolescents (16-20 Years): 0.80-2.10

Reference range: Adults: 20-50 Years: 0.70-2.04, 50-90 Years: 0.40-1.81, Reference range in Pregnancy: First Trimester : 0.81-1.90, Second Trimester : 1.0-2.60

Increased Levels: Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG).
Decreased Levels: Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

Comments: Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4). It also helps to monitor treatment of Hyperthyroidism with Thiouracil or other anti-thyroid drugs.

Reference Range: Males : 4.6-10.5, Females : 5.5-11.0, > 60 Years: 5.0-10.70, Cord : 7.40-13.10, Children: 1-3 Days : 11.80-22.60, 1-2 Weeks : 9.90-16.60, 1-4 Months: 7.20-14.40, 1-5 Years : 7.30-15.0, 5-10 Years: 6.4-13.3

1-15 Years: 5.60-11.70, Newborn Screen: 1-5 Days: >7.5, 6 Days : >6.5
Increased Levels: Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia, Increased transthyretin, estrogen therapy, pregnancy.
Decreased Levels: Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.

Comments: TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, caloric intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after ~ 3 weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality.

Reference range in Pregnancy: I- trimester: 0.1-2.5; II -trimester: 0.2-3.0; III- trimester: 0.3-3.0
Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks: 1.7-9.1

Increased Levels: Primary hypothyroidism, Subclinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance.
Decreased Levels: Graves disease, Autonomous thyroid hormone secretion, TSH deficiency.



Printed By : spectrum
Printed On : 09 Aug, 2023 04:24 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajaji Nagar, Opp. St. Theresa Hospital, Bangalore - 10



Name	: MRS. LEELA	UHID	: 0908230021	Bill Date	: 09-Aug-2023 08:16 AM
Age / Gender	: 50 years / Female			Sample Col. Date	: 09-Aug-2023 08:16 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 09-Aug-2023 02:28 PM
Reg. No.	: 0908230021			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Glycosylated Haemoglobin (HbA1c)-Whole Blood EDTA	6.80	%	Non diabetic adults :<5.7 At risk (Prediabetes) : 5.7 - 6.4 Diagnosing Diabetes :>= 6.5 Diabetes Excellent Control : 6-7 Fair to good Control : 7-8 Unsatisfactory Control :8-10 Poor Control :>10	HPLC
Estimated Average Glucose(eAG)	148.46	mg/dL		Calculated

Note: 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

Comments: HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.



Printed By : spectrum
Printed On : 09 Aug, 2023 04:24 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

SCAN FOR LOCATION



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajaji Nagar, Opp. St. Theresa Hospital, Bangalore - 10

+91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostics.org Page 4 of 4

Name : MRS. LEELA	UHID : 0908230021	Bill Date : 09-Aug-2023 08:16 AM
Age / Gender : 50 years / Female		Sample Col. Date : 09-Aug-2023 08:16 AM
Ref. By Dr. : Dr. APOLO CLINIC	0908230021	Result Date : 09-Aug-2023 02:28 PM
Reg. No. : 0908230021		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Complete Haemogram-Whole Blood EDTA				
Haemoglobin (HB)	12.2	g/dL	Female:12.0-15.0	Spectrophotometer
Red Blood Cell (RBC)	4.60	million/cumm	3.50 - 5.50	Volumetric Impedance
Packed Cell Volume (PCV)	37.2	%	Female: 36.0-45.0	Electronic Pulse
Mean corpuscular volume (MCV)	80.9	fL	78.0- 94.0	Calculated
Mean corpuscular hemoglobin (MCH)	26.5	pg	27.50-32.20	Calculated
Mean corpuscular hemoglobin concentration (MCHC)	32.8	%	33.00-35.50	Calculated
Red Blood Cell Distribution Width SD (RDW-SD)	33.6	fL	40.0-55.0	Volumetric Impedance
Red Blood Cell Distribution CV (RDW-CV)	14.5	%	Female: 12.20-16.10	Volumetric Impedance
Mean Platelet Volume (MPV)	9.0	fL	8.0-15.0	Volumetric Impedance
Platelet	2.3	lakh/cumm	1.50-4.50	Volumetric Impedance
Platelet Distribution Width (PDW)	15.6	%	8.30 - 56.60	Volumetric Impedance
White Blood cell Count (WBC)	8020.0	cells/cumm	Female: 4000.0-11000.0	Volumetric Impedance
Neutrophils	63.0	%	40.0-75.0	Light scattering/Manual
Lymphocytes	23.0	%	20.0-40.0	Light scattering/Manual
Eosinophils	6.0	%	0.0-6.0	Light scattering/Manual
Monocytes	7.0	%	0.0-8.0	Light scattering/Manual
Basophils	1.0	%	0.0-1.0	Light scattering/Manual
Absolute Neutrophil Count	4.50	10 ³ /uL	2.0- 7.0	Calculated



Name : MRS. LEELA	UHID : 0908230021	Bill Date : 09-Aug-2023 08:16 AM
Age / Gender : 50 years / Female		Sample Col. Date : 09-Aug-2023 08:16 AM
Ref. By Dr. : Dr. APOLO CLINIC	0908230021	Result Date : 09-Aug-2023 02:28 PM
Reg. No. : 0908230021		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Absolute Lymphocyte Count	2.75	10 ³ /uL	1.0-3.0	Calculated
Absolute Monocyte Count	0.45	10 ³ /uL	0.20-1.00	Calculated
Absolute Eosinophil Count	300	cells/cumm	40-440	Calculated
Absolute Basophil Count	0.02	10 ³ /uL	0.0-0.10	Calculated
Erythrocyte Sedimentation Rate (ESR)	13	mm/hr	Female: 0.0-20.0	Westergren

Peripheral Smear Examination-Whole Blood EDTA

Method: (Microscopy-Manual)

- RBC'S : Normocytic Normochromic.
- WBC'S : Are normal in total number, morphology and distribution.
- Platelets : Adequate in number and normal in morphology.
No abnormal cells or hemoparasites are present.
- Impression : Normocytic Normochromic Blood picture.



Printed By : spectrum
Printed On : 09 Aug, 2023 04:24 pm



Name	: MRS. LEELA	UHID	: 0908230021	Bill Date	: 09-Aug-2023 08:16 AM
Age / Gender	: 50 years / Female			Sample Col. Date	: 09-Aug-2023 08:16 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 09-Aug-2023 02:28 PM
Reg. No.	: 0908230021			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
LFT-Liver Function Test -Serum				
Bilirubin Total-Serum	0.84	mg/dL	0.2-1.0	Caffeine Benzoate
Bilirubin Direct-Serum	0.16	mg/dL	0.0-0.2	Diazotised Sulphanilic Acid
Bilirubin Indirect-Serum	0.68	mg/dL	0.0-1.10	Direct Measure
Aspartate Aminotransferase (AST/SGOT)-Serum	18.00	U/L	15.0-37.0	UV with Pyridoxal - 5 - Phosphate
Alanine Aminotransferase (ALT/SGPT)-Serum	20.00	U/L	14.0-59.0	UV with Pyridoxal - 5 - Phosphate
Alkaline Phosphatase (ALP)-Serum	82.00	U/L	45.0-117.0	PNPP,AMP-Buffer
Protein, Total-Serum	6.86	g/dL	6.40-8.20	Biuret/Endpoint-With Blank
Albumin-Serum	4.05	g/dL	3.40-5.00	Bromocresol Purple
Globulin-Serum	2.81	g/dL	2.0-3.50	Calculated
Albumin/Globulin Ratio-Serum	1.44	Ratio	0.80-1.20	Calculated



Printed By : spectrum
Printed On : 09 Aug, 2023 04:24 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajaji Nagar, Opp. St. Theresa Hospital, Bangalore - 10

+91 77604 97644 | 080 2337 1555 | info@spectrumdiagnostics.org | www.spectrumdiagnostics.org Page 7 of 7

SCAN FOR LOCATION



Name	: MRS. LEELA	UHID	: 0908230021	Bill Date	: 09-Aug-2023 08:16 AM
Age / Gender	: 50 years / Female			Sample Col. Date	: 09-Aug-2023 08:16 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 09-Aug-2023 02:28 PM
Reg. No.	: 0908230021			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Fasting Blood Sugar (FBS)- Plasma	122	mg/dL	60.0-110.0	Hexo Kinase

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula $C_6H_{12}O_6$. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high. Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes : Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol ,Dietary – Intake of excessive carbohydrates and foods with high glycemic index ? Exercise in between samples ? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.

Gamma-Glutamyl Transferase (GGT)-Serum	14.00	U/L	Female: 5.0-55.0	Other g-Glut-3-carboxy-4 nitro
--	-------	-----	------------------	--------------------------------

Comments: Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system, and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of serum GGT levels in detecting alcohol-induced liver disease. Elevated serum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.



Printed By : spectrum
Printed On : 09 Aug, 2023 04:24 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajaji Nagar, Opp. St. Theresa Hospital, Bangalore - 10

+91 77604 97644 | 080 2337 1555 | info@spectrumdiagnostics.org | www.spectrumdiagnostics.org Page 8 of 8



Name	: MRS. LEELA	UHID	: 0908230021	Bill Date	: 09-Aug-2023 08:16 AM
Age / Gender	: 50 years / Female			Sample Col. Date	: 09-Aug-2023 08:16 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 09-Aug-2023 02:28 PM
Reg. No.	: 0908230021	0908230021		Report Status	: Final
C/o	: Apollo Clinic				


Test Name	Result	Unit	Reference Value	Method
Blood Group & Rh Typing-Whole Blood EDTA				
Blood Group	B			Slide/Tube agglutination
Rh Type	Positive			Slide/Tube agglutination

Note: Confirm by tube or gel method.

Comments: ABO blood group system, the classification of human blood based on the inherited properties of red blood cells (erythrocytes) as determined by the presence or absence of the antigens A and B, which are carried on the surface of the red cells. Persons may thus have type A, type B, type O, or type AB blood.



Printed By : spectrum
Printed On : 09 Aug, 2023 04:24 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajaji Nagar, Opp. St. Theresa Hospital, Bangalore - 10

+91 77604 97644 | 080 2337 1555  info@spectrumdiagnostics.org  www.spectrumdiagnostics.org Page 9 of 9

SCAN FOR LOCATION

