



भारत सरकार
Unique Identification
Authority of India
Government
of India

E-Aadhaar Letter

नगरपालिका/Enrollment No: 1050/0833767706

Indira Chaudhary (उत्तर लिंगी)
W/O Narendra Kumar Bhadya, 40, Mohalia poliwala,
Bathod, Sikar,
Rajasthan - 332301

Date 05/01/2014

6386 6912 0570



भारत सरकार/Your Aadhaar No.
www.aadhaar.gov.in

- आधार देश भर में मान्य है।
- आधार के लिए आपको एक ही बार नामांकन दर्ज करवाने की आवश्यकता है।
- कृपया अपना नवीनतम मोबाइल नंबर तथा ई-मेल पता दर्ज कराएं। इससे आपको विभिन्न सुविधाएं प्राप्त करने में सक्षम बनने में मदद मिलेगी।

भारत सरकार
GOVERNMENT OF INDIA



इन्द्रा चौधरी
Indira Chaudhary
जन्म तिथि/DOB: 10/06/1986
लिंगी / FEMALE



नगरपालिका

- आधार पहचान का प्रमाण है, नागरिकता का नहीं।
- आधार का प्रमाण ऑनलाइन और ऑनलाइन नहीं प्राप्त किया जा सकता।
- आधार एक इलेक्ट्रॉनिक रूप से उत्पन्न होता है।

INFORMATION

- Aadhaar is a proof of identity, not of citizenship.
- To establish identity, authenticate online.
- This is electronically generated letter.

- Aadhaar is valid throughout the country.
- You need to enrol only once for Aadhaar.
- Please update your mobile number and e-mail address. This will help you to avail various services in future.

भारतीय विशिष्ट पहचान प्राधिकरण
UNIQUE IDENTIFICATION AUTHORITY OF INDIA

पता:
W/O नरेंद्र कुमार भडिया,
40, मोहलिया पोलीवाला,
बाथोड, सिकर,
राजस्थान - 332301
Address:
W/O Narendra Kumar Bhadya, 40,
Mohalia poliwala, Bathod, Sikar,
Rajasthan - 332301

Dr. U. C. GUPTA
MBBS, MD (Physician)
RMC No. 291

Signature

आधार-आम आदमी का अधिकार
0570

Aadhaar-Aam Admi ka Adhikar
0570



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com

General Physical Examination

Date of Examination: 03/06/2023

Name: INDRA CHAUDHARY Age: 36 YRS DOB: 10/06/1986 Sex: Female

Referred By: BANK OF BARODA

Photo ID: AADHAR ID #: 0570

Ht: 161 (cm)

Wt: 66 (Kg)

Chest (Expiration): 88 (cm)

Abdomen Circumference: 75 (cm)

Blood Pressure: 110/65 mm Hg

PR: 79 / min

RR: 18 / min

Temp: Afebrile

BMI 25.5

Eye Examination:

| | | | |
|----|-----|-----|-----|
| RE | 6/6 | N/6 | NCB |
| LE | 6/6 | N/6 | NCB |

Other: No

On examination he/she appears physically and mentally fit: Yes / No

Signature Of Examinee : [Signature]

Name of Examinee: INDRA CHAUDHARY

Signature Medical Examiner : [Signature]

Name Medical Examiner DR. U.C. GUPTA

Dr. U. C. GUPTA
MBBS, MD (Physician)
RMC No. 291



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave - II, Near Axis Bank
 Central Spine, Vidhyadhar Nagar, Jaipur - 302023
 +91 141 4824885 maxcarediagnostics1@gmail.com



| | | | |
|-------------------------------------|--------------------------------|--------------------|----------|
| NAME :- Mrs. INDRA CHAUDHARY | Patient ID :-1223409 | Date :- 03/06/2023 | 09:39:25 |
| Age :- 36 Yrs 11 Mon 24 Days | Ref. By Doctor:-BANK OF BARODA | | |
| Sex :- Female | Lab/Hosp :- | | |
| | Company :- | Mr.MEDIWHEEL | |

Final Authentication : 03/06/2023 17:14:20

HAEMATOLOGY

| Test Name | Value | Unit | Biological Ref Interval |
|---|-------|---------------------------|-------------------------|
| FULL BODY HEALTH CHECKUP BELOW 40 FEMAL | | | |
| HAEMOGARAM | | | |
| HAEMOGLOBIN (Hb) | 12.1 | g/dL | 12.0 - 15.0 |
| TOTAL LEUCOCYTE COUNT | 4.50 | /cumm | 4.00 - 10.00 |
| DIFFERENTIAL LEUCOCYTE COUNT | | | |
| NEUTROPHIL | 55.0 | % | 40.0 - 80.0 |
| LYMPHOCYTE | 40.0 | % | 20.0 - 40.0 |
| EOSINOPHIL | 2.0 | % | 1.0 - 6.0 |
| MONOCYTE | 3.0 | % | 2.0 - 10.0 |
| BASOPHIL | 0.0 | % | 0.0 - 2.0 |
| TOTAL RED BLOOD CELL COUNT (RBC) | 4.46 | $\times 10^6/\mu\text{L}$ | 3.80 - 4.80 |
| HEMATOCRIT (HCT) | 37.30 | % | 36.00 - 46.00 |
| MEAN CORP VOLUME (MCV) | 83.0 | fL | 83.0 - 101.0 |
| MEAN CORP HB (MCH) | 27.2 | pg | 27.0 - 32.0 |
| MEAN CORP HB CONC (MCHC) | 32.6 | g/dL | 31.5 - 34.5 |
| PLATELET COUNT | 323 | $\times 10^3/\mu\text{L}$ | 150 - 410 |
| RDW-CV | 12.9 | % | 11.6 - 14.0 |

RAVIMEENA

Technologist

Page No: 1 of 16

DR.TANU RUNGTA

MD (Pathology)

RMC No. 17226



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



| | | | |
|-------------------------------------|--------------------------------|--------------------|----------|
| NAME :- Mrs. INDRA CHAUDHARY | Patient ID :-1223409 | Date :- 03/06/2023 | 09:39:25 |
| Age :- 36 Yrs 11 Mon 24 Days | Ref. By Doctor:-BANK OF BARODA | | |
| Sex :- Female | Lab/Hosp :- | | |
| | Company :- Mr.MEDIWHEEL | | |

Final Authentication : 03/06/2023 17:14:20

HAEMATOLOGY

Erythrocyte Sedimentation Rate (ESR)

08

mm in 1st hr

00 - 20

Method:- Westergreen

The erythrocyte sedimentation rate (ESR or sed rate) is a relatively simple, inexpensive, non-specific test that has been used for many years to help detect inflammation associated with conditions such as infections, cancers, and autoimmune diseases. ESR is said to be a non-specific test because an elevated result often indicates the presence of inflammation but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other tests, such as C-reactive protein. ESR is used to help diagnose certain specific inflammatory diseases, including temporal arteritis, systemic vasculitis and polymyalgia rheumatica. (For more on these, read the article on Vasculitis.) A significantly elevated ESR is one of the main test results used to support the diagnosis. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as



RAVIMEENA

Technologist

Page No: 2 of 16

DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



| | | | |
|-------------------------------------|--------------------------------|--------------------|----------|
| NAME :- Mrs. INDRA CHAUDHARY | Patient ID :-1223409 | Date :- 03/06/2023 | 09:39:25 |
| Age :- 36 Yrs 11 Mon 24 Days | Ref. By Doctor:-BANK OF BARODA | | |
| Sex :- Female | Lab/Hosp :- | | |
| | Company :- Mr.MEDIWHEEL | | |

(CBC): Methodology: TLC,DLC Fluorescent Flow cytometry, HB SLS method,TRBC,PCV,PLT Hydrodynamically focused Impedance. and MCH,MCV,MCHC,MENZER INDEX are calculated. InstrumentName: Sysmex 6 part fully automatic analyzer XN-L,Japan



RAVIMEENA

Page No: 3 of 16



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



| | | | |
|-------------------------------------|--------------------------------|--------------------|----------|
| NAME :- Mrs. INDRA CHAUDHARY | Patient ID :-1223409 | Date :- 03/06/2023 | 09:39:25 |
| Age :- 36 Yrs 11 Mon 24 Days | Ref. By Doctor:-BANK OF BARODA | | |
| Sex :- Female | Lab/Hosp :- | | |
| | Company :- | Mr.MEDIWHEEL | |

Final Authentication : 03/06/2023 17:14:20


BIOCHEMISTRY

| Test Name | Value | Unit | Biological Ref Interval |
|--|-----------------|-------|-------------------------|
| FASTING BLOOD SUGAR (Plasma) Method:- GOD POD | 112.0 | mg/dl | 70.0 - 115.0 |
| Impaired glucose tolerance (IGT) | 111 - 125 mg/dL | | |
| Diabetes Mellitus (DM) | > 126 mg/dL | | |

Instrument Name: HORIBA CA60 Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels(hypoglycemia) may result from excessive insulin therapy or various liver diseases .

BI. OOD SUGAR PP (Plasma) 122.0 mg/dl 70.0 - 140.0
Method - GOD PAP

Instrument Name: HORIBA Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels(hypoglycemia) may result from excessive insulin therapy or various liver diseases .



RAVIMEENA

Technologist
Page No: 4 of 16

DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics@gmail.com



| | | | |
|-------------------------------------|--------------------------------|--------------------|----------|
| NAME :- Mrs. INDRA CHAUDHARY | Patient ID :-1223409 | Date :- 03/06/2023 | 09:39:25 |
| Age :- 36 Yrs 11 Mon 24 Days | Ref. By Doctor:-BANK OF BARODA | | |
| Sex :- Female | Lab/Hosp :- | | |
| | Company :- Mr.MEDIWHEEL | | |

Final Authentication : 03/06/2023 17:14:20

HAEMATOLOGY

| Test Name | Value | Unit | Biological Ref Interval |
|--|-------|-------|--|
| GLYCOSYLATED HEMOGLOBIN (HbA1C) Method:- CAPILLARY with EDTA | 5.6 | mg% | Non-Diabetic < 6.0 Good Control 6.0-7.0 Weak Control 7.0-8.0 Poor control > 8.0 |
| MEAN PLASMA GLUCOSE Method:- Calculated Parameter | 110 | mg/dL | 68 - 125 |

INTERPRETATION

AS PER AMERICAN DIABETES ASSOCIATION (ADA)

Reference Group HbA1c in %

Non diabetic adults >=18 years < 5.7

At risk (Prediabetes) 5.7 - 6.4

Diagnosing Diabetes >= 6.5

CLINICAL NOTES

In vitro quantitative determination of HbA1c in whole blood is utilized in long term monitoring of glycemia. The HbA1c level correlates with the mean glucose concentration prevailing in the course of the patient's recent history (approx. 6-8 weeks) and therefore provides much more reliable information for glycemia monitoring than do determinations of blood glucose or urinary glucose. It is recommended that the determination of HbA1c be performed at intervals of 4-6 weeks during Diabetes Mellitus therapy. Results of HbA1c should be assessed in conjunction with the patient's medical history, clinical examinations and other findings. Some of the factors that influence HbA1c and its measurement [Adapted from Gallagher et al.]

1. Erythropoiesis

- Increased HbA1c: iron, vitamin B12 deficiency, decreased erythropoiesis
- Decreased HbA1c: administration of erythropoietin, iron, vitamin B12, reticulocytosis, chronic liver disease

2. Altered Haemoglobin-Genetic or chemical alterations in hemoglobin: hemoglobinopathies, HbF, methemoglobin, may increase or decrease HbA1c

3. Glycation

- Increased HbA1c: alcoholism, chronic renal failure, decreased intraerythrocytic pH
- Decreased HbA1c: certain hemoglobinopathies, increased intra-erythrocyte pH

4. Erythrocyte destruction

- Increased HbA1c: increased erythrocyte life span: Splenectomy
- Decreased A1c: decreased RBC life span: hemoglobinopathies, splenomegaly, rheumatoid arthritis or drugs such as antiretrovirals, ribavirin & dapsone.

5. Others

- Increased HbA1c: hyperbilirubinemia, carbamylated hemoglobin, alcoholism, large doses of aspirin, chronic opiate use, chronic renal failure
- Decreased HbA1c: hypertriglyceridemia, reticulocytosis, chronic liver disease, aspirin, vitamin C and E, splenomegaly, rheumatoid arthritis or drugs

Note

- Shortened RBC life span -HbA1c test will not be accurate when a person has a condition that affects the average lifespan of red blood cells (RBCs), such as hemolytic anemia or blood loss. When the lifespan of RBCs in circulation is shortened, the A1c result is falsely low and is an unreliable measurement of a person's average glucose over time.
- Abnormal forms of hemoglobin - The presence of some hemoglobin variants, such as hemoglobin S in sickle cell anemia, may affect certain methods for measuring A1c. In these cases, fructosamine can be used to monitor glucose control.

Advised:

- To follow patient for glycemic control test like fructosamine or glycated albumin may be performed instead
 - Hemoglobin HPLC screen to analyze abnormal hemoglobin variant.
- estimated Average Glucose (eAG) based on value calculated according to National Glycohemoglobin Standardization Program (NGSP) criteria.

RAVIMEENA

Technologist

Page No: 5 of 16

Tanu

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mrs. INDRA CHAUDHARY

Age :- 36 Yrs 11 Mon 24 Days

Sex :- Female

Patient ID :-1223409

Date :- 03/06/2023

09:39:25

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 03/06/2023 17:14:20

HAEMATOLOGY

BLOOD GROUP ABO

Method:- Haemagglutination reaction

"O" POSITIVE



RAVIMEENA

Technologist

Page No: 6 of 16

DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



| | | | |
|-------------------------------------|--------------------------------|--------------------|----------|
| NAME :- Mrs. INDRA CHAUDHARY | Patient ID :-1223409 | Date :- 03/06/2023 | 09:39:25 |
| Age :- 36 Yrs 11 Mon 24 Days | Ref. By Doctor:-BANK OF BARODA | | |
| Sex :- Female | Lab/Hosp :- | | |
| | Company :- | Mr.MEDIWHEEL | |

Final Authentication : 03/06/2023 17:14:20

BIOCHEMISTRY

| Test Name | Value | Unit | Biological Ref Interval |
|-----------|-------|------|-------------------------|
|-----------|-------|------|-------------------------|

LIPID PROFILE

TOTAL CHOLESTEROL 148.00 mg/dl
Desirable <200
Borderline 200-239
High > 240
Method:- CHOD-PAP methodology

InstrumentName MISPA PLUS **Interpretation:** Cholesterol measurements are used in the diagnosis and treatments of lipid lipoprotein metabolism disorders

TRIGLYCERIDES 99.80 mg/dl
Normal <150
Borderline high 150-199
High 200-499
Very high >500
Method:- GPO-PAP

InstrumentName Randox Rx Imola **Interpretation :** Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction.

DIRECT HDL CHOLESTEROL 37.00 mg/dl
MALE- 30-70
FEMALE - 30-85
Method:- Direct clearance Method

Instrument Name Rx Daytona plus **Interpretation:** An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.

LDL CHOLESTEROL 94.37 mg/dl
Optimal <100
Near Optimal/above optimal 100-129
Borderline High 130-159
High 160-189
Very High > 190
Method - Calculated Method

VI.DL CHOLESTEROL 19.96 mg/dl
0.00 - 80.00
Method- Calculated

T.CHOLESTEROL/HDL CHOLESTEROL RATIO 4.00
0.00 - 4.90
Method - Calculated

LDL / HDL CHOLESTEROL RATIO 2.55
0.00 - 3.50
Method - Calculated

TOTAL LIPID 453.20 mg/dl
400.00 - 1000.00
Method - CALCULATED

1. Measurements in the same patient can show physiological& analytical variations. Three serialsamples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL& LDL Cholesterol.

2. As per NCEP guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended

3. Low HDL levels are associated with Coronary Heart Disease due to insufficient HDL being available to participate in reverse cholesterol transport.

RAVIMEENA

Tanu Rungta

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226

Technologist
Page No: 7 of 16



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mrs. INDRA CHAUDHARY

Age :- 36 Yrs 11 Mon 24 Days

Sex :- Female

Patient ID :-1223409

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Date :- 03/06/2023 09:39:25

Final Authentication : 03/06/2023 17:14:20

BIOCHEMISTRY

transport, the process by which cholesterol is eliminated from peripheral tissues.

Comments: 1- ATP III suggested the addition of Non HDL Cholesterol (Total Cholesterol – HDL Cholesterol) as an indicator of all atherogenic lipoproteins (mainly LDL & VLDL). The Non HDL Cholesterol is used as a secondary target of therapy in persons with triglycerides ≥ 200 mg/dL. The goal for Non HDL Cholesterol in those with increased triglyceride is 30 mg/dL above that set for LDL Cholesterol.
2 -For calculation of CHD risk, history of smoking, any medication for hypertension & current B.P. levels are required.



RAVIMEENA

Technologist

Page No: 8 of 16

DR.TANU RUNGTA

MD (Pathology)

RMC No. 17226



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



| | | | |
|-------------------------------------|--------------------------------|--------------------|----------|
| NAME :- Mrs. INDRA CHAUDHARY | Patient ID :-1223409 | Date :- 03/06/2023 | 09:39:25 |
| Age :- 36 Yrs 11 Mon 24 Days | Ref. By Doctor:-BANK OF BARODA | | |
| Sex :- Female | Lab/Hosp :- | | |
| | Company :- Mr.MEDIWHEEL | | |

Final Authentication : 03/06/2023 17:14:20

BIOCHEMISTRY

LIVER PROFILE WITH GGT

| | | | |
|---|-------|-------|--|
| SERUM BILIRUBIN (TOTAL) Method- DMSO/Diazo | 0.60 | mg/dL | Infants : 0.2-8.0 mg/dL Adult - Up to - 1.2 mg/dL |
| SERUM BILIRUBIN (DIRECT) Method - DMSO/Diazo | 0.19 | mg/dL | Up to 0.40 mg/dL |
| SERUM BILIRUBIN (INDIRECT) Method - Calculated | 0.41 | mg/dl | 0.30-0.70 |
| SGOT Method- IFCC | 24.2 | U/L | 0.0 - 40.0 |
| SGPT Method- IFCC | 14.1 | U/L | 0.0 - 35.0 |
| SERUM ALKALINE PHOSPHATASE Method- DGKC - SCE | 69.10 | U/L | 64.00 - 306.00 |
| InstrumentName MISPA PLUS Interpretation :Measurements of alkaline phosphatase are of use in the diagnosis, treatment and investigation of hepatobiliary disease and in bone disease associated with increased osteoblastic activity. Alkaline phosphatase is also used in the diagnosis of parathyroid and intestinal disease | | | |
| SERUM GAMMA GT Method- Szasz methodology Instrument Name Randox Rx Imola Interpretation Elevations in GGT levels are seen earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or post-hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal) are observed with infectious hepatitis. | 16.50 | U/L | 5.00 - 32.00 |
| SERUM TOTAL PROTEIN Method- Direct Biuret Reagent | 6.94 | g/dl | 6.00 - 8.40 |
| SERUM ALBUMIN Method - Bromocresol Green | 4.74 | g/dl | 3.50 - 5.50 |
| SERUM GLOBULIN Method - CALCULATION | 2.20 | gm/dl | 2.20 - 3.50 |
| A/G RATIO | 2.15 | | 1.30 - 2.50 |

Interpretation : Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

Note :- These are group of tests that can be used to detect the presence of liver disease, distinguish among different types of liver disorders, gauge the extent of known liver damage, and monitor the response to treatment. Most liver diseases cause only mild symptoms initially, but these diseases must be detected early. Some tests are associated with functionality (e.g., albumin), some with cellular integrity (e.g., transaminase), and some with conditions linked to the biliary tract (gamma-glutamyl transferase and alkaline phosphatase). Conditions with elevated levels of ALT and AST include hepatitis A,B ,C ,paracetamol toxicity etc. Several biochemical tests are useful in the evaluation and management of patients with hepatic dysfunction. Some or all of these measurements are also carried out (usually about twice a year for routine cases) on those individuals taking certain medications, such as anticonvulsants, to ensure that the medications are not adversely impacting the person's liver.

RAVIMEENA

Technologist
Page No: 9 of 16

DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



| | | | |
|-------------------------------------|--------------------------------|--------------------|----------|
| NAME :- Mrs. INDRA CHAUDHARY | Patient ID :-1223409 | Date :- 03/06/2023 | 09:39:25 |
| Age :- 36 Yrs 11 Mon 24 Days | Ref. By Doctor:-BANK OF BARODA | | |
| Sex :- Female | Lab/Hosp :- | | |
| | Company :- Mr.MEDIWHEEL | | |

Final Authentication : 03/06/2023 17:14:20

BIOCHEMISTRY

RFT / KFT WITH ELECTROLYTES

SERUM UREA 27.60 mg/dl 10.00 - 50.00
Method:- Urease/GLDH

InstrumentName: HORIBA CA 60 **Interpretation :** Urea measurements are used in the diagnosis and treatment of certain renal and metabolic diseases.

SERUM CREATININE 0.80 mg/dl Males : 0.6-1.50 mg/dl
Method - Jaffe's Method Females : 0.6 -1.40 mg/dl

Interpretation :
Creatinine is measured primarily to assess kidney function and has certain advantages over the measurement of urea. The plasma level of creatinine is relatively independent of protein ingestion, water intake, rate of urine production and exercise. Depressed levels of plasma creatinine are rare and not clinically significant.

SERUM URIC ACID 4.94 mg/dl 2.40 - 7.00

InstrumentName: HORIBA YUMIZEN CA60 Daytona plus **Interpretation: Elevated Urate:** High purine diet, Alcohol, Renal insufficiency, Drugs, Polycythaemia vera, Malignancies, Hypothyroidism, Rare enzyme defects, Down's syndrome, Metabolic syndrome, Pregnancy, Gout.

SODIUM 140.7 mmol/L 135.0 - 150.0
Method - ISE

Interpretation: Decreased sodium - Hyponatraemia Causes include: fluid or electrolyte loss, Drugs, Oedematous states, Legionnaire's disease and other chest infections, pseudonatremia, Hyperlipidaemias and paraproteinaemias, endocrine diseases, SIADH.

POTASSIUM 4.51 mmol/L 3.50 - 5.50
Method:- ISE

Interpretation: A. Elevated potassium (hyperkalaemia) • Artefactual, Physiologic elevation, Drugs, Pathological states, Renal failure Adrenocortical insufficiency, metabolic acidoses, very high platelet or white cell counts B. Decreased potassium (hypokalaemia) Drugs, Liqueuric, Diarrhoea and vomiting, Metabolic alkalosis, Corticosteroid excess, Oedematous state, Anorexia nervosa/bulimia

CHLORIDE 97.8 mmol/L 94.0 - 110.0
Method - ISF

Interpretation: Used for Electrolyte monitoring.

SERUM CALCIUM 9.23 mg/dL 8.80 - 10.20
Method:- Arsenazo III Method

InstrumentName: MISPA PLUS **Interpretation:** Serum calcium levels are believed to be controlled by parathyroid hormone and vitamin D. Increases in serum PTH or vitamin D are usually associated with hypercalcemia. Hypocalcemia may be observed in hypoparathyroidism, nephrosis and pancreatitis.

SERUM TOTAL PROTEIN 6.94 g/dl 6.00 - 8.40
Method:- Buret Reagent

Technologist
Page No: 10 of 16

Tanu
DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



| | | | |
|-------------------------------------|--------------------------------|--------------------|----------|
| NAME :- Mrs. INDRA CHAUDHARY | Patient ID :-1223409 | Date :- 03/06/2023 | 09:39:25 |
| Age :- 36 Yrs 11 Mon 24 Days | Ref. By Doctor:-BANK OF BARODA | | |
| Sex :- Female | Lab/Hosp :- | | |
| | Company :- Mr.MEDIWHEEL | | |

Final Authentication : 03/06/2023 17:14:20

BIOCHEMISTRY


| | | | |
|--|------|-------|-------------|
| SERUM ALBUMIN Method- Bromocresol Green | 4.74 | g/dl | 3.50 - 5.50 |
| SERUM GLOBULIN Method- CALCULATION | 2.20 | gm/dl | 2.20 - 3.50 |
| A/G RATIO | 2.15 | | 1.30 - 2.50 |

Interpretation : Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

INTERPRETATION

Kidney function tests are group of tests that can be used to evaluate how well the kidneys are functioning. Creatinine is a waste product that comes from protein in the diet and also comes from the normal wear and tear of muscles of the body. In blood, it is a marker of GFR .in urine, it can remove the need for 24-hourcollections for many analytes or be used as a quality assurance tool to assess the accuracy of a 24-hour collection Higher levels may be a sign that the kidneys are not working properly. As kidney disease progresses, the level of creatinine and urea in the bloodincreases. Certain drugs are nephrotoxic hence KFT is done before and after initiation of treatment with these drugs.

Low serum creatinine values are rare; they almost always reflect low muscle mass.



RAVIMEENA

Technologist

Page No: 11 of 16

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



| | | | |
|-------------------------------------|--------------------------------|--------------------|----------|
| NAME :- Mrs. INDRA CHAUDHARY | Patient ID :-1223409 | Date :- 03/06/2023 | 09:39:25 |
| Age :- 36 Yrs 11 Mon 24 Days | Ref. By Doctor:-BANK OF BARODA | | |
| Sex :- Female | Lab/Hosp :- | | |
| | Company :- Mr.MEDIWHEEL | | |

Final Authentication : 03/06/2023 17:14:20

CLINICAL PATHOLOGY

| Test Name | Value | Unit | Biological Ref Interval |
|--------------------------------------|-------------|------|-------------------------|
| Urine Routine | | | |
| <u>PHYSICAL EXAMINATION</u> | | | |
| COLOUR | PALE YELLOW | | PALE YELLOW |
| APPEARANCE | Clear | | Clear |
| <u>CHEMICAL EXAMINATION</u> | | | |
| REACTION(PH) | 7.0 | | 5.0 - 7.5 |
| SPECIFIC GRAVITY | 1.010 | | 1.010 - 1.030 |
| PROTEIN | NIL | | NIL |
| SUGAR | NIL | | NIL |
| BILIRUBIN | NEGATIVE | | NEGATIVE |
| UROBILINOGEN | NORMAL | | NORMAL |
| KETONES | NEGATIVE | | NEGATIVE |
| NITRITE | NEGATIVE | | NEGATIVE |
| <u>MICROSCOPY EXAMINATION</u> | | | |
| RBC/HPF | NIL | /HPF | NIL |
| WBC/HPF | 2-3 | /HPF | 2-3 |
| EPITHELIAL CELLS | 2-3 | /HPF | 2-3 |
| CRYSTALS/HPF | ABSENT | | ABSENT |
| CAST/HPF | ABSENT | | ABSENT |
| AMORPHOUS SEDIMENT | ABSENT | | ABSENT |
| BACTERIAL FLORA | ABSENT | | ABSENT |
| YEAST CELL | ABSENT | | ABSENT |
| OTHER | ABSENT | | ABSENT |

RAVIMEENA

Technologist
Page No: 12 of 16

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mrs. INDRA CHAUDHARY

Age :- 36 Yrs 11 Mon 24 Days

Sex :- Female

Patient ID :-1223409

Date :- 03/06/2023

09:39:25

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 03/06/2023 17:14:20

CLINICAL PATHOLOGY

URINE SUGAR (FASTING)
Collected Sample Received

Nil

Nil



RAVIMEENA

Technologist

Page No: 13 of 16

DR. TANU RUNGTA

MD (Pathology)

RMC No. 17226



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



| | | | |
|-------------------------------------|--------------------------------|--------------------|----------|
| NAME :- Mrs. INDRA CHAUDHARY | Patient ID :-1223409 | Date :- 03/06/2023 | 09:39:25 |
| Age :- 36 Yrs 11 Mon 24 Days | Ref. By Doctor:-BANK OF BARODA | | |
| Sex :- Female | Lab/Hosp :- | | |
| | Company :- Mr.MEDIWHEEL | | |

Final Authentication : 03/06/2023 17:14:20

TOTAL THYROID PROFILE

IMMUNOASSAY

| Test Name | Value | Unit | Biological Ref Interval |
|-----------|-------|------|-------------------------|
|-----------|-------|------|-------------------------|

THYROID-TRIIODOTHYRONINE T3
Method- ECLIA

0.95 ng/ml.

0.70 - 2.04

NOTE-TSH levels are subject to circadian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result. Transient increase in TSH levels or abnormal TSH levels can be seen in some non thyroidal conditions, simultaneous measurement of TSH with free T4 is useful in evaluating differential diagnosis

INTERPRETATION-Ultra Sensitive 4th generation assay 1.Primary hyperthyroidism is accompanied by ↑serum T3 & T4 values along with ↓ TSH level. 2.Low TSH, high FT4 and TSH receptor antibody (TRAb) +ve seen in patients with Graves disease 3.Low TSH, high FT4 and TSH receptor antibody (TRAb) -ve seen in patients with Toxic adenoma/Toxic Multinodular goiter 4.High TSH, Low FT4 and Thyroid microsomal antibody increased seen in patients with Hashimotos thyroiditis 5.High TSH, Low FT4 and Thyroid microsomal antibody normal seen in patients with Iodine deficiency/Congenital T4 synthesis deficiency 6.Low TSH, Low FT4 and TRH stimulation test -Delayed response seen in patients with Tertiary hypothyroidism 7.Primary hypothyroidism is accompanied by ↓ serum T3 and T4 values & ↑serum TSH levels 8.Normal T4 levels accompanied by ↑ T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis 9.Normal cr T3 & T4 10.Normal T3 & T4 along with ↑ TSH indicate mild / Subclinical Hyperthyroidism . 11.Normal T3 & T4 along with ↓ TSH is seen in Hypothyroidism . 12.Normal T3 & T4 levels with ↑ TSH indicate Mild / Subclinical Hypoth

DURING PREGNANCY - REFERENCE RANGE for TSH IN uIU/mL (As per American Thyroid Association) 1st Trimester : 0.10-2.50 uIU/mL 2nd Trimester : 0.20-3.00 uIU/mL 3rd Trimester : 0.30-3.00 uIU/mL. The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy.

REMARK-Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test. Abnormal thyroid test findings often found in critically ill patients should be repeated after the critical nature of the condition is resolved. TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age, and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly. ***

THYROID THYRONINE (T4)
Method- ECLIA

6.15 uIU/mL

5.10 - 14.10

NOTE-TSH levels are subject to circadian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result. Transient increase in TSH levels or abnormal TSH levels can be seen in some non thyroidal conditions, simultaneous measurement of TSH with free T4 is useful in evaluating differential diagnosis

INTERPRETATION-Ultra Sensitive 4th generation assay 1.Primary hyperthyroidism is accompanied by ↑serum T3 & T4 values along with ↓ TSH level. 2.Low TSH, high FT4 and TSH receptor antibody (TRAb) +ve seen in patients with Graves disease 3.Low TSH, high FT4 and TSH receptor antibody (TRAb) -ve seen in patients with Toxic adenoma/Toxic Multinodular goiter 4.High TSH, Low FT4 and Thyroid microsomal antibody increased seen in patients with Hashimotos thyroiditis 5.High TSH, Low FT4 and Thyroid microsomal antibody normal seen in patients with Iodine deficiency/Congenital T4 synthesis deficiency 6.Low TSH, Low FT4 and TRH stimulation test -Delayed response seen in patients with Tertiary hypothyroidism 7.Primary hypothyroidism is accompanied by ↓ serum T3 and T4 values & ↑serum TSH levels 8.Normal T4 levels accompanied by ↑ T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis 9.Normal cr T3 & T4 10.Normal T3 & T4 along with ↑ TSH indicate mild / Subclinical Hyperthyroidism . 11.Normal T3 & T4 along with ↓ TSH is seen in Hypothyroidism . 12.Normal T3 & T4 levels with ↑ TSH indicate Mild / Subclinical Hypoth

DURING PREGNANCY - REFERENCE RANGE for TSH IN uIU/mL (As per American Thyroid Association) 1st Trimester : 0.10-2.50 uIU/mL 2nd Trimester : 0.20-3.00 uIU/mL 3rd Trimester : 0.30-3.00 uIU/mL. The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy.

REMARK-Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test. Abnormal thyroid test findings often found in critically ill patients should be repeated after the critical nature of the condition is resolved. TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age, and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly.

TSH
Method- ECLIA

1.224 uIU/mL

0.350 - 5.500

NOTE-TSH levels are subject to circadian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result. Transient increase in TSH levels or abnormal TSH levels can be seen in some non thyroidal conditions, simultaneous measurement of TSH with free T4 is useful in evaluating differential diagnosis

INTERPRETATION-Ultra Sensitive 4th generation assay 1.Primary hyperthyroidism is accompanied by ↑serum T3 & T4 values along with ↓ TSH level.

RAVIMEENA

Technologist

Page No: 15 of 16

Tanu

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



📍 B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 ✉ maxcarediagnostics1@gmail.com



| | | | |
|-------------------------------------|--------------------------------|--------------------|----------|
| NAME :- Mrs. INDRA CHAUDHARY | Patient ID :-1223409 | Date :- 03/06/2023 | 09:39:25 |
| Age :- 36 Yrs 11 Mon 24 Days | Ref. By Doctor:-BANK OF BARODA | | |
| Sex :- Female | Lab/Hosp :- | | |
| | Company :- Mr.MEDIWHEEL | | |

Final Authentication : 03/06/2023 17:14:20

IMMUNOASSAY


2. Low TSH,high FT4 and TSH receptor antibody(TRAb) +ve seen in patients with Graves disease
3. Low TSH,high FT4 and TSH receptor antibody(TRAb) -ve seen in patients with Toxic adenoma/Toxic Multinodular goiter
4. HighTSH,Low FT4 and Thyroid microsomal antibody increased seen in patients with Hashimoto's thyroiditis
5. HighTSH,Low FT4 and Thyroid microsomal antibody normal seen in patients with Iodine deficiency/Congenital T4 synthesis deficiency
6. Low TSH,Low FT4 and TRH stimulation test -Delayed response seen in patients with Tertiary hypothyroidism
7. Primary hypothyroidism is accompanied by ↓ serum T3 and T4 values & ↑serum TSH levels
8. Normal T4 levels accompanied by ↑ T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis
9. Normal or, T3 & ↑T4 levels indicate T4 Thyrotoxicosis (problem is conversion of T4 to T3)
10. Normal T3 & T4 along with ↓ TSH indicate mild / Subclinical Hyperthyroidism .
11. Normal T3 & ↓ T4 along with ↑ TSH is seen in Hypothyroidism
12. Normal T3 & T4 levels with ↑ TSH indicate Mild / Subclinical Hypothyroidism
13. Slightly ↑ T3 levels may be found in pregnancy and in estrogen therapy while ↓ levels may be encountered in severe illness , malnutrition , renal failure and during therapy with drugs like propranolol
14. Although ↑ TSH levels are nearly always indicative of Primary Hypothyroidism ,rarely they can result from TSH secreting pituitary tumours.

DURING PREGNANCY - REFERENCE RANGE for TSH IN uIU/mL (As per American Thyroid Association)

- 1st Trimester : 0.10-2.50 uIU/mL
- 2nd Trimester : 0.20-3.00 uIU/mL
- 3rd Trimester : 0.30-3.00 uIU/mL

The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy.

REMARK-Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test. Abnormal thyroid test findings often found in critically ill patients should be repeated after the critical nature of the condition is resolved. TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age, and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly.



RAVIMEENA

Technologist

Page No: 16 of 16

Tanu

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



| | | | |
|--------|----------------------|---------|------------|
| NAME: | MRS. INDRA CHOUDHARY | AGE/SEX | 36 YRS/F |
| REF.BY | BANK OF BARODA | DATE | 03/06/2023 |

CHEST X RAY (PA VIEW)

Bilateral lung fields appear clear.

Bilateral costo-phrenic angles appear clear.

Cardiothoracic ratio is normal.

Thoracic soft tissue and skeletal system appear unremarkable.

Soft tissue shadows appear normal.

IMPRESSION: No significant abnormality is detected.

DR.SHALINI GOEL
M.B.B.S, D.N.B (Radiodiagnosis)
RMC No.: 21954







| | |
|--------------------------------------|--------------------------------|
| MRS. INDRA CHOUDHARY | Age: 36 Y/F |
| Registration Date: 03/06/2023 | Ref. by: BANK OF BARODA |

ULTRASOUND OF WHOLE ABDOMEN

Liver is of normal size (11.5 cm). Echo-texture is normal. No focal space occupying lesion is seen within liver parenchyma. Intrahepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is well distended. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape (9.8 cm). Echotexture is normal. No focal lesion is seen.

Kidneys are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. No focal lesion is seen. Collecting system does not show any dilatation or calculus.

Right kidney is measuring approx. 10.7 x 4.0 cm.

Left kidney is measuring approx. 11.8 x 4.4 cm.

Urinary bladder does not show any calculus or mass lesion.

Uterus is retroverted and normal in size (measuring approx. 7.3 x 4.0 x 3.8 cm).

Myometrium shows normal echo -pattern. No focal space occupying lesion is seen. Endometrial echo is normal. Endometrial thickness is 6.4 mm.

A well-defined, thick-walled, avascular, cystic lesion of size 33 x 28 x 34 mm (AP x TR x CC) is noted in left ovary with internal septations giving fishnet appearance – suggestive of hemorrhagic ovarian cyst.
Right ovary is normal.

No enlarged nodes are visualized. No retro-peritoneal lesion is identified.

No significant free fluid is seen in pouch of Douglas.

NOTE: No evidence of probe tenderness is noted in RIF

IMPRESSION: Left ovarian hemorrhagic cyst as described above.

DR.SHALINI GOEL
M.B.B.S, D.N.B (Radiodiagnosis)
RMC no.: 21954


Dr. SHALINI GOEL
MBBS, DNB (Radiologist)
RMC No. 21954
P-3 Health Solutions LLP

P3 HEALTH SOLUTIONS LLP

B-14, V/dhyanager Nagar, Enclave, Phase-2, Jaipur
12229451323779/Mrs Indra Chaudhary 36Yrs/Female Kgs/31 Cms BP: ___/___ mmHg
Ref.: BANK OF BARODA Test Date: 03-Jun-2023(10:47:13) Notch: 50Hz 0.05Hz - 100Hz 10mm/mV 25mm/Sec

HR: 54 bpm

PR Interval: 154 ms
QRS Duration: 108 ms
QT/QTc: 426/404ms
P-QRS-T Axis: 19 - -1 - 21 (Deg)



FINDINGS: Abnormal ECG with indication of Sinus Bradycardia
Vent Rate : 54 bpm; PR Interval : 154 ms; QRS Duration : 108 ms; QT/QTc Int : 426/404 ms
P-QRS-T axis: 19 - -1 - 21 (Deg)
Comments :

Borderline sinus bradycardia

SM

Dr. Naresh Kumar Mohanka
RMC No.: 35703
MBBS, DIP. CARDIO (ESCORTS)
D.E.M. (RCGP-UK)

B-14, Vidhyadhar Nagar Enclave, Phase -2, Jaipur

1323258/MRS INDRA CHAUDHARY

36 Yrs/Female 0 Kg/0 Cms

Date: 03-Jun-2023 10:51:39 AM

Ref. By : BANK OF BARODA

Protocol : BRUCE

Medication :
Objective :


History :

| Stage | StageTime (Min:Sec) | PhaseTime (Min:Sec) | Speed (mph) | Grade (%) | METS | H.R. (bpm) | B.P. (mmHg) | R.P.P. x100 | PVC | Comments |
|----------|---------------------|---------------------|-------------|-----------|------|------------|-------------|-------------|-----|----------|
| Supine | | | | | 1.0 | 55 | 120/80 | 66 | - | |
| Standing | | | | | 1.0 | 74 | 120/80 | 88 | - | |
| HV | | | | | 1.0 | 67 | 120/80 | 80 | - | |
| ExStart | | | | | 1.0 | 77 | 120/80 | 92 | - | |
| Stage 1 | 3:01 | 3:02 | 1.7 | 10.0 | 4.7 | 102 | 130/80 | 132 | - | |
| Stage 2 | 3:01 | 6:02 | 2.5 | 12.0 | 7.1 | 129 | 140/80 | 180 | - | |
| Stage 3 | 3:01 | 9:02 | 3.4 | 14.0 | 10.2 | 160 | 150/85 | 240 | - | |
| PeakEx | 0:06 | 9:07 | 4.2 | 16.0 | 10.3 | 160 | 150/85 | 240 | - | |
| Recovery | 1:00 | | 0.0 | 0.0 | 4.3 | 106 | 150/85 | 158 | - | |
| Recovery | 2:00 | | 0.0 | 0.0 | 1.0 | 80 | 160/90 | 128 | - | |
| Recovery | 3:00 | | 0.0 | 0.0 | 1.0 | 74 | 150/85 | 111 | - | |
| Recovery | 4:00 | | 0.0 | 0.0 | 1.0 | 74 | 140/85 | 103 | - | |

Findings :

Exercise Time : 09:06
 Max HR Attained : 160 bpm 87% of Max Predictable HR 184
 Max BP : 160/90(mmHg)
 Max Workload attained : 10.3 (Good Effort Tolerance)

T.M. IS NEARLY NORMAL



Advice/Comments:

Stm

[Signature]

Dr. Naresh Kumar Mohanka
 RMC No.: 35703
 MBBS, DIP. CARDIO (ESCORTS)
 D.E.M. (RCGP-UK)




HR: 55 bpm
METS: 1.0
BP: 120/80

MPHR: 29% of 184
Speed: 0.0 mph
Grade: 0.0%

Raw ECG
BRUCE
(1.0-35)Hz

Ex Time 00:35
BLC : On
Notch : On

Supine
10.0 mm/mV
25 mm/Sec.



HR: 74 bpm
METS: 1.0
BP: 120/80

MPHR: 40% of 184
Speed: 0.0 mph
Grade: 0.0%




Raw ECG
BRUCE
(1.0-35)HZ

Ex Time 01:08
BLC : On
Notch : On

Standing
10.0 mm/mV
25 mm/Sec.



V5
0.1



B-14, Vidhyadhar Nagar Enclave, Phase -2, Jaipur

1323258/MRS INDRA CHAUDHARY

36 Yrs/Female

0 Kg/0 Cms

Date: 03-Jun-2023 10:51:39 AM
4X 107 ms Post J





HR: 67 bpm
METS: 1.0
BP: 120/80

MPHR: 36% of 184
Speed: 0.0 mph
Grade: 0.0%

Raw ECG
BRUCE
(1.0-35)HZ

Ex Time 01:24
BLC : On
Notch : On

HV
10.0 mm/mV
25 mm/Sec.



B-14, Vidhyadhar Nagar Enclave, Phase -2, Jaipur
1323258/MRS INDRA CHAUDHARY
36 Yrs/Female
0 Kg/0 Cms
Date: 03-Jun-2023 10:51:39 AM
4X 107ms Post J



HR: 76 bpm
METS: 1.0
BP: 120/80

MPHR: 41% of 184
Speed: 0.0 mph
Grade: 0.0%

Raw ECG
BRUCE
(1.0-35)Hz

Ex Time 02:03
BLC : On
Notch : On

ExStart
10.0 mm/mv
25 mm/Sec.



B-14, Vidhyadhar Nagar Enclave, Phase -2, Jaipur

1323258/MRS INDRA CHAUDHARY

36 Yrs/Female

0 Kg/0 Cms

Date: 03-Jun-2023 10:51:39 AM
4X 107 ms Post U



HR: 102 bpm
METS: 4.7
BP: 130/80

APHR: 55% of 184
Speed: 1.7 mph
Grade: 10.0%

Raw ECG
BRUCE
(1.0-35)Hz

EX Time 02:59
BLC :On
Notch :On

BRUCE: Stage 1(3:00)
10.0 mm/mV
25 mm/Sec.



1323258/MRS INDRA CHAUDHARY

03-Jun-2023 10:51:39 AM



HR: 129 bpm
METs: 7.1
BP: 140/80

MPHR: 70% of 184
Speed: 2.5 mph
Grade: 12.0%

Raw ECG
BRUCE
(1.0-35)HZ

Ex Time 05:59
BLC : On
Notch : On

BRUCE: Stage 2(3:00)
10.0 mm/mV
25 mm/Sec.





HR: 161 bpm
METs: 10.2
BP: 150/85

MPHR: 87% of 184
Speed: 3.4 mph
Grade: 14.0%

Raw ECG
BRUCE
(1.0-35)Hz

Ex Time 08:59
BLC : On
Notch : On

BRUCE: Stage 3(3:00)
10.0 mm/mV
25 mm/Sec.



B-14, Vidhyadhar Nagar Enclave, Phase -2, Jaipur

1323258/MRS INDRA CHAUDHARY

36 Yrs/Female

0 Kg/0 Cms

Date: 03-Jun-2023 10:51:39 AM
4X 107ms Post J




HR: 160 bpm
METS: 10.3
BP: 150/85

MPHR: 86% of 184
Speed: 4.2 mph
Grade: 16.0%

Raw ECG
BRUCE
(1.0-35)Hz

EX Time 09:04
BLC : On
Notch : On

BRUCE: PeakEx(0:04)
10.0 mm/mV
25 mm/Sec



B-14, Vidhyadhar Nagar Enclave, Phase -2, Jaipur

1323258/MRS INDRRA CHAUDHARY

36 Yrs./Female

0 Kg/0 Cms

Date: 03-Jun-2023 10:51:39 AM

4X 107ms Post-J

HR: 106 bpm

METS: 4.4

BP: 150/85

MPHR: 57% of 184

Speed: 0.0 mph

Grade: 0.0%

Raw ECG

BRUCE

(1.0-35)Hz

Ex Time 09:06



BLC :On

Notch :On

Recovery(1:00)

10.0 mm/mV

25 mm/Sec.



B-14, Vidhyadhar Nagar Enclave, Phase -2, Jaipur

1323258/MRS INDRA CHAUDHARY
36 Yrs/Female
0 Kg/0 Cms

Date: 03-Jun-2023 10:51:39 AM
4X 107 ms Post J

HR: 78 bpm
METS: 1.0
BP: 160/90

MPHR: 42% of 184
Speed: 0.0 mph
Grade: 0.0%





Raw ECG
BRUCE
(1.0-35)HZ

Ex Time 09:06
BLC : On
Notch : On

Recovery(2:00)
10.0 mm/mv
25 mm/Sec.



V5
0.2





HR: 74 bpm
METs: 1.0
BP: 150/85

MpHR: 40% of 184
Speed: 0.0 mph
Grade: 0.0%

Raw ECG
BRUCE
(1.0-3.5)Hz

Ex Time 09:06
BLC : On
Notch : On

Recovery(3:00)
10.0 mm/mV
25 mm/Sec





HR: 72 bpm
METs: 1.0
BP: 140/85

MpHR: 39% of 184
Speed: 0.0 mph
Grade: 0.0%

Raw ECG
BRUCE
(1.0-35)Hz

Ex Time 09:06
BLC :On
Notch :On

Recovery(4:00)
10.0 mm/mV
25 mm/Sec.



I II III aVR aVL aVF V1 V2 V3 V4 V5 V6

