



Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 GN: U85110UP2003PLC193493

Patient Name	: Mrs.BANDANA SINGH	Registered On	: 10/Nov/2024 09:24:10
Age/Gender	: 31 Y 0 M 0 D / F	Collected	: 10/Nov/2024 09:38:42
UHID/MR NO	: CALI.0000051596	Received	: 10/Nov/2024 11:34:58
Visit ID	: CALI0170162425	Reported	: 10/Nov/2024 14:31:45
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

DEPARTMENT OF HAEM ATOLOGY

MEDIWHEEL BANK OF BARODA FEM ALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
Blood Group (ABO & Rh typing) **	, Blood			
Blood Group	В			ERYTHROCYTE MAGNETIZED TECHNOLOGY/ TUBE AGGLUTINA
Rh (Anti-D)	POSITIVE			ERYTHROCYTE MAGNETIZED TECHNOLOGY/ TUBE AGGLUTINA
Complete Blood Count (CBC) ** , M	/hole Blood			
Haemoglobin	12.00	g/ dl	1 Day- 14.5-22.5 g/dl 1 Wk- 13.5-19.5 g/dl 1 Mo- 10.0-18.0 g/dl 3-6 Mo- 9.5-13.5 g/dl 0.5-2 Yr- 10.5-13.5 g/dl 2-6 Yr- 11.5-15.5 g/dl 6-12 Yr- 11.5-15.5 g/dl 12-18 Yr 13.0-16.0 g/dl Male- 13.5-17.5 g/dl Female- 12.0-15.5 g/dl	COLORIMETRICMETHOD (CYANIDE-FREE REAGENT)
TLC (WBC) DLC	7,800.00	/Qu mm	4000-10000	IMPEDANCE METHOD
Polymorphs (Neutrophils)	67.00	%	40-80	FLOW CYTOMETRY
Lymphocytes	27.00	%	20-40	FLOW CYTOMETRY
Monocytes	3.00	%	2-10	FLOW CYTOMETRY
Eosinophils	3.00	%	1-6	FLOW CYTOMETRY
Basophils ESR	0.00	%	<1-2	FLOW CYTOMETRY
Observed	16.00	MM/1H	10-19 Yr 8.0 20-29 Yr 10.8 30-39 Yr 10.4 40-49 Yr 13.6 50-59 Yr 14.2 60-69 Yr 16.0 70-79 Yr 16.5 80-91 Yr 15.8	









Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 GN: U85110UP2003PLC193493

Patient Name	: Mrs.BANDANA SINGH	Registered On	: 10/Nov/2024 09:24:10
Age/Gender	: 31 Y 0 M 0 D / F	Collected	: 10/Nov/2024 09:38:42
UHID/MR NO	: CALI.0000051596	Received	: 10/Nov/2024 11:34:58
Visit ID	: CALI0170162425	Reported	: 10/Nov/2024 14:31:45
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

DEPARTMENT OF HAEM ATOLOGY

MEDIWHEEL BANK OF BARODA FEM ALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
			Pregnancy	
			Early gestation - 48 (62	
			if anaemic) Leter gestation - 70 (95	
			if anaemic)	,
Corrected	10.00	Mm for 1st hr.	,	
PCV (HCT)	37.00	%	40-54	
Platelet count				
Platelet Count	1.80	LACS/ cu mm	1.5-4.0	ELECTRONIC
				IMPEDANCE/MICROSCOPIC
PDW (Platelet Distribution width)	15.70	fL	9-17	ELECTRONIC IMPEDANCE
P-LOR (Platelet Large Cell Patio)	50.50	%	35-60	ELECTRONIC IMPEDANCE
PCT (Platelet Hematocrit)	0.23	%	0.108-0.282	ELECTRONIC IM PEDANCE
MPV (Mean Platelet Volume)	13.20	fL	6.5-12.0	ELECTRONIC IM PEDANCE
RBCCount				
RBC Count	4.23	Mill./cumm	3.7-5.0	ELECTRONIC IMPEDANCE
Blood Indices (MCV, MCH, MCHC)				
MCV	88.70	fl	80-100	CALCULATED PARAMETER
MCH	28.30	pg	27-32	CALCULATED PARAMETER
МОНС	31.90	%	30-38	CALCULATED PARAMETER
RDW-CV	14.90	%	11-16	ELECTRONIC IM PEDANCE
RDW-SD	47.60	fL	35-60	ELECTRONIC IMPEDANCE
Absolute Neutrophils Count	5,226.00	/cu mm	3000-7000	
Absolute Eosinophils Count (AEC)	234.00	/cu mm	40-440	

Dr. Neetu Kushwaha MD.PATH

View Reports on









Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 QN: U85110UP2003PLC193493

Patient Name	: Mrs.BANDANA SINGH	Registered On	: 10/Nov/2024 09:24:11
Age/Gender	: 31 Y 0 M 0 D / F	Collected	: 10/Nov/2024 09:38:42
UHID/MR NO	: CALI.0000051596	Received	: 10/Nov/2024 11:41:22
Visit ID	: CALI0170162425	Reported	: 10/Nov/2024 13:19:59
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interva	al Method
GLUCOSE FASTING ** , Plasma Glucose Fasting	93.30	5	< 100 Normal 100-125 Pre-diabetes ≥ 126 Diabetes	GOD POD

Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions.b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential.c) I.G.T = Impaired Glucose Tolerance.

CLINICAL SIGNIFICANCE:- Glucose is the major source of energy in the body. Lack of insulin or resistance to it section at the cellular level causes diabetes. Therefore, the blood glucose levels are very high. Elevated serum glucose levels are observed in diabetes mellitus and may be associated with pancreatitis, pituitary or thyroid dysfunction and liver disease. Hypoglycaemia occurs most frequently due to over dosage of insulin.

Gucose PP**	110.20	mg/ dl	<140 Normal	GOD POD
Sample:Plasma After Meal			140-199 Pre-diabetes	
			>200 Diabetes	

Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions.b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential.c) I.G.T = Impaired Glucose Tolerance.

GLYCOSYLATED HAEMOGLOBIN (HBA1C) ** , EDTA BLOOD

Glycosylated Haemoglobin (HbA1c)	5.00	%NGSP
Glycosylated Haemoglobin (HbA1c)	31.00	mmol/mol/IFCC
Estimated Average Glucose (eAG)	96	mg/dl

Interpretation:

NOTE:-

• eAG is directly related to A1c.



HPLC(NGSP)



Page 3 of 12

View Reports on



Те



CHANDAN DIAGNOSTIC CENTRE

Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 QN: U85110UP2003PLC193493

Patient Name	: Mrs.BANDANA SINGH	Registered On	: 10/Nov/2024 09:24:11
Age/Gender	: 31 Y 0 M 0 D / F	Collected	: 10/Nov/2024 09:38:42
UHID/MR NO	: CALI.0000051596	Received	: 10/Nov/2024 11:41:22
Visit ID	: CALI0170162425	Reported	: 10/Nov/2024 13:19:59
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

est Name	Result	Unit	Bio. Ref. Interval	Method

- An A1c of 7% -the goal for most people with diabetes-is the equivalent of an eAG of 154 mg/dl.
- eAG may help facilitate a better understanding of actual daily control helping you and your health care provider to make necessary changes to your diet and physical activity to improve overall diabetes mnagement.

The following ranges may be used for interpretation of results. However, factors such as duration of diabetes, adherence to therapy and the age of the patient should also be considered in assessing the degree of blood glucose control.

Haemoglobin A1C (%)NGSP	mmol/mol / IFCC Unit	eAG (mg/dl)	Degree of Glucose Control Unit
> 8	>63.9	>183	Action Suggested*
7-8	53.0 -63.9	154-183	Fair Control
< 7	<63.9	<154	Goal**
6-7	42.1 -63.9	126-154	Near-normal glycemia
< 6%	<42.1	<126	Non-diabetic level

*High risk of developing long term complications such as Retinopathy, Nephropathy, Neuropathy, Cardiopathy, etc. **Some danger of hypoglycemic reaction in Type 1diabetics. Some glucose intolerant individuals and "subclinical" diabetics may demonstrate HbA1C levels in this area.

N.B.: Test carried out on Automated VARIANT II TURBO HPLC Analyser.

<u>Clinical Implications:</u>

*Values are frequently increased in persons with poorly controlled or newly diagnosed diabetes.

*With optimal control, the HbA 1c moves toward normal levels.

*A diabetic patient who recently comes under good control may still show higher concentrations of glycosylated hemoglobin. This level declines gradually over several months as nearly normal glycosylated *Increases in glycosylated hemoglobin occur in the following non-diabetic conditions: a. Iron-deficiency anemia b. Splenectomy

c. Alcohol toxicity d. Lead toxicity

*Decreases in A 1c occur in the following non-diabetic conditions: a. Hemolytic anemia b. chronic blood loss

*Pregnancy d. chronic renal failure. Interfering Factors:

*Presence of Hb F and H causes falsely elevated values. 2. Presence of Hb S, C, E, D, G, and Lepore (autosomal recessive mutation resulting in a hemoglobinopathy) causes falsely decreased values.

BUN (I	Blood Urea Nitrogen) * *	
Sample	:Serum	

11.20

mg/dL 7.0-23.0

CALCULATED



Page 4 of 12

View Reports on







Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 QN: U85110UP2003PLC193493

Patient Name Age/Gender	:Mrs.BANDANA SINGH :31 Y 0 M 0 D / F		Registere Collected		: 10/Nov/202 : 10/Nov/202	
UHID/MR NO	: CALI.0000051596		Received		: 10/Nov/202	
Visit ID	: CALI0170162425		Reported		: 10/Nov/202	
Ref Doctor	: Dr.Mediwheel - Arcofemi H	ealth Care Ltd.	Status		: Final Report	
	-		Sidius		. i mai neport	
		PARTMENT C				
	MEDIWHEEL	BANK OF BAR				
Test Name		Result	U	nit	Bio. Ref. Interva	al Method
Interpretation:						
-	UN levels can be seen in the fo	llowing:				
High-protein diet, D	ehydration, Aging, Certain medic	cations, Burns, Ga	strointestin	nal (GI) b	bleeding.	
Low BUN levels c	an be seen in the following:					
Low-protein diet, ov	verhydration, Liver disease.					
reatinine * *		0.63	mg/dl	0.5-1.2	20	MODIFIED JAFFES
ample:Serum Interpretation: The significance of s	single creatinine value must be int	erpreted in light o	of the patien	ts muscle	e mass. A patient v	vith a greater muscle
ample:Serum Interpretation: The significance of s mass will have a hig absolute creatinine o could be affected mi	single creatinine value must be int ther creatinine concentration. The concentration. Serum creatinine co ildly and may result in anomalous	erpreted in light o trend of serum cro oncentrations may	of the patien eatinine con y increase v	ts muscle ncentration when an 2	e mass. A patient v ons over time is m ACE inhibitor (AC	vith a greater muscle ore important than CE) is taken. The assa
Interpretation: The significance of s mass will have a hig absolute creatinine of could be affected mi lipemic.	ther creatinine concentration. The concentration. Serum creatinine co	erpreted in light o trend of serum cro oncentrations may	of the patien eatinine con y increase v	ts muscle ncentration when an 2	e mass. A patient v ons over time is m ACE inhibitor (AC hilic antibodies, h	vith a greater muscle ore important than CE) is taken. The assa
Interpretation: The significance of s mass will have a hig absolute creatinine of could be affected mid lipemic. Interpretation: Note:- Elevated uric acid	ther creatinine concentration. The concentration. Serum creatinine co	erpreted in light o trend of serum cro oncentrations may values if serum sa 4.47 4.47	of the patien eatinine cor y increase v amples hav mg/dl	tts muscle ncentratio when an A re heterop 2.5-6.0	e mass. A patient v ons over time is m ACE inhibitor (AC hilic antibodies, h	vith a greater muscle ore important than CE) is taken. The assay emolyzed, icteric or
Interpretation: The significance of s mass will have a hig absolute creatinine of could be affected mid lipemic. Interpretation: Note:- Elevated uric acid	ther creatinine concentration. The concentration. Serum creatinine co ildly and may result in anomalous I levels can be seen in the follow rotein diet, alcohol), Chronic kidm	erpreted in light o trend of serum cro oncentrations may values if serum sa 4.47 4.47	of the patien eatinine cor y increase v amples hav mg/dl	tts muscle ncentratio when an A re heterop 2.5-6.0	e mass. A patient v ons over time is m ACE inhibitor (AC hilic antibodies, h	vith a greater muscle ore important than CE) is taken. The assay emolyzed, icteric or
Interpretation: The significance of s mass will have a hig absolute creatinine of could be affected mi- lipemic. Interpretation: Note:- Elevated uric acid Drugs, Diet (high-pri- FT (WITH GAMM	ther creatinine concentration. The concentration. Serum creatinine concentration. Serum creating concentration. Serum	erpreted in light o trend of serum cro oncentrations may values if serum sa 4.47 4.47	of the patien eatinine cor y increase v amples hav mg/dl	tts muscle ncentratio when an A re heterop 2.5-6.0	e mass. A patient v ons over time is m ACE inhibitor (AC hilic antibodies, h	vith a greater muscle ore important than CE) is taken. The assay emolyzed, icteric or
Interpretation: The significance of significa	ther creatinine concentration. The concentration. Serum creatinine concentration. Serum creatinine concentration. Serum result in anomalous a levels can be seen in the follow rotein diet, alcohol), Chronic kidm A GT) ** , <i>Serum</i> Aminotransferase (AST)	erpreted in light o trend of serum cro oncentrations may values if serum sa 4.47 wing: ney disease, Hyper	of the patien eatinine cor y increase v amples hav mg/dl	its muscle ncentratio when an A re heterop 2.5-6.0	e mass. A patient v ons over time is m ACE inhibitor (AC hilic antibodies, h	vith a greater muscle ore important than CE) is taken. The assay emolyzed, icteric or URICASE
Interpretation: The significance of s mass will have a hig absolute creatinine of could be affected mi- lipemic. Interpretation: Note:- Elevated uric acid Drugs, Diet (high-pr FT (WITH GAMIM SGOT / Aspartate A SGPT / Alanine Am Gamma GT (GGT)	ther creatinine concentration. The concentration. Serum creatinine co- ildly and may result in anomalous I levels can be seen in the follow rotein diet, alcohol), Chronic kidm A GT) ** , <i>Serum</i> Aminotransferase (AST) inotransferase (ALT)	erpreted in light o trend of serum cro oncentrations may values if serum sa 4.47 wing: hey disease, Hyper 54.60 58.30 99.00	of the patien eatinine cor y increase v amples hav mg/ dl rtension, O U/L	e heterop 2.5-6.0 besity. 35 40 11-50	e mass. A patient v ons over time is m ACE inhibitor (AC hilic antibodies, h	vith a greater muscle ore important than CE) is taken. The assay emolyzed, icteric or URICASE
Interpretation: The significance of s mass will have a hig absolute creatinine of could be affected mi- lipemic. ric Acid ** ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p: FT (WITH GAMM SGOT / Aspartate A SGPT / Alanine Am Gamma GT (GGT) Protein	ther creatinine concentration. The concentration. Serum creatinine co- ildly and may result in anomalous I levels can be seen in the follow rotein diet, alcohol), Chronic kidm A GT) ** , <i>Serum</i> Aminotransferase (AST) inotransferase (ALT)	erpreted in light of trend of serum cre oncentrations may values if serum sa 4.47 wing: hey disease, Hyper 54.60 58.30 99.00 5.95	of the patien eatinine cor y increase v amples hav mg/dl mg/dl u/L U/L U/L U/L gm/dl	tts muscle ncentratio when an A re heterop 2.5-6.0 besity. >besity. <35 <40 11-50 6.2-8.0	e mass. A patient v ons over time is m ACE inhibitor (AC hilic antibodies, h	vith a greater muscle ore important than (E) is taken. The assay emolyzed, icteric or URICASE URICASE
Interpretation: The significance of s mass will have a hig absolute creatinine of could be affected mi- lipemic. ric Acid ** ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-pri- FT (WITH GAMM SGOT / Aspartate A SGPT / Alanine Am Gamma GT (GGT) Protein Albumin	ther creatinine concentration. The concentration. Serum creatinine co- ildly and may result in anomalous I levels can be seen in the follow rotein diet, alcohol), Chronic kidm A GT) ** , <i>Serum</i> Aminotransferase (AST) inotransferase (ALT)	erpreted in light of trend of serum creations may values if serum sa 4.47 wing: hey disease, Hyper 54.60 58.30 99.00 5.95 3.90	of the patien eatinine cor y increase v amples hav mg/dl mg/dl u/L U/L U/L gm/dl gm/dl	ets muscle ncentratio when an <i>A</i> e heterop 2.5-6.0 besity. vbesity. <35 <40 11-50 6.2-8.0 3.4-5.4	e mass. A patient v ons over time is m ACE inhibitor (AC hilic antibodies, h	vith a greater muscle ore important than CE) is taken. The assay emolyzed, icteric or URICASE IFCCWITHOUT P5P IFCCWITHOUT P5P IFCCWITHOUT P5P OPTIMIZED SZAZING BURET B.C.G.
Interpretation: The significance of s mass will have a hig absolute creatinine of could be affected mi- lipemic. ric Acid ** ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p: FT (WITH GAMM SGOT / Aspartate A SGPT / Alanine Am Gamma GT (GGT) Protein	ther creatinine concentration. The concentration. Serum creatinine co- ildly and may result in anomalous I levels can be seen in the follow rotein diet, alcohol), Chronic kidm A GT) ** , <i>Serum</i> Aminotransferase (AST) inotransferase (ALT)	erpreted in light of trend of serum cre oncentrations may values if serum sa 4.47 wing: hey disease, Hyper 54.60 58.30 99.00 5.95	of the patien eatinine cor y increase v amples hav mg/dl mg/dl u/L U/L U/L U/L gm/dl	tts muscle ncentratio when an A re heterop 2.5-6.0 besity. >besity. <35 <40 11-50 6.2-8.0	e mass. A patient v ons over time is m ACE inhibitor (AC hilic antibodies, h	vith a greater muscle ore important than (E) is taken. The assay emolyzed, icteric or URICASE URICASE









Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 GN: U85110UP2003PLC193493

Patient Name	: Mrs.BANDANA SINGH	Registered On	: 10/Nov/2024 09:24:11
Age/Gender	: 31 Y 0 M 0 D / F	Collected	: 10/Nov/2024 09:38:42
UHID/MR NO	: CALI.0000051596	Received	: 10/Nov/2024 11:41:22
Visit ID	: CALI0170162425	Reported	: 10/Nov/2024 13:19:59
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

Test Name	Result	U	nit Bio. Ref. In	terval Method
Alkaline Phosphatase (Total)	140.53	U/L	42.0-165.0	PNP/ AMP KINETIC
Bilirubin (Total)	1.25	mg/ dl	0.3-1.2	JENDRASSIK & GROF
Bilirubin (Direct)	0.65	mg/ dl	< 0.30	JENDRASSIK & GROF
Bilirubin (Indirect)	0.60	mg/dl	<0.8	JENDRASSIK & GROF
LIPID PROFILE (MINI)**, Serum				
Cholesterol (Total)	173.00	mg/ dl	<200 Desirable 200-239 Borderline > 240 High	CHOD-PAP High
HDL Cholesterol (Good Cholesterol)	69.30	mg/ dl	30-70	DIRECT ENZYMATIC
LDL Cholesterol (Bad Cholesterol)	86	mg/ dl	< 100 Optimal 100-129 Nr. Optimal/ Above Op 130-159 Borderline 160-189 High > 190 Very High	
VLDL	17.20	mg/ dl	10-33	CALCULATED
Triglycerides	86.00	mg/dl	< 150 Normal 150-199 Borderline 200-499 High >500 Very High	GPO-PAP

Dr. Anupam Singh (MBBS MD Pathology)



Page 6 of 12







Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 QN: U85110UP2003PLC193493

Patient Name	: Mrs.BANDANA SINGH	Registered On	: 10/Nov/2024 09:24:11
Age/Gender	: 31 Y 0 M 0 D / F	Collected	: 10/Nov/2024 14:46:40
UHID/MR NO	: CALI.0000051596	Received	: 10/Nov/2024 16:09:16
Visit ID	: CALI0170162425	Reported	: 10/Nov/2024 17:13:39
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

DEPARTMENT OF CLINICAL PATHOLOGY

MEDIWHEEL BANK OF BARODA FEM ALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
URINE EXAMINATION, ROUTINE ** ,	Urine			
Color	PALEYELLOW			
Specific Gravity	1.015			
Reaction PH	Acidic (5.0)			DIPSTICK
Appearance	QLEAR			
Protein	ABSENT	mg %	<10 Absent 10-40 (+) 40-200 (++) 200-500 (+++)	DIPSTICK
Sugar	ABSENT	gms%	>500 (++++) <0.5 (+) 0.5-1.0 (++) 1-2 (+++) >2 (++++)	DIPSTICK
Ketone	ABSENT	mg/ dl	Serum-0.1-3.0 Urine-0.0-14.0	BIOCHEMISTRY
Bile Salts	ABSENT			
Bile Pigments	ABSENT			
Bilirubin	ABSENT			DIPSTICK
Leucocyte Esterase	ABSENT			DIPSTICK
Urobilinogen(1:20 dilution)	ABSENT			
Nitrite	ABSENT			DIPSTICK
Blood	ABSENT			DIPSTICK
Microscopic Examination:				
Epithelial œlls	1-2/h.p.f			MICROSCOPIC EXAMINATION
Pus œlls	OCCASIONAL			
RBC:	ABSENT			MICROSCOPIC EXAMINATION
Cast	ABSENT			
Crystals	ABSENT			MICROSCOPIC EXAMINATION
Others	ABSENT			
SUGAR, FASTING STAGE ** , Urine				
Sugar, Fasting stage	ABSENT	gms%		



View Reports on







Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 CIN: U85110UP2003PLC193493

ľ	Patient Name	: Mrs.BANDANA SINGH	Registered On	: 10/Nov/2024 09:24:11
	Age/Gender	: 31 Y 0 M 0 D / F	Collected	: 10/Nov/2024 14:46:40
	UHID/MR NO	: CALI.0000051596	Received	: 10/Nov/2024 16:09:16
	Visit ID	: CALI0170162425	Reported	: 10/Nov/2024 17:13:39
	Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

DEPARTMENT OF CLINICAL PATHOLOGY

MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method

Interpretation:

(+) < 0.5 (++) 0.5-1.0 (+++) 1-2 (++++) > 2

SUGAR, PP STAGE** , Urine

Sugar, PP Stage

ABSENT

Interpretation:

(+) < 0.5 gms% (++) 0.5-1.0 gms% (+++) 1-2 gms% (++++) > 2 gms%



Page 8 of 12

Dr. Mamta Barthwal MD(Micro-Biology)





 \odot





Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 QN: U85110UP2003PLC193493

Patient Name	: Mrs.BANDANA SINGH	Registered On	: 10/Nov/2024 09:24:11
Age/Gender	: 31 Y 0 M 0 D / F	Collected	: 10/Nov/2024 09:38:42
UHID/MR NO	: CALI.0000051596	Received	: 10/Nov/2024 11:27:34
Visit ID	: CALI0170162425	Reported	: 10/Nov/2024 13:11:19
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

DEPARTMENT OF IMMUNOLOGY

MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
THYROID PROFILE - TOTAL** , Serum				
T3, Total (tri-iodothyronine)	104.65	ng/ dl	84.61–201.7	ALD
T4, Total (Thyroxine)	8.00	ug/dl	3.2-12.6	ALIA
TSH (Thyroid Stimulating Hormone)	1.850	μIU/mL	0.27 - 5.5	ALD
Interpretation:				
•		0.3-4.5 μIU/n	nL First Trimeste	er
		0.5-4.6 μIU/n	nL Second Trime	ster
		0.8-5.2 μIU/n	nL Third Trimest	er
		0.5-8.9 μIU/n	nL Adults	55-87 Years
		0.7-27 μIU/n	nL Premature	28-36 Week
		2.3-13.2 μIU/m		
		0.7-64 μIU/n	· · · · · · · · · · · · · · · · · · ·	·
		1-39 μIU		0-4 Days
		1.7-9.1 μIU/n	nL Child	2-20 Week

1) Patients having low T3 and T4 levels but high TSH levels suffer from primary hypothyroidism, cretinism, juvenile myxedema or autoimmune disorders.

2) Patients having high T3 and T4 levels but low TSH levels suffer from Grave's disease, toxic adenoma or sub-acute thyroiditis.

3) Patients having either low or normal T3 and T4 levels but low TSH values suffer from iodine deficiency or secondary hypothyroidism.

4) Patients having high T3 and T4 levels but normal TSH levels may suffer from toxic multinodular goiter. This condition is mostly a symptomatic and may cause transient hyperthyroidism but no persistent symptoms.

5) Patients with high or normal T3 and T4 levels and low or normal TSH levels suffer either from T3 toxicosis or T4 toxicosis respectively.

6) In patients with non thyroidal illness abnormal test results are not necessarily indicative of thyroidism but may be due to adaptation to the catabolic state and may revert to normal when the patient recovers.

7) There are many drugs for eg. Glucocorticoids, Dopamine, Lithium, Iodides, Oral radiographic dyes, etc. which may affect the thyroid function tests.

8) Generally when total T3 and total T4 results are indecisive then Free T3 and Free T4 tests are recommended for further confirmation along with TSH levels.

Dr. Anupam Singh (MBBS MD Pathology)



Page 9 of 12







Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 QN: U85110UP2003PLC193493

Patient Name	: Mrs.BANDANA SINGH	Registered On	: 10/Nov/2024 09:24:12
Age/Gender	: 31 Y 0 M 0 D / F	Collected	: 2024-11-10 12:50:45
UHID/MR NO	: CALI.0000051596	Received	: 2024-11-10 12:50:45
Visit ID	: CALI0170162425	Reported	: 10/Nov/2024 12:51:03
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

DEPARTMENT OF X-RAY

MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

X-RAY DIGITAL CHEST PA (500 mA COMPUTERISED UNIT SPOT FILM DEVICE)

DIGITAL CHEST P-A VIEW

- Costo-phrenic angles are bilaterally clear.
- Trachea is central in position.
- Cardiac size & contours are normal.
- Hilar shadows are normal.
- Pulmonary vascularity & distribution are normal.
- Pulmonary parenchyma did not reveal any significant lesion.

IMPRESSION:

• NO SIGNIFICANT DIAGNOSTIC ABNORMALITY SEEN.

Dr. Pankaj Kumar Gupta (M.B.B.S D.M.R.D)



Page 10 of 12







Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 QN: U85110UP2003PLC193493

Patient Name	: Mrs.BANDANA SINGH	Registered On	: 10/Nov/2024 09:24:12
Age/Gender	: 31 Y 0 M 0 D / F	Collected	: 2024-11-10 10:39:40
UHID/MR NO	: CALI.0000051596	Received	: 2024-11-10 10:39:40
Visit ID	: CALI0170162425	Reported	: 10/Nov/2024 10:42:18
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd. -	Status	: Final Report

DEPARTMENT OF ULTRASOUND

MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

ULTRASOUND WHOLE ABDOM EN (UPPER & LOWER)

LIVER

• The liver is normal in size ~14.2 cm in longitudinal span and has a normal homogenous echotexture. No focal lesion is seen.

PORTAL SYSTEM

- The intra hepatic portal channels are normal.
- The portal vein is not dilated.
- Porta hepatis is normal.

BILIARY SYSTEM

- The intra-hepatic biliary radicles are normal.
- Common duct is not dilated.
- The gall bladder is normal in size and has regular walls. Lumen of the gall bladder is anechoic.

PANCREAS

• The pancreas is normal in size and shape and has a normal homogenous echotexture. Pancreatic duct is not dilated.

KIDNEYS

- Right kidney is normal in size ~9.4 x 4.3 cm position and cortical echotexture. Corticomedullary demarcation is maintained.
- Left kidney is normal in size ~9.7 x 4.5 cm position and cortical echotexture. Corticomedullary demarcation is maintained.
- The collecting system of both the kidneys are not dilated.

SPLEEN

• The spleen is normal in size ~9.6 cm and has a normal homogenous echo-texture.

ILIAC FOSSAE & PERITONEUM

- Scan over the iliac fossae does not reveal any fluid collection or mass.
- No free fluid is noted in peritoneal cavity.
- Visualized bowel loops are gaseous and grossly appear normal in caliber, peristalsis and wall thickness.

URINARY BLADDER









Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 QN: U85110UP2003PLC193493

Patient Name	: Mrs.BANDANA SINGH	Registered On	: 10/Nov/2024 09:24:12
Age/Gender	: 31 Y 0 M 0 D / F	Collected	: 2024-11-10 10:39:40
UHID/MR NO	: CALI.0000051596	Received	: 2024-11-10 10:39:40
Visit ID	: CALI0170162425	Reported	: 10/Nov/2024 10:42:18
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

DEPARTMENT OF ULTRASOUND

MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

 The urinary bladder is normal. Bladder wall is normal in thickness and is regular. No calculus seen.

UTERUS

- The uterus is anteverted and normal in size ~ 6.4 x 4.6 x 2.9 cm.
- It has a homogenous myometrial echotexture.
- The endometrium is seen in midline. ET measures ~5.9 mm.
- Cervix is normal.

ADNEXA & OVARIES

- Both ovaries are normal in size volume and echotexture.
- Right ovary measures ~2.8 x 2.3 x 1.5 cm (volume~4.8 cc).
- Left ovary measures ~3.1 x 2.7 x 1.8 cm (volume~5.2 cc).
- Adnexa are normal.

FINAL IMPRESSION:-

NO SIGNIFICANT ABNORMALITY SEEN IN PRESENT STUDY.

Adv: Clinico-pathological correlation and follow-up.

*** End Of Report ***

(**) Test Performed at Chandan Speciality Lab.

Result/s to Follow: STOOL, ROUTINE EXAMINATION, ECG / EKG, Tread Mill Test (TMT), PAP SMEAR FOR CYTOLOGICAL EXAMINATION





This report is not for medico legal purpose. If clinical correlation is not established, kindly repeat the test at no additional cost within seven days

Facilities: MRI, CT scan, DR X-ray, Ultrasound, Sonomammography, Digital Mammography, ECG (Bedside also), 2D Echo, TMT, Holter, OPG, EEG, NCV, EMG & BERA, Audiometry, BMD, PFT, Fibroscan, Bronchoscopy, Colonoscopy and Endoscopy, Allergy Testing, Biochemistry & Immunoassay, Hematology, Microbiology & Serology, Histopathology & Immunohistochemistry, Cytogenetics and Molecular Diagnostics and Health Checkups 365 Days Open

*Facilities Available at Select Location Page 12 of 12



View Reports on Chandan 24x7 App

