Age / Sex : 27 Year(s) / Female

 PID No.
 : MED111393737
 Register On
 : 26/11/2022 8:39 AM

 SID No.
 : 422079429
 Collection On
 : 26/11/2022 9:43 AM

Report On : 26/11/2022 5:53 PM

: 05/12/2022 5:38 PM

Type : OP Printed On

Ref. Dr : MediWheel



Investigation HAEMATOLOGY	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood/Spectrophotometry)	13.3	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	42.3	%	37 - 47
RBC Count (EDTA Blood)	4.82	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (EDTA Blood)	87.8	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood)	27.7	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood)	31.5	g/dL	32 - 36
RDW-CV (EDTA Blood)	15.5	%	11.5 - 16.0
RDW-SD (EDTA Blood)	47.63	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood)	10900	cells/cu.mm	4000 - 11000
Neutrophils (EDTA Blood)	56.4	%	40 - 75
Lymphocytes (EDTA Blood)	33.4	%	20 - 45
Eosinophils (EDTA Blood)	4.6	%	01 - 06
Monocytes (EDTA Blood)	5.3	%	01 - 10



 PID No.
 : MED111393737
 Register On
 : 26/11/2022 8:39 AM

 SID No.
 : 422079429
 Collection On
 : 26/11/2022 9:43 AM

 Age / Sex
 : 27 Year(s) / Female
 Report On
 : 26/11/2022 5:53 PM

Printed On

MEDALL

< 20

Type : OP

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
Basophils (Blood)	0.3	%	00 - 02
INTERPRETATION: Tests done on Automated Five l	Part cell counter. All a	abnormal results are r	eviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood)	6.15	10^3 / μ1	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood)	3.64	10^3 / μ1	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood)	0.50	10^3 / μ1	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood)	0.58	10^3 / μ1	< 1.0
Absolute Basophil count (EDTA Blood)	0.03	10^3 / μΙ	< 0.2
Platelet Count (EDTA Blood)	271	10^3 / μΙ	150 - 450
MPV (EDTA Blood)	11.3	fL	8.0 - 13.3
PCT (EDTA Blood/Automated Blood cell Counter)	0.31	%	0.18 - 0.28

: 05/12/2022 5:38 PM



4

APPROVED BY

mm/hr

ESR (Erythrocyte Sedimentation Rate)

(Citrated Blood)

Age / Sex : 27 Year(s) / Female

 PID No.
 : MED111393737
 Register On : 26/11/2022 8:39 AM

 SID No.
 : 422079429
 Collection On : 26/11/2022 9:43 AM

Report On : 26/11/2022 5:53 PM

: 05/12/2022 5:38 PM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation BIOCHEMISTRY	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	0.55	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.21	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.34	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	24.15	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	23.81	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	16.78	U/L	< 38
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	80.6	U/L	42 - 98
Total Protein (Serum/Biuret)	7.60	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	5.22	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	2.38	gm/dL	2.3 - 3.6
A : G RATIO (Serum/Derived)	2.19		1.1 - 2.2



VERIFIED BY



 PID No.
 : MED111393737
 Register On
 : 26/11/2022 8:39 AM

 SID No.
 : 422079429
 Collection On
 : 26/11/2022 9:43 AM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	163.59	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	108.25	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

: 05/12/2022 5:38 PM

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual_circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	42.45	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/Calculated)	99.4	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >=190
VLDL Cholesterol (Serum/Calculated)	21.7	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	121.1	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220



DR SHAMIM JAVED MD PATHOLOGY KMC 88902

 PID No.
 : MED111393737
 Register On
 : 26/11/2022 8:39 AM

 SID No.
 : 422079429
 Collection On
 : 26/11/2022 9:43 AM

Printed On

Type : OP

Ref. Dr : MediWheel



InvestigationObservedUnitBiologicalValueReference Interval

INTERPRETATION: 1.Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

: 05/12/2022 5:38 PM

Total Cholesterol/HDL Cholesterol Ratio
(Serum/Calculated)

3.9

Optimal: < 3.3

Low Risk: 3.4 - 4

Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio 2.6 Optimal: < 2.5

(TG/HDL) Mild to moderate risk: 2.5 - 5.0

(Serum/Calculated) High Risk: > 5.0

LDL/HDL Cholesterol Ratio

2.3 Optimal: 0.5 - 3.0

(Serum/Calculated)
Borderline: 3.1 - 6.0
High Risk: > 6.0



VERIFIED BY



> : MED111393737 Register On : 26/11/2022 8:39 AM : 422079429

Collection On : 26/11/2022 9:43 AM

Printed On

Age / Sex : 27 Year(s) / Female Report On : 26/11/2022 5:53 PM **Type** : OP : 05/12/2022 5:38 PM

Ref. Dr : MediWheel

PID No.

SID No.



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/ <i>HPLC</i>)	5.4	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 108.28 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.



VERIFIED BY



PID No. : MED111393737 Register On : 26/11/2022 8:39 AM

: 422079429 SID No. Collection On : 26/11/2022 9:43 AM

Age / Sex : 27 Year(s) / Female Report On : 26/11/2022 5:53 PM

Printed On

: OP

Type

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
-	Value		Reference Interval

: 05/12/2022 5:38 PM

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.40 ng/ml 0.7 - 2.04

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

4.2 - 12.0T4 (Tyroxine) - Total 9.08 μg/dl

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

μIU/mL 0.35 - 5.50TSH (Thyroid Stimulating Hormone) 1.21

(Serum/ECLIA)

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester: 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



PID No. : MED111393737 Register On : 26/11/2022 8:39 AM : 422079429 SID No. Collection On : 26/11/2022 9:43 AM

Age / Sex : 27 Year(s) / Female Report On : 26/11/2022 5:53 PM

Printed On

Type : OP

Ref. Dr : MediWheel

: 05/12/2022 5:38 PM

Investigation CLINICAL PATHOLOGY PHYSICAL EXAMINATION (URINE COMPLETE)	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Colour (Urine)	Yellow		Yellow to Amber
Appearance (Urine)	Clear		Clear
Volume(CLU) (Urine)	15		
<u>CHEMICAL EXAMINATION (URINE</u> <u>COMPLETE)</u>			
рН	5.5		4.5 - 8.0

(Urine) Ketone Negative Negative (Urine)

1.010

Normal

Urobilinogen

Specific Gravity

(Urine)

(Urine)

Negative Blood

(Urine) Nitrite Negative

(Urine)

Bilirubin Negative

(Urine) (Urine)

Protein Negative Negative

> Dr Anusha.K.S Sr.Consultant Pathologist Reg No: 100674 **APPROVED BY**

1.002 - 1.035

Normal

Negative

Negative

Negative

PID No. : MED111393737 : 422079429 SID No.

Age / Sex : 27 Year(s) / Female

Type : OP

: MediWheel

Register On : 26/11/2022 8:39 AM

Collection On : 26/11/2022 9:43 AM

Report On : 26/11/2022 5:53 PM

Printed On : 05/12/2022 5:38 PM



<u>Investigation</u>	<u>Observed</u> <u>Unit</u> <u>Value</u>	<u>Biological</u> <u>Reference Interval</u>
Glucose	Negative	Negative

(Urine/GOD - POD)

Leukocytes(CP) Negative

Ref. Dr

MICROSCOPIC EXAMINATION (URINE COMPLETE)

NIL 0-2/hpf Pus Cells

(Urine)

Epithelial Cells 0-2 /hpf **NIL**

(Urine)

RBCs NIL /HPF **NIL**

(Urine)

NIL Others

(Urine)

INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.

/hpf NIL Casts NIL

(Urine)

NIL NIL Crystals /hpf

(Urine)

Dr Anusha.K.S Sr.Consultant Pathologist Reg No: 100674 APPROVED BY

PID No. : MED111393737

SID No. : 422079429 **Age / Sex** : 27 Year(s) / Female

Type : OP

Ref. Dr : MediWheel

Register On : 26/11/2022 8:39 AM

Collection On : 26/11/2022 9:43 AM

Report On : 26/11/2022 5:53 PM

Printed On : 05/12/2022 5:38 PM

MEDALI

InvestigationObservedUnitBiologicalValueReference Interval

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING

(EDTA Blood/Agglutination)

'B' 'Positive'



 PID No.
 : MED111393737
 Register On
 : 26/11/2022 8:39 AM

 SID No.
 : 422079429
 Collection On
 : 26/11/2022 9:43 AM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	9.2		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	99.23	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

: 05/12/2022 5:38 PM

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine) (Urine - F/GOD - POD)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD-PAP)	92.96	mg/dL	70 - 140

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Blood Urea Nitrogen (BUN)	6.6	mg/dL	7.0 - 21
(Serum/Urease UV / derived)			
Creatinine	0.71	mg/dL	0.6 - 1.1

(Serum/Modified Jaffe)

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine etc

Uric Acid 2.92 mg/dL 2.6 - 6.0 (Serum/Enzymatic)



VERIFIED BY



APPROVED BY

-- End of Report --



Name	Mrs.Prity kumari	ID	MED111393737
Age & Gender	27/FEMALE	Visit Date	26/11/2022
Ref Doctor Name	MediWheel		

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER shows normal in shape, size and has uniform echopattern. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER shows normal shape and has clear contents. Gall bladder wall is of normal thickness. CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN shows normal shape, size and echopattern.

No demonstrable Para -aortic lymphadenopathy.

KIDNEYS move well with respiration and have normal shape, size and echopattern. Cortico- medullary differentiations are well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney	10.3	1.7
Left Kidney	11.0	1.8

URINARY BLADDER shows normal shape and wall thickness. It has clear contents. No evidence of diverticula.

UTERUS is anteverted and normal in size. It has uniform myometrial echopattern.

Endometrial thickness measures 7mm

Uterus measures as follows: LS: 8.2cms AP: 3.6cms TS: 4.3cms.

OVARIES are normal in size, shape and echotexture. No focal lesion seen.

Ovaries measure as follows: **Right ovary**: 3.9 x 1.9cms **Left ovary**: 2.7 x 1.8cms

POD & adnexae are free.

No evidence of ascites/pleural effusion.

IMPRESSION:

REPORT DISCLAIMER

- 1.This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- $11. Disputes, if any \ , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only. \\$



Name	Mrs.Prity kumari	ID	MED111393737
Age & Gender	27/FEMALE	Visit Date	26/11/2022
Ref Doctor Name	MediWheel		

> NO SIGNIFICANT ABNORMALITY DETECTED.

DR. APARNA CONSULTANT RADIOLOGIST A/vp

REPORT DISCLAIMER

^{1.}This is only a radiological imperssion.Like other investigations, radiological investication also have limitation. Therefore radiological reports should be interpreted in correlation with clinical and pathological findings.

^{2.} The results reported here in are subject to interpretation by qualified medical professionals only.

^{3.} Customer identities are accepted provided by the customer or their representative.

^{4.}information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.

^{5.}If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.

^{6.}Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.

^{7.}Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,

^{8.}If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.

^{9.}Liability is limited to the extend of amount billed.

^{10.}Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.

^{11.}Disputes,if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.