# Dr. Goyal's Path Lab & Imaging Centre

B-51, 6	Ganesh Nagar,	Opp. Janpath	Corner, New S	Sanga 🕝 😝	neral	30 <b>20 9</b> 1	ysical	Examin	ation
Tolo .	0444 220224	C 4040707	0007040707						


Tele: 0141-2293346, 4049787, 9887049787

Vebsite: www.drgoyalspathlab.com	E-mail	: drg	oyalpiy	yush@g	mail.com
Vebsite : www.drgoyalspathlab.com   Date of Examination	n:	A .	09	-20	22

Name: <u>LOKCSH MCCNA</u> Age: <u>32</u> DOB: <u>13-06-1990</u> Sex: <u>Ma</u> le
Referred By: 806
Photo ID:AADHAR ID#:attached
Ht: <u>16 7</u> (cm) Wt: <u>96</u> (Kg)
Chest (Expiration): 105 (cm) Abdomen Circumference: 120 (cm)
Blood Pressure: 129/83 mm Hg PR: 18/min RR: 17/min Temp: website
вмі 34.4
Eye Examination: Niston somal 6/6, N/6
Other: Dol ugoificant
On examination he/she appears physically and mentally fit: Yes/No
Signature Of Examine :Name of Examinee:
Name Medical Examiner








भारत सरकार Unique Identification Authority of India Government of India

नामांकन क्रम/ Enrolment No.: 2036/81005/58359

लोकेश मीना-Lokesh Meena S/O: Babu Lal Meena gaon- jharoli post- raneela tah- bamanwas Iharoli Ranila Sawai Madhopur Rajasthan - 322211 9610510967





आपका आधार क्रमांक / Your Aadhaar No. :

8513 8044 9001

मेरा आधार, मेरी पहचान



#### भारत सरकार

Government of India



लोकेश मीना Lokesh Meena जन्म तिथि/DOB: 13/06/1990 TENY MALE

8513 8044 9001









- आधार पहचान का प्रमाण है, नागरिकता का नहीं ।
- पहचान का प्रमाण ऑनलाइन ऑथेन्टिकेशन दारा प्राप्त करें ।
- यह एक इलेक्ट्रॉनिक प्रक्रिया द्वारा बना हुआ पत्र है ।

#### INFORMATION

- Aadhaar is a proof of identity, not of citizensh
- To establish identity, authenticate online.
- This is electronically generated letter.
- आधार देश भर में मान्य है।
- आधार भविष्य में सरकारी और गैर-सरकारी सेवाओं का लाभ उठाने में उपयोगी होगा।
- Aadhaar is valid throughout the country .
- Aadhaar will be helpful in availing Governme and Non-Government services in future

Dr. Piyush Goyal M.B.B.S., D.M.R.D. RMC Res. No., 017996



भारतीय विशिष्ट पहुंचान प्राधिकरण

Unique Identification Authority of India

आत्मजः बाबूला्ल\_मीना, गोय- झाडोली पोस्ट- रामीला तह-जारमणः बाबुरास्त नाना, गाव- झाडा बामनवास, झाडोली, सवाई माधोपुर, राजस्थान - 322211

Address:

S/O: Babu Lai Meena, gaon- jharoli post-raneela tah- bamarwas, Jharoli, Sawai Madhopur, Rajasthan - 322211



8513 8044 9001







Allengers ECG (Pisces)(PI\$212160118 12347 / MR LOKESH MEENA / 32 Yrs / M/ Non Smoker
Heart Rate: 92 bpm / / Refd By:: BOB / Tested On: 10-Sep-22 12:00:13 / HF 0.05 Hz - LF 100 Hz / Notch 50 Hz / Sn 1.00 Cm/mV / Sw 25 DR. GOYALS PATH LAB & IMAGING CENTER **≤**[ avRV Dr. Names No. 35 103 (ARCHO) ( 465 avF S ECG mm/s

## Dr. Goyal's Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date :- 10/09/2022 10:04:51

NAME :- Mr. LOKESH MEENA

Sex / Age :- Male

Sample Type :- EDTA

32 Yrs

Company :- MediWheel

Patient ID :-12222285

Ref. By Dr:- BOB

Lab/Hosp :-

Final Authentication: 10/09/2022 15:08:13

Sample Collected Time 10/09/2022 10:21:30

HAEMATOLOGY

W-97 117 117 117 117 117 117 117 117 117 1	HAEMAIO	LUGI	
Test Name	Value	Unit	Biological Ref Interval
BOB PACKAGE BELOW 40MALE			
HAEMOGARAM			
HAEMOGLOBIN (Hb)	13.1	g/dL	13.0 - 17.0
TOTAL LEUCOCYTE COUNT	8.98	/cumm	4.00 - 10.00
DIFFERENTIAL LEUCOCYTE COUNT			
NEUTROPHIL	56.1	%	40.0 - 80.0
LYMPHOCYTE	33.8	%	20.0 - 40.0
EOSINOPHIL	5.6	%	1.0 - 6.0
MONOCYTE	4.3	%	2.0 - 10.0
BASOPHIL	0.2	%	0.0 - 2.0
NEUT#	5.04	10^3/uL	1.50 - 7.00
LYMPH#	3.04	10^3/uL	1.00 - 3.70
EO#	0.50 H	10^3/uL	0.00 - 0.40
MONO#	0.38	10^3/uL	0.00 - 0.70
BASO#	0.02	10^3/uL	0.00 - 0.10
TOTAL RED BLOOD CELL COUNT (RBC)	4.10 L	x10^6/uL	4.50 - 5.50
HEMATOCRIT (HCT)	38.80 L	%	40.00 - 50.00
MEAN CORP VOLUME (MCV)	94.7	fL	83.0 - 101.0
MEAN CORP HB (MCH)	32.1 H	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	33.9	g/dL	31.5 - 34.5
PLATELET COUNT	368	x10^3/uL	150 - 410
RDW-CV	16.1 H	%	11.6 - 14.0
MENTZER INDEX	23.10		

The Mentzer index is used to differentiate iron deficiency anemia from beta thalassemia trait. If a CBC indicates microcytic anemia, these are two of the most likely causes, making it necessary to distinguish between them.

If the quotient of the mean corpuscular volume divided by the red blood cell count is less than 13, thalassemia is more likely. If the result is greater than 13, then iron-deficiency anemia is more likely.

BANWARI Technologist

Page No: 1 of 12





Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 10/09/2022 10:04:51 Date

NAME :- Mr. LOKESH MEENA

Ref. By Dr:- BOB

Patient ID: -12222285

Sex / Age :- Male

32 Yrs

Lab/Hosp :-

Company:- MediWheel

Sample Type :- EDTA

Sample Collected Time 10/09/2022 10:21:30

Final Authentication: 10/09/2022 15:08:13

#### HAEMATOLOGY

Value **Test Name** Unit **Biological Ref Interval** 

Erythrocyte Sedimentation Rate (ESR)

24 H

mm/hr.

00 - 13

(ESR) Methodology: Measurment of ESR by cells aggregation.

Instrument Name : Indepedent form Hematocrit value by Automated Analyzer (Roller-20)

: ESR test is a non-specific indicator ofinflammatory disease and abnormal protein states. Interpretation

The test in used to detect, follow course of a certain disease (e.g.-tuberculosis, rheumatic fever, myocardial infarction

Levels are higher in pregnency due to hyperfibrinogenaemia.

The "3-figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia (CRC) Methodology TLC DLC Fluorescent Flow cytometry, HB SLS method, TRBC, PCV, PLT Hydrodynamically focused Impedance. and MCH, MCV, MCHC, MENTZER INDEX are calculated. InstrumentName: Sysmex 6 part fully automatic analyzer XN-L, Japan

BANWARI **Technologist** 

Page No: 2 of 12





Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 10/09/2022 10:04:51

Patient ID: -12222285

NAME :- Mr. LOKESH MEENA

Ref. By Dr:- BOB

Sex / Age :- Male

Lab/Hosp :-

Company :- MediWheel

Sample Type :- EDTA, KOx/Na FLUORIDE-F, KSawhpale-Collegue ETPR: UR09E,028 INE29830

Final Authentication: 10/09/2022 15:40:04

#### HAEMATOLOGY

**Test Name** 

Value

Unit

**Biological Ref Interval** 

**BLOOD GROUP ABO** 

"A" POSITIVE

BLOOD GROUP ABO Methodology: Haemagglutination reaction Kit Name: Monoclonal agglutinating antibodies (Span clone).

FASTING BLOOD SUGAR (Plasma) Method:- GOD PAP

96.4

mg/dl

75.0 - 115.0

Impaired glucose tolerance (IGT)	111 - 125 mg/dL	
Diabetes Mellitus (DM)	> 126 mg/dL	

Instrument Name: Randox Rx Imola Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases .

BLOOD SUGAR PP (Plasma)

110.8

mg/dl

70.0 - 140.0

Method:- GOD PAP Instrument Name: Randox Rx Imola Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases .

URINE SUGAR (FASTING)
Collected Sample Received

Nil

Nil

URINE SUGAR PP Collected Sample Received

Nil

Nil

BANWARI, MUKESHSINGH, SURENDRAMEENA **Technologist** 

Page No: 3 of 12



Dr. Rashmi Bakshi MBBS. MD (Path) RMC No. 17975/008828 Dr. Chandrika Gupta



Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 10/09/2022 10:04:51

NAME :- Mr. LOKESH MEENA

Sex / Age :- Male 32 Yrs

Company :- MediWheel Sample Type :- STOOL

Patient ID: -12222285

Ref. By Dr:- BOB

Lab/Hosp:-

Final Authentication: 10/09/2022 12:23:50

#### **CLINICAL PATHOLOGY**

Sample Collected Time 10/09/2022 10:21:30

**Test Name** 

Value

Unit

**Biological Ref Interval** 

#### STOOL ANALYSIS

#### PHYSICAL EXAMINATION

COLOUR

CONSISTENCY

**MUCUS** 

**BLOOD** 

MICROSCOPIC EXAMINATION

RBC's

WBC/HPF

MACROPHAGES

OVA

CYSTS

**TROPHOZOITES** 

CHARCOT LEYDEN CRYSTALS

OTHERS Collected Sample Received

YELLOW BROWN

SEMI SOLID

ABSENT

ABSENT

NIL /HPF

/HPF

ABSENT

NIL.

ABSENT

ABSENT

ABSENT

ABSENT

ABSENT

SURENDRAMEENA **Technologist** 

Page No: 4 of 12



Dr. Chandrika Gupta MBBS.MD (Path) RMC NO. 21021/008037

'CONDITIONS OF REPORTING SEE OVER LEAF"



Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 10/09/2022 10:04:51 Date

NAME :- Mr. LOKESH MEENA

Sex / Age :- Male

32 Yrs

Company :- MediWheel

Sample Type :- PLAIN/SERUM

Patient ID: -12222285

Ref. By Dr:- BOB

Lab/Hosp :-

Sample Collected Time 10/09/2022 10:21:30

Final Authentication: 10/09/2022 14:07:43

#### BIOCHEMISTRY

	DIOCERDINA		
Test Name	Value	Unit	Biological Ref Interval
LIPID PROFILE			
TOTAL CHOLESTEROL Method:- Enzymatic Endpoint Method	159.32	mg/dl	Desirable <200 Borderline 200-239 High> 240
TRIGLYCERIDES Method:- GPO-PAP	91.32	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500
DIRECT HDL CHOLESTEROL Method:- Direct clearance Method	31.97	mg/dl	Low < 40 High > 60
DIRECT LDL CHOLESTEROL Method:- Direct clearance Method	112.13	mg/dl	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190
VLDL CHOLESTEROL Method:- Calculated	18.26	mg/dl	0.00 - 80.00
T.CHOLESTEROL/HDL CHOLESTEROL RATIO Method:- Calculated	4.98 H		0.00 - 4.90
LDL / HDL CHOLESTEROL RATIO Method:- Calculated	3.51 H		0.00 - 3.50
TOTAL LIPID Method:- CALCULATED TOTAL CHOINESTROY Instrument Name: Randov Ry Imola	470.42	mg/dl	400.00 - 1000.00

TOTAL CHOLESTEROL InstrumentName: Randox Rx Imola Interpretation: Cholesterol measurements are used in the diagnosis and treatments of lipid lipoprotein meta-

TRIGLYCERIDES InstrumentName: Randox Rx Imola Interpretation: Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction.

DIRECT HDLCHOLESTERO InstrumentName:Randox Rx Imola Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.

DIRECT LDL-CHOLESTEROLInstrumentName: Randox Rx Imola Interpretation: Accurate measurement of LDL-Cholesterol is of vital importance in therapies which focus on lipid reduction to prevent atherosclerosis or reduce its progress and to avoid plaque rupture. TOTAL LIPID AND VLDL ARE CALCULATED

MUKESHSINGH

Page No: 5 of 12





Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 10/09/2022 10:04:51 Date

NAME :- Mr. LOKESH MEENA

Sex / Age :- Male

Company :- MediWheel Sample Type :- PLAIN/SERUM Patient ID: -12222285

Ref. By Dr:- BOB

Lab/Hosp:-

Final Authentication: 10/09/2022 14:07:43

Sample Collected Time 10/09/2022 10:21:30

	BIOCHEM	ISTRY	
Test Name	Value	Unit	Biological Ref Interval
LIVER PROFILE WITH GGT			
SERUM BILIRUBIN (TOTAL) Method:- Colorimetric method	0.42	mg/dl	Up to - 1.0 Cord blood <2 mg/dL Premature < 6 days <16mg/dL Full-term < 6 days= 12 mg/dL 1month - <12 months <2 mg/dL 1-19 years <1.5 mg/dL Adult - Up to - 1.2 Ref-(ACCP 2020)
SERUM BILIRUBIN (DIRECT) Method:- Colorimetric Method	0.10	mg/dL	Adult - Up to 0.25 Newborn - <0.6 mg/dL >- 1 month - <0.2 mg/dL
SERUM BILIRUBIN (INDIRECT) Method:- Calculated	0.32	mg/dl	0.30-0.70
SGOT Method:- IFCC	56.0 H	U/L	Men- Up to - 37.0 Women - Up to - 31.0
SGPT Method:- IFCC	83.9 H	U/L	Men- Up to - 40.0 Women - Up to - 31.0
SERUM ALKALINE PHOSPHATASE Method:-AMP Buffer	77.90	IU/L	30.00 - 120.00
SERUM GAMMA GT Method:- IFCC	45.40	U/L	11.00 - 50.00
SERUM TOTAL PROTEIN Method:- Biuret Reagent	7.04	g/dl	6.40 - 8.30
SERUM ALBUMIN Method:- Bromocresol Green	4.24	g/dl	3.80 - 5.00
SERUM GLOBULIN Method:- CALCULATION	2.80	gm/dl	2.20 - 3.50
A/G RATIO	1.51		1.30 - 2.50

Total BilirubinMethodology:Colorimetric method InstrumentName:Randox Rx Imola Interpretation An increase in bilirubin concentration in the serum occurs in toxic or infectious diseases of the liver e.g. hepatitis B or obstruction of the bile duct and in rhesus incompatible babies. High levels of unconjugated bilirubin indicate that too much haemoglobin is being destroyed or that the liver is not actively treating the haemoglobin it is receiving.

AST Aspartate Aminotransferase Methodology: IFCC InstrumentName: Randox Rx Imola Interpretation: Elevated levels of AST can signal myocardial infarction, hepatic disease, muscular dystrophy and organ damage. Although heart muscle is found to have the most activity of the enzyme, significant activity has also been seen in the brain, liver, gastric mucosa, adipose tissue and kidneys of hum ALT Alanine Aminotransferase Methodology: IFCCInstrumentName:Randox Rx Imola Interpretation: The enzyme ALT has been found to be in highest concentrations in the liver, with decreasing concentrations found in kidney, heart, skeletal muscle, pancreas, spleen and lung tissue respectively. Elevated levels of the transaminases can indicate myocardial infarction, hepatic disease, muscular

dystrophy and organ damage.

Alkaline Phosphatase Methodology:AMP Buffer InstrumentName:Randox Rx Imola Interpretation:Measurements of alkaline phosphatase are of use in the diagnosis, treatment and investigation of hepatobilary disease and in bone disease associated with increased osteoblastic activity. Alkaline phosphatase is also used in the diagnosis of parathyroid and intestinal disease.

TOTAL PROTEIN Methodology: Biuret Reagent InstrumentName: Randox Rx Imola Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

ALBUMIN (ALB) Methodology: Bromocresol Green InstrumentName:Randox Rx Imola Interpretation: Albumin measurements are used in the diagnosis and treatment of numerous diseases involving

MUKESHSINGH

Page No: 6 of 12





Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date :- 10/09/2022 10:04:51

NAME :- Mr. LOKESH MEENA

Sex / Age :- Male 32 Yrs

Company :- MediWheel

Sample Type :- PLAIN/SERUM

Patient ID: -12222285

Ref. By Dr:- BOB

Lab/Hosp:-

Sample Collected Time 10/09/2022 10:21:30

Final Authentication: 10/09/2022 14:07:43

**BIOCHEMISTRY** 

Test Name Value Unit Biological Ref Interval

primarily the liver or kidneys. Globulin & A/G ratio is calculated.

Instrument Name Randox Rx Imola Interpretation: Elevations in GGT levels areseen earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or post-hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal) are observed with infectious hepatitis.

MUKESHSINGH

Page No: 7 of 12





Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 10/09/2022 10:04:51

NAME :- Mr. LOKESH MEENA

Sex / Age :- Male

Sample Type :- PLAIN/SERUM

32 Yrs

Company :- MediWheel

Ref. By Dr:- BOB Lab/Hosp :-

Patient ID: -12222285

Sample Collected Time 10/09/2022 10:21:30

Final Authentication: 10/09/2022 14:07:43

BIOCHEMISTRY

DIOCHEMICIKI			
Test Name	Value	Unit	Biological Ref Interval
SERUM CREATININE Method:- Colorimetric Method	0.90	mg/dl	Men - 0.6-1.30 Women - 0.5-1.20
SERUM URIC ACID Method:- Enzymatic colorimetric	5.86	mg/dl	Men - 3.4-7.0 Women - 2.4-5.7

MUKESHSINGH

Page No: 8 of 12





Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date :- 10/09/2022 10:04:51

NAME :- Mr. LOKESH MEENA

Sex / Age :- Male 32 Yrs

Company :- MediWheel

Sample Type :- PLAIN/SERUM

Patient ID :-12222285

Ref. By Dr:- BOB

Lab/Hosp :-

Sample Collected Time 10/09/2022 10:21:30

Final Authentication: 10/09/2022 14:07:43

#### BIOCHEMISTRY

	DIOCHEN	HSIKI			
Test Name	Value	Unit	Biological Ref Interval		
BLOOD UREA NITROGEN (BUN)	18.3	mg/dl	0.0 - 23.0		

MUKESHSINGH

Page No: 9 of 12





Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date :- 10/09/2022 10:04:51

NAME :- Mr. LOKESH MEENA

Sex / Age :- Male

Sample Type :- EDTA

32 Yrs

Company :- MediWheel

Patient ID :-12222285

Ref. By Dr:- BOB

Lab/Hosp :-

Sample Collected Time 10/09/2022 10:21:30

Final Authentication : 10/09/2022 15:08:13

HAEMATOLOGY

Test Name Value Unit Biological Ref Interval

GLYCOSYLATED HEMOGLOBIN (HbA1C)

Method:- HPLC

6.3 H

%

Non-diabetic: < 5.7 Pre-diabetics: 5.7-6.4

Diabetics: = 6.5 or higher ADA Target: 7.0

ADA Target: 7.0 Action suggested: > 6.5

Instrument name: ARKRAY's ADAMS Lite HA 8380V, JAPAN.

Test Interpretation:

HbA1C is formed by the condensation of glucose with n-terminal valine residue of each beta chain of HbA to form an unstable schiff base. It is the major fraction, constituting approximately 80% of HbA1c. Formation of glycated hemoglobin (GHb) is essentially irreversible and the concentration in the blood depends on both the lifespan of the red blood cells (RBC) (120 days) and the blood glucose concentration. The GHb concentration represents the integrated values for glucose overthe period of 6 to 8 weeks. GHb values are free of day to day glucose fluctuations and are unaffected by recent exercise or food ingestion. Concentration of plasmaglucose concentration in GHb depends on the time interval, with more recent values providing a larger contribution than earlier values. The interpretation of GHbdepends on RBC having a normal life span. Patients with hemolytic disease or other conditions with shortened RBC survival exhibit a substantial reduction of GHb. High GHb have been reported in iron deficiency anemia. GHb has been firmly established as an index of long term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. The absolute risk of retinopathy and nephropathy are directly proportional to themean of HbA1C. Genetic variants (e.g. HbS trait, HbC trait), elevated HbF and chemically modified derivatives of hemoglobin can affect the accuracy of HbA1c measurements. The effects vary depending on the specific Hb vatiant or derivative and the specific HbA1c method.

Ref by ADA 2020

MEAN PLASMA GLUCOSE

Method:- Calculated Parameter

134 H

mg/dL

Non Diabetic < 100 mg/dL Prediabetic 100- 125 mg/dL Diabetic 126 mg/dL or Higher

BANWARI Technologist

Page No: 10 of 12





Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 10/09/2022 10:04:51 Date NAME :- Mr. LOKESH MEENA

Sex / Age :- Male 32 Yrs

Company :- MediWheel

Patient ID :-12222285

Ref. By Dr:- BOB

Lab/Hosp:-

Sample Type :- URINE

Sample Collected Time 10/09/2022 10:21:30

Final Authentication: 10/09/2022 12:23:50

#### **CLINICAL PATHOLOGY**

Test Name	Value	Unit	Biological Ref Interval
Urine Routine			
PHYSICAL EXAMINATION			
COLOUR	PALE YEL	LOW	PALE YELLOW
APPEARANCE	Clear		Clear
CHEMICAL EXAMINATION			
REACTION(PH)	5.5		5.0 - 7.5
SPECIFIC GRAVITY	1.025		1.010 - 1.030
PROTEIN	NIL		NIL
SUGAR	NIL		NIL
BILIRUBIN	NEGATIV	E	NEGATIVE
UROBILINOGEN	NORMAL		NORMAL
KETONES	NEGATIV	E	NEGATIVE
NITRITE	NEGATIV	E	NEGATIVE
MICROSCOPY EXAMINATION			
RBC/HPF	NIL	/HPF	NIL
WBC/HPF	2-3	/HPF	2-3
EPITHELIAL CELLS	1-2	/HPF	2-3
CRYSTALS/HPF	ABSENT		ABSENT
CAST/HPF	ABSENT		ABSENT
AMORPHOUS SEDIMENT	ABSENT		ABSENT
BACTERIAL FLORA	ABSENT		ABSENT
YEAST CELL	ABSENT		ABSENT
OTHER	ABSENT		

SURENDRAMEENA **Technologist** 

Page No: 11 of 12





Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgovalspathlab.com | E-mail: drgovalpiyush@gmail.com



:- 10/09/2022 10:04:51 Date NAME :- Mr. LOKESH MEENA Patient ID: -12222285 Ref. By Dr:- BOB

Sex / Age :- Male

Sample Type :- PLAIN/SERUM

32 Yrs

Lab/Hosp:-

Company :- MediWheel

Sample Collected Time 10/09/2022 10:21:30

Final Authentication: 10/09/2022 12:49:15

#### **IMMUNOASSAY**

Test Name	Value	Unit	Biological Ref Interval
TOTAL THYROID PROFILE			
SERUM TOTAL T3 Method:- Chemiluminescence(Competitive immunoassay)	1.280	ng/ml	0.600 - 1.810
SERUM TOTAL T4 Method:- Chemiluminescence(Competitive immunoassay)	7.860	ug/dl	4.500 - 10.900
SERUM TSH ULTRA Method:- Enhanced Chemiluminescence Immunoassay	2.410	μIU/mL	0.550 - 4.780

Interpretation: Triiodothyronine (T3) contributes to the maintenance of the euthyroid state. A decrease in T3 concentration of up to 50% occurs in a variety of clinical situations, including acute and chronic disease. Although T3 results alone cannot be used to diagnose hypothyroidism, T3 concentration may be more sensitive than thyroxine (T4) for hyperthyroidism. Consequently, the total T3 assay can be used in conjunction with other assays to aid in the differential diagnosis of thyroid disease. T3 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, Free T3 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake, or T4 uptake can be used with the total T3 result to calculate the free T3 index and estimate the concentration of free T3.

Interpretation: The measurement of Total T4 aids in the differential diagnosis of thyroid disease. While >99.9% of T4 is protein-bound, primarily to thyroxine-binding globulin (TBG), it is the free fraction that is biologically active. In most patients, the total T4 concentration is a good indicator of thyroid status. T4 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, free T4 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake may be used with the total T4 result to calculate the free T4 index (FT4I) and estimate the concentration of free T4. Some drugs and some nonthyroidal patient conditions are known to alter TT4 concentrations in vivo.

Interpretation: TSH stimulates the production of thyroxine (T4) and trilodothyronine (T3) by the thyroid gland. The diagnosis of overt hypothyroidism by the finding of a low total T4 or free T4 concentration is readily confirmed by a raised TSH concentration. Measurement of low or undetectable TSH concentrations may assist the diagnosis of hyperthyroidism, where concentrations of T4 and T3 are elevated and TSH secretion is suppressed. These have the advantage of discriminating between the concentrations of TSH observed in thyrotoxicosis, compared with the low, but detectable, concentrations that occur in subclinical hyperthyroidism. The performance of this assay has not been established for neonatal specimens. Some drugs and some nonthyroidal patient conditions are known to alter TSH concentrations in vivo.

#### INTERPRETATION

PREGNANCY	REFERENCE RANGE FOR TSH IN uIU/mL (As per American Thyroid Association)
1st Trimester	0.10-2.50
2nd Trimester	0.20-3.00
3rd Trimester	0.30-3.00

\*\*\* End of Report \*\*\*

NARENDRAKUMAR **Technologist** 

Page No: 12 of 12





## Dr. Goya Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 10/09/2022 10:04:51

NAME :- Mr. LOKESH MEENA

Sex / Age :- Male

Company :- MediWheel

32 Yrs

Patient ID :-12222285 Ref. By Doctor:-BOB

Lab/Hosp:-

Final Authentication: 10/09/2022 13:04:09

**BOB PACKAGE BELOW 40MALE** 

#### X RAY CHEST PA VIEW:

Both lung fields appears clear.

Bronchovascular markings appear normal.

Trachea is in midline.

Both the hilar shadows are normal.

Both the C.P. angles is clear.

Both the domes of diaphragm are normally placed.

Bony cage and soft tissue shadows are normal.

Heart shadows appear normal.

Impression :- Normal Study

(Please correlate clinically and with relevant further investigations)

\*\*\* End of Report \*\*\*

Dr. Piyush Goyal (D.M.R.D.) BILAL

Page No: 1 of 1

Dr. Piyush Goyal M.B.B.S., D.M.R.D. RMC Reg No. 017996

Dr. Poonam Supta MBBS, MD (Radio Diagnosis) RMO No. 32495

Dr. Ashish Choudhary

MBBS, MD (Radio Diagnosis) Fetal Medicine Consultant FMF ID - 260517 | RMC No 22430 Dr. Rathod Hetali Amrutlal MBBS, M.D. (Radio-Diagnosis) RMC No. 17163

Transcript by.



## Dr. Goyal's

### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Tele: 0141-2293346, 4049787, 9887049787 Website: www.drgoyalspathlab.com | E-mail: dago

:- 10/09/2022 10:04:51 NAME :- Mr. LOKESH MEENA

Sex / Age :- Male 32 Yrs Company :- MediWheel

Patient ID: -12222285 Ref. By Doctor:-BOB

Lab/Hosp:-



Final Authentication: 10/09/2022 14:43:52

#### **BOB PACKAGE BELOW 40MALE** 2D ECHO OPTION TMT (ADULT/CHILD)

#### 2D-ECHOCARDIOGRAPHY M.MODE WITH DOPPLER STUDY:

			_FAIR TRANST	HORACIC ECHOCAR	IDIOGRAPHICV	VINDOW MORE	HOLOGY:		
MITRAL VA	ALVE	NORMAL		TRICUS	TRICUSPID VALVE			NORMAL	
AORTIC VA	ALVE	NOR	MAL	PULMO	NARY VALVE		NORMA	1	
		M.MODE	EXAMITATION	۷:			,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,		
AO	27	mm	LA	32	Mm	IVS-D	6	mm	

				61111				_
.VEF	62%			RWMA	•	ABSENT		
RVWT		mm.	EDV		МІ	LVVS		ml
LVPW-D	6	mm	LVPW-S	14	Mm	RV		mm
IVS-S	11	mm	LVID	40	Mm	LVSD	26	mm
AO	27	mm	LA	32	Mm	IVS-D	6	mm

		CHA	AMBERS:	
LA	NORMAL	RA	NORMAL	
LV	NORMAL	RV	NORMAL	
PERICARDIU	JM	NORMAL		
		COLOU	D DODDIED	

				COL	OUR DOPPLER:			
	MI	TRAL VAL	VE					
E VELOCITY	0.99	m/sec	PEAK	PEAK GRADIENT		Mn	n/hg	
A VELOCITY	0.56	m/sec	MEA	N GRADIEN	Т		Mm/hg	
MVA BY PHT		Cm2	MVA	BY PLANIM	ETRY		Cm2	
MITRAL REGURGITAT	ION		+:		ABSENT			
	AC	RTIC VAL	VE					
PEAK VELOCITY	1.27	m	/sec	PEAK GE	PEAK GRADIENT		mm/hg	
AR VMAX		m	/sec	ec MEAN GRADIENT		mı	mm/hg	
AORTIC REGURGITAT	ION			ABSENT				
	TRIC	USPID VA	LVE					
PEAK VELOCITY	0.60	)	m/sec	PEAK GRADIENT			mm/hg	
MEAN VELOCITY			m/sec	MEAN GRADIENT			mm/hg	
VMax VELOCITY								
TRICUSPID REGURGIT	TATION			ABSENT				
	PU	MONARY	VALVE					
PEAK VELOCITY		0.9		M/sec.	PEAK GRADIENT		Mm/hg	
MEAN VALOCITY					MEAN GRADIENT		Mm/hg	
PULMONARY REGUR	GITATION				ABSENT			

Page No: 1 of 2

ANITASHARMA





## Dr. Goyal's Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanga Tele: 0141-2293346, 4049787, 9887049787 Website: www.drgoyalspathlab.com | E-mail: dr

Date

:- 10/09/2022 10:04:51

NAME :- Mr. LOKESH MEENA

32 Yrs

Sex / Age :- Male

Company :- MediWheel

Patient ID: -12222285

Ref. By Doctor:-BOB

Lab/Hosp:-



Final Authentication: 10/09/2022 14:43:52

#### Impression--

- 1. Normal LV size & contractility
- LVEF 70 %. 2. No RWMA,
- 3. Normal cardiac chamber.
- 4. Normal valve
- 5. No clot, no vegetation, no pericardial effusion.

\*\*\* End of Report \*\*\*

Page No: 2 of 2

**ANITASHARMA** 





### Dr. Goyal Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 10/09/2022 10:04:51

NAME

:- Mr. LOKESH MEENA

Sex / Age :- Male

32 Yrs

Company :- MediWheel

Patient ID :-12222285 Ref. By Doctor:-BOB

Lab/Hosp :-

Final Authentication: 10/09/2022 13:44:14

**BOB PACKAGE BELOW 40MALE** 

#### **USG WHOLE ABDOMEN**

Liver is enlarged in size 16 cm. Echo-texture is bright. No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is of normal size. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal: No focal lesion is seen within pancreas.

Spleen is of normal size and shape. Echotexture is normal. No focal lesion is seen.

Kidneys are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. No focal lesion is seen. Collecting system does not show any dilatation or calculus.

Urinary bladder is partially filled.

Prostate is normal in size with normal echo-texture and outline.

No significant free fluid is seen in peritoneal cavity.

.RIF / LIF shows gas filled bowel loops.

#### IMPRESSION: .

Hepatomegaly with Grade I Fatty liver Needs clinical correlation for further evaluation

\*\*\* End of Report \*\*\*

Page No: 1 of 1

ANITASHARMA

Dr. Piyush Goyal . M.B.B.S., D.M.R.D. RMC Reg No. 017996

Dr. Poonam Gupta MBBS, MD (Radio Diagnosis) RMC No. 32495

Dr. Ashish Choudhary MBBS, MD (Radio Diagnosis) Fetal Medicine Consultant

MBBS, M.D. (Radio-Diagnosis) RMC No. 17163 FMF ID - 260517 | RMC No 22430

Dr. Rathod Hetali Amrutlal

Transcript by.