Printed On

 Age / Sex
 : 41 Year(s) / Male
 Report On
 : 25/05/2024 6:01 PM

Type : OP

PID No.

Ref. Dr : MediWheel



InvestigationObservedUnitBiologicalValueReference Interval

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING

 $({\rm EDTA~Blood} Agglutination)$

Remark: Test to be confirmed by gel method.

'O' 'Positive'





: 28/05/2024 6:44 PM



PID No. : MED111936776 : 712415793 SID No. Collection On : 25/05/2024 11:38 AM Age / Sex : 41 Year(s) / Male Report On

Printed On

Type : OP

RDW-SD

Neutrophils

Lymphocytes

Total WBC Count (TC)

 $({\rm EDTA}\ Blood \textit{Derived from Impedance})$

(Blood/Impedance Variation & Flow Cytometry)

(Blood/Impedance Variation & Flow Cytometry)

(Derived)

Ref. Dr : MediWheel



: 25/05/2024 6:01 PM

: 28/05/2024 6:44 PM

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HAEMATOLOGY			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood'Spectrophotometry)	15.7	g/dL	13.5 - 18.0
INTERPRETATION: Haemoglobin values vary in Mer blood loss, renal failure etc. Higher values are often due			
PCV (Packed Cell Volume) / Haematocrit (EDTA Blood/Derived)	45.8	%	42 - 52
RBC Count (EDTA Blood/Automated Blood cell Counter)	5.56	mill/cu.mm	4.7 - 6.0
MCV (Mean Corpuscular Volume) (EDTA Blood/Derived from Impedance)	82.0	fL	78 - 100
MCH (Mean Corpuscular Haemoglobin) (EDTA Blood/Derived)	28.2	pg	27 - 32
MCHC (Mean Corpuscular Haemoglobin concentration) (EDTA Blood/Derived)	34.3	g/dL	32 - 36
RDW-CV (Derived)	12.7	%	11.5 - 16.0

36.45

8150

56

34





fL

%

%

cells/cu.mm



39 - 46

4000 - 11000

40 - 75

20 - 45

PID No. : MED111936776

: 712415793 SID No. Age / Sex : 41 Year(s) / Male

Type : OP

Ref. Dr : MediWheel Register On : 25/05/2024 8:57 AM

Collection On : 25/05/2024 11:38 AM

: 25/05/2024 6:01 PM

Report On

: 28/05/2024 6:44 PM **Printed On**



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Eosinophils (Blood/Impedance Variation & Flow Cytometry)	04	%	01 - 06
Monocytes (Blood/Impedance Variation & Flow Cytometry)	06	%	01 - 10
Basophils (Blood/Impedance Variation & Flow Cytometry)	00	%	00 - 02
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	4.56	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	2.77	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.33	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.49	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.00	10^3 / μl	< 0.2
Platelet Count (EDTA Blood/Derived from Impedance)	196	10^3 / μl	150 - 450
MPV (Blood/ <i>Derived</i>)	12.0	fL	7.9 - 13.7
PCT	0.24	%	0.18 - 0.28

10



ESR (Erythrocyte Sedimentation Rate)

(Citrated Blood/Automated ESR analyser)



mm/hr



< 15

 PID No.
 : MED111936776
 Register On
 : 25/05/2024 8:57 AM

 SID No.
 : 712415793
 Collection On
 : 25/05/2024 11:38 AM

 Age / Sex
 : 41 Year(s) / Male
 Report On
 : 25/05/2024 6:01 PM

Type : OP Printed On : 28/05/2024 6:44 PM

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulfanilic Acid)	0.5	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.2	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.30	mg/dL	0.1 - 1.0
Total Protein (Serum/Biuret)	7.3	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.9	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	2.40	gm/dL	2.3 - 3.6
A: G Ratio (Serum/Derived)	2.04		1.1 - 2.2
INTERPRETATION: Remark : Electrophoresis is the	e preferred method		
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC / Kinetic)	41	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC / Kinetic)	54	U/L	5 - 41
Alkaline Phosphatase (SAP) (Serum/PNPP / Kinetic)	96	U/L	53 - 128
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	37	U/L	< 55







 PID No.
 : MED111936776
 Register On
 : 25/05/2024 8:57 AM

 SID No.
 : 712415793
 Collection On
 : 25/05/2024 11:38 AM

 Age / Sex
 : 41 Year(s) / Male
 Report On
 : 25/05/2024 6:01 PM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/Oxidase / Peroxidase method)	223	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol phosphate oxidase / peroxidase)	207	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

: 28/05/2024 6:44 PM

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual_circulating level of triglycerides during most part of the day.

1			
HDL Cholesterol (Serum/Immunoinhibition)	35	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	146.6	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	41.4	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	188.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220







PID No. : MED111936776 Register On : 25/05/2024 8:57 AM : 712415793 SID No. Collection On : 25/05/2024 11:38 AM

Age / Sex : 41 Year(s) / Male Report On : 25/05/2024 6:01 PM

Printed On

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	Observed Unit	<u>Biological</u>
	<u>Value</u>	Reference Interval

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

: 28/05/2024 6:44 PM

Total Cholesterol/HDL Cholesterol Ratio (Serum/Calculated)	6.4	Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
Triglyceride/HDL Cholesterol Ratio	5.9	Optimal: < 2.5

Triglyceride/HDL Cholesterol Ratio Mild to moderate risk: 2.5 - 5.0 (TG/HDL) High Risk: > 5.0

(Serum/Calculated)

LDL/HDL Cholesterol Ratio 4.2 Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 (Serum/Calculated) High Risk: > 6.0









 PID No.
 : MED111936776
 Register On
 : 25/05/2024 8:57 AM

 SID No.
 : 712415793
 Collection On
 : 25/05/2024 11:38 AM

Printed On

Age / Sex : 41 Year(s) / Male Report On : 25/05/2024 6:01 PM

Type : OP

Ref. Dr : MediWheel

medall

Investigation Glycosylated Haemoglobin (HbA1c)	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HbA1C (Whole Blood/HPLC)	7.2	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

: 28/05/2024 6:44 PM

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 159.94 mg/dl

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.







: MediWheel

PID No. : MED111936776 Register On : 25/05/2024 8:57 AM : 712415793 SID No. Collection On : 25/05/2024 11:38 AM

Report On

Age / Sex : 41 Year(s) / Male

Type : OP

Ref. Dr

25/05/2024 6:01 PM

: 28/05/2024 6:44 PM **Printed On**



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	7.9		
Glucose Fasting (FBS) (Plasma - F/GOD- POD)	111	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Nil Urine sugar, Fasting Nil

(Urine - F)

Glucose Postprandial (PPBS) 259 mg/dL 70 - 140

(Plasma - PP/GOD - POD)

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Sugar (PP-2 hours) (Urine - PP)	Positive (+++)		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	7.9	mg/dL	7.0 - 21
Creatinine (Serum/Laffe Kinetic)	1.0	mg/dL	0.9 - 1.3

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetylcyteine, chemotherapeutic agent such as flucytosine etc.

Uric Acid 5.4 mg/dL 3.5 - 7.2

(Serum/Uricase/Peroxidase)







Report On

Type : OP Printed On : 28/05/2024 6:44 PM

Ref. Dr : MediWheel

Age / Sex : 41 Year(s) / Male



<u>Investigation</u>	<u>Observed</u> <u>l</u>	<u>Jnit</u>	<u>Biological</u>
-	<u>Value</u>		Reference Interval

: 25/05/2024 6:01 PM

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.23 ng/ml 0.7 - 2.04

(Serum/Chemiluminescent Immunometric Assay (CLIA))

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Thyroxine) - Total 11.69 Microg/dl 4.2 - 12.0

(Serum/Chemiluminescent Immunometric Assay

(CLIA))

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 1.403 µIU/mL 0.35 - 5.50

(Serum/Chemiluminescent Immunometric Assay (CLIA))

INTERPRETATION: Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.







PID No. : MED111936776

SID No. : 712415793

Age / Sex : 41 Year(s) / Male

Type : OP

Ref. Dr : MediWheel

Register On : 25/05/2024 8:57 AM

Collection On : 25/05/2024 11:38 AM

Report On : 25/05/2024 6:01 PM

Printed On : 28/05/2024 6:44 PM



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>IMMUNOASSAY</u>			
Total PSA (Serum/Chemiluminescent Immunometric Assay (CLIA))	0.89	ng/ml	Normal: 0.0 - 4.0 Inflammatory & Non Malignant conditions of Prostate & genitourinary system: 4.01 - 10.0 Suspicious of Malignant disease of Prostate: > 10.0

INTERPRETATION: REMARK: PSA alone should not be used as an absolute indicator of malignancy.







 PID No.
 : MED111936776
 Register On
 : 25/05/2024 8:57 AM

 SID No.
 : 712415793
 Collection On
 : 25/05/2024 11:38 AM

 Age / Sex
 : 41 Year(s) / Male
 Report On
 : 25/05/2024 6:01 PM

Type : OP Printed On : 28/05/2024 6:44 PM

Ref. Dr : MediWheel

Glucose

(Urine)

Ketone

Blood (Urine)



Investigation	<u>Observed</u> <u>Unit</u> <u>Value</u>	<u>Biological</u> Reference Interval
CLINICAL PATHOLOGY	 -	
PHYSICAL EXAMINATION		
Colour (Urine/Physical examination)	Pale yellow	Yellow to Amber
Volume (Urine/Physical examination)	30	ml
Appearance (Urine)	Clear	
CHEMICAL EXAMINATION		
pH (Urine)	6.0	4.5 - 8.0
Specific Gravity (Urine/Dip Stick - Reagent strip method)	1.005	1.002 - 1.035
Protein (Urine/Dip Stick - Reagent strip method)	Negative	Negative

Nil

Nil

Nil

(Urine/Dip Stick Reagent strip method)			
Leukocytes (Urine)	Negative	leuco/uL	Negative
Nitrite (Urine/Dip Stick - Reagent strip method)	Nil		Nil
Bilirubin (Urine)	Negative	mg/dL	Negative







Nil

Nil

Nil

 PID No.
 : MED111936776
 Register On
 : 25/05/2024 8:57 AM

 SID No.
 : 712415793
 Collection On
 : 25/05/2024 11:38 AM

Age / Sex : 41 Year(s) / Male **Report On** : 25/05/2024 6:01 PM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Urobilinogen (Urine/Dip Stick - Reagent strip method)	Normal		Within normal limits
<u>Urine Microscopy Pictures</u>			
RBCs (Urine/Microscopy)	Nil	/hpf	NIL
Pus Cells (Urine/Microscopy)	1-2	/hpf	< 5
Epithelial Cells (Urine/Microscopy)	2-3	/hpf	No ranges
Others (Urine)	Nil		Nil

: 28/05/2024 6:44 PM







PID No. : MED111936776 Register On : 25/05/2024 8:57 AM : 712415793 SID No. Collection On : 25/05/2024 11:38 AM Age / Sex : 41 Year(s) / Male Report On : 25/05/2024 6:01 PM

: 28/05/2024 6:44 PM **Printed On**

: OP

Type

Ref. Dr : MediWheel



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Stool Analysis - ROUTINE			
Colour (Stool)	Brown		Brown
Blood (Stool)	Not present		Not present
Mucus (Stool)	Not present		Not present
Reaction (Stool)	Alkaline		Alkaline
Consistency (Stool)	Semi solid		Semi solid
Ova (Stool)	Nil		Nil
Others (Stool)	Nil		Nil
Cysts (Stool)	Nil		Nil
Trophozoites (Stool)	Nil		Nil
RBCs (Stool)	Nil	/hpf	Nil
Pus Cells (Stool)	1-2	/hpf	Nil
Macrophages (Stool)	Nil		Nil
Epithelial Cells (Stool)	Nil	/hpf	Nil







APPROVED BY

-- End of Report --



Name	Mr.GOVINDA NAYAK	ID	MED111936776
Age & Gender	41/MALE	Visit Date	25/05/2024
Ref Doctor Name	MediWheel		

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in size and shows slightly increased echotexture.

No evidence of focal lesion or intrahepatic biliary ductal dilatation.

Hepatic and portal vein radicals are normal.

GALL BLADDER show normal shape and has clear contents.

Gall bladder wall is of normal thickness. CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern.

No evidence of ductal dilatation or calcification.

SPLEEN show normal shape, size and echopattern.

No demonstrable Para-aortic lymphadenopathy.

KIDNEYS move well with respiration and have normal shape, size and echopattern.

Cortico- medullary differentiations are well madeout.

No evidence of calculus or hydronephrosis.

	Bipolar length (cms)	Parenchymal thickness	
		(cms)	
Right Kidney	9.4	1.4	
Left Kidney	10.0	1.5	

URINARY BLADDER show normal shape and wall thickness.

It has clear contents. No evidence of diverticula.

PROSTATE is normal size.

No evidence of ascites.

IMPRESSION:

> GRADE I FATTY CHANGES IN LIVER.

CONSULTANT RADIOLOGISTS

REPORT DISCLAIMER

- 1.This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11.Disputes, if any, with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



Name	Mr.GOVINDA NAYAK	ID	MED111936776
Age & Gender	41/MALE	Visit Date	25/05/2024
Ref Doctor Name	MediWheel		

DR. ANITHA ADARSH

DR. MOHAN B

AA/MS

REPORT DISCLAIMER

- 1.This is only a radiologincal imperssion.Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11.Disputes,if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



Name	Mr. GOVINDA NAYAK	ID	MED111936776
Age & Gender	41Y/M	Visit Date	May 25 2024 8:57AM
Ref Doctor	MediWheel		

X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

Impression: No significant abnormality detected.

Dr. Anitha Adarsh Consultant Radiologist