

YOUR CLINICAL TEST RESULTS

Mr. SUKHENDU HALDER

MALE / 43 Yrs / AGMC.0000387882 / AGMCAH47476

MEDI WHEEL FULL BODY HCK - MALE (ABOVE 40 YRS WITH TMT/ECHO)

Date : 25/05/2024

COMPLETE HAEMOGRAM PROFILE

Test Name	Result	Unit	Level	Range
Hemoglobin	12.9 *	g/dl	●	13.0-17.0
RBC COUNT	4.45 *	Million/ul	●	4.5-5.5
Hematocrit - Hct:	39.2 *	%	●	41-53
MCV	88.0	fl	●	83-101
MCH	29.0	pg	●	27-32
MCHC 32.9 % ●	31.5-34.5	RDW 14.0 % ●		11.8-14.0
WBC Count	8500	/cu mm	●	4000-10000
Platelet Count	1.56	lacs/cum m	●	1.5-4.0
Neutrophils	57	%	●	40-80
Lymphocytes	33	%	●	20-40
Monocytes	06	%	●	2-10
Eosinophils	04	%	●	01-06
Basophils	00	%	●	0-0
RBC:	Normocytic Normochromic cells			
Platelets:	Adequate.			
ERYTHROCYTE SEDIMENTATION RATE (ESR)	15	mm/1st hr	●	0-15

URINE ROUTINE AND MICROSCOPY

Test Name	Result	Unit	Level	Range
Volume:	40	mL		
Colour:	Pale straw			
Appearance	Slightly Turbid			
Specific Gravity 1.015 * ●	1.016-1.025	Reaction: Acidic		
Protein :	Absent			
Sugar:	Absent			
Ketone	Absent			



Within Normal Range



Boderline High/Low



Outside Range

YOUR CLINICAL TEST RESULTS

Mr. SUKHENDU HALDER

MALE / 43 Yrs / AGMC.0000387882 / AGMCAH47476

MEDI WHEEL FULL BODY HCK - MALE (ABOVE 40 YRS WITH TMT/ECHO)

Date : 25/05/2024

Bile Pigments:	Absent		
Blood:	Absent		
RBC	Nil	/hpf	0-2
Pus Cells	2-4	/hpf	
Epithelial Cells	2-4 /h.p.f	/hpf	
Yeast Cells	Absent		
Bacteria:	Absent		
Casts:	Not Found		
Crystals:	Found		
Ca-oxalate	+	/lpf	

URINE SUGAR - POST PRANDIAL (QUALITATIVE)

Test Name	Result	Unit	Level	Range
URINE GLUCOSE(POST PRANDIAL)	Trace			

URINE SUGAR- FASTING(QUALITATIVE)

Test Name	Result	Unit	Level	Range
URINE GLUCOSE(FASTING)	Nil			

BLOOD GROUPING AND TYPING (ABO AND RH)

Test Name	Result	Unit	Level	Range
ABO Group:	A			
Rh (D) Type:	POSITIVE			

LIVER FUNCTION TEST (PACKAGE)

Test Name	Result	Unit	Level	Range
ALT(SGPT) - SERUM	91 *	U/L	●	0-50
ALBUMIN - SERUM	4.6	g/dL	●	3.5-5.1



Within Normal Range



Boderline High/Low



Outside Range

YOUR CLINICAL TEST RESULTS

Mr. SUKHENDU HALDER

MALE / 43 Yrs / AGMC.0000387882 / AGMCAH47476

MEDI WHEEL FULL BODY HCK - MALE (ABOVE 40 YRS WITH TMT/ECHO)

Date : 25/05/2024

ALKALINE PHOSPHATASE - SERUM	55	U/L	●	43-115
AST (SGOT) - SERUM	61 *	U/L	●	0-50
BILIRUBIN TOTAL - SERUM	0.8	mg/dL	●	0.3-1.2

LIPID PROFILE TEST (PACKAGE)

Test Name	Result	Unit	Level	Range
CHOLESTEROL - SERUM	163	mg/dL	●	0-200

CREATININE - SERUM

Test Name	Result	Unit	Level	Range
CREATININE - SERUM	1.2	mg/dL	●	0.9-1.3

LIVER FUNCTION TEST (PACKAGE)

Test Name	Result	Unit	Level	Range
GGTP: GAMMA GLUTAMYL TRANSPEPTIDASE - SERUM	42	U/L	●	0-55

GLUCOSE - PLASMA (FASTING)

Test Name	Result	Unit	Level	Range
GLUCOSE - PLASMA (FASTING)	111 *	mg/dL	●	70-99

GLUCOSE - PLASMA (POST PRANDIAL)

Test Name	Result	Unit	Level	Range
GLUCOSE - PLASMA (POST PRANDIAL)	171 *	mg/dL	●	70-140

HBA1C (GLYCOSYLATED HAEMOGLOBIN)-WHOLE BLOOD

Test Name	Result	Unit	Level	Range
HBA1C (GLYCOSYLATED HAEMOGLOBIN)-WHOLE BLOOD	5.8 *	%	●	Nondiabetic : 4 - 5.6 % Prediabetics : 5.7 - 6.4%

Diabetes : $\geq 6.5\%$

ADA Therapeutic goal : $<7\%$

LIPID PROFILE TEST (PACKAGE)

Test Name	Result	Unit	Level	Range
	Within Normal Range ●		Boderline High/Low ●	Outside Range

YOUR CLINICAL TEST RESULTS

Mr. SUKHENDU HALDER

MALE / 43 Yrs / AGMC.0000387882 / AGMCAH47476

MEDI WHEEL FULL BODY HCK - MALE (ABOVE 40 YRS WITH TMT/ECHO)

Date : 25/05/2024

HDL CHOLESTEROL - SERUM	48	mg/dL	●	30-70
LDL CHOLESTEROL -SERUM	99	mg/dL	●	Optimal: <100
VLDL CHOLESTEROL - SERUM (Calculated)	16	mg/dL	●	0-35

LIVER FUNCTION TEST (PACKAGE)

Test Name	Result	Unit	Level	Range
PROTEIN TOTAL - SERUM	7.7	g/dL	●	6.4-8.3
GLOBULIN: (CALCULATED) - SERUM	3.1	g/dL	●	1.8-3.6

THYROID PROFILE - I(T3,T4 AND TSH)

Test Name	Result	Unit	Level	Range
TOTAL T3: TRI IODOTHYRONINE - SERUM	1.37	ng/ml	●	0.87-1.78
TOTAL T4: THYROXINE - SERUM	10.65	µg/dL	●	5.48-14.28

LIPID PROFILE TEST (PACKAGE)

Test Name	Result	Unit	Level	Range
TRIGLYCERIDES - SERUM	108	mg/dL	●	0-150

THYROID PROFILE - I(T3,T4 AND TSH)

Test Name	Result	Unit	Level	Range
TSH: THYROID STIMULATING HORMONE - SERUM	3.12	µIU/mL	●	0.38-5.33

URIC ACID - SERUM

Test Name	Result	Unit	Level	Range
URIC ACID - SERUM	7.2	mg/dL	●	2.6-7.2

LIPID PROFILE TEST (PACKAGE)

Test Name	Result	Unit	Level	Range
VLDL CHOLESTEROL - SERUM (Calculated)	16	mg/dL	●	0-35

LIVER FUNCTION TEST (PACKAGE)

Test Name	Result	Unit	Level	Range
	Within Normal Range		Boderline High/Low	Outside Range

YOUR CLINICAL TEST RESULTS

Mr. SUKHENDU HALDER

MALE / 43 Yrs / AGMC.0000387882 / AGMCAH47476

MEDI WHEEL FULL BODY HCK - MALE (ABOVE 40 YRS WITH TMT/ECHO)

Date : 25/05/2024

BILIRUBIN CONJUGATED (DIRECT) - SERUM	0.2	mg/dL	●	0.0-0.2
---------------------------------------	-----	-------	---	---------

PROSTATIC SPECIFIC ANTIGEN (PSA TOTAL) - SERUM

Test Name	Result	Unit	Level	Range
PROSTATIC SPECIFIC ANTIGEN (PSA TOTAL) - SERUM	0.46	ng/mL	●	0.00-6.50

BUN (BLOOD UREA NITROGEN)

Test Name	Result	Unit	Level	Range
BUN (BLOOD UREA NITROGEN)	12.3	mg/dL	●	7.0-18.0

LIVER FUNCTION TEST (PACKAGE)

Test Name	Result	Unit	Level	Range
A/G - RATIO	1.5		●	1.0-2.0

2D ECHOCARDIOGRAPHY

- Good biventricular systolic function.

ECG

Normal Sinus Rhythm

Normal ECG.

ULTRASOUND SCREENING WHOLE ABDOMEN

Hepatomegaly with grade I fatty liver.

X-RAY CHEST PA

Normal study.



Within Normal Range



Boderline High/Low



Outside Range



Name	: MR.RAJU M	TID/SID	: UMR1577070/ 27661344
Age / Gender	: 40 Years / Male	Registered on	: 25-May-2024 / 09:57 AM
Ref.By	: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS	Collected on	: 25-May-2024 / 10:00 AM
Req.No	: BIL4292613	Reported on	: 25-May-2024 / 15:27 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF CLINICAL PATHOLOGY

Complete Urine Examination (CUE), Urine

Investigation	Observed Value	Biological Reference Intervals
Physical Examination		
Colour Method:Physical	Pale Yellow	Straw to Yellow
Appearance Method:Physical	Clear	Clear
Chemical Examination		
Reaction and pH Method:pH- Methyl red & Bromothymol blue	5.5	4.6-8.0
Specific gravity Method:Bromothymol Blue	1.015	1.003-1.035
Protein Method:Tetrabromophenol blue	Negative	Negative
Glucose Method:Glucose oxidase/Peroxidase	Positive (+++)	Negative
Blood Method:Peroxidase	Positive (Trace)	Negative
Ketones Method:Sodium Nitroprusside	Negative	Negative
Bilirubin Method:Dichloroanilinediazonium	Negative	Negative
Leucocytes Method:3 hydroxy5 phenylpyrrole + diazonium	Positive (+)	Negative
Nitrites Method:Diazonium + 1,2,3,4 tetrahydrobenzo (h) quinolin 3-ol	Negative	Negative
Urobilinogen Method:Dimethyl aminobenzaldehyde	0.2	0.2-1.0 mg/dl
Microscopic Examination		
Pus cells (leukocytes) Method:Microscopy	10-15	2 - 3 /hpf
Epithelial cells Method:Microscopy	0-1	2 - 5 /hpf
RBC (erythrocytes) Method:Microscopy	2-3	Absent
Casts Method:Microscopy	Absent	Occasional hyaline casts may be seen



Name : **MR.RAJU M** TID/SID : UMR1577070/ 27661344
Age / Gender : 40 Years / Male Registered on : 25-May-2024 / 09:57 AM
Ref.By : ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 25-May-2024 / 10:00 AM
Req.No : BIL4292613 Reported on : 25-May-2024 / 15:27 PM
Reference : Arcofemi Health Care Ltd -

TEST REPORT

Crystals Absent Phosphate, oxalate, or urate crystals may be seen
Method:Microscopy
Others Bacteria (+) Nil
Method:Microscopy
Note Kindly correlate clinically

Method: Semi Quantitative test ,For CUE

Reference: Godkar Clinical Diagnosis and Management by Laboratory Methods, First South Asia edition. Product kit literature.

Interpretation:

The complete urinalysis provides a number of measurements which look for abnormalities in the urine. Abnormal results from this test can be indicative of a number of conditions including kidney disease, urinary tract infection or elevated levels of substances which the body is trying to remove through the urine . A urinalysis test can help identify potential health problems even when a person is asymptomatic. All the abnormal results are to be correlated clinically.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Dr.M.G.Satish
Consultant Pathologist





Name : **MR.RAJU M** TID/SID : UMR1577070/ 27661345
Age / Gender : 40 Years / Male Registered on : 25-May-2024 / 09:57 AM
Ref.By : ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 25-May-2024 / 10:00 AM
Req.No : BIL4292613 Reported on : 25-May-2024 / 16:15 PM
Reference : Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF HEMATOPATHOLOGY

Blood Grouping ABO And Rh Typing, EDTA Whole Blood

Parameter	Results
Blood Grouping (ABO)	A
Rh Typing (D)	POSITIVE

Method: Hemagglutination Tube Method by Forward & Reverse Grouping

Reference: Tulip kit literature

Interpretation: The ABO grouping and Rh typing test determines blood type grouping (A,B, AB, O) and the Rh factor (positive or negative). A person's blood type is based on the presence or absence of certain antigens on the surface of their red blood cells and certain antibodies in the plasma. ABO antigens are poorly expressed at birth, increase gradually in strength and become fully expressed around 1 year of age.

Note: Records of previous blood grouping/Rh typing not available. Please verify before transfusion.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Debleena Thakur

Dr Debleena Thakur
Consultant Pathologist





Name	: MR.RAJU M	TID/SID	: UMR1577070/ 27661345
Age / Gender	: 40 Years / Male	Registered on	: 25-May-2024 / 09:57 AM
Ref.By	: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS	Collected on	: 25-May-2024 / 10:00 AM
Req.No	: BIL4292613	Reported on	: 25-May-2024 / 14:33 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF HEMATOPATHOLOGY

Erythrocyte Sedimentation Rate (ESR), Sodium Citrate Whole Blood

Investigation	Observed Value	Biological Reference Intervals
Erythrocyte Sedimentation Rate	20	<=15 mm/hour
Method:Microphotometrical capillary using stopped flow kinetic analysis		

Complete Blood Count (CBC), EDTA Whole Blood

Investigation	Observed Value	Biological Reference Interval
Hemoglobin	16.3	13.0-18.0 g/dL
Method:Spectrophotometry		
Packed Cell Volume	49.4	40-54 %
Method:Derived from Impedance		
Red Blood Cell Count.	5.64	4.3-6.0 Mill/Cumm
Method:Impedance Variation		
Mean Corpuscular Volume	87.5	78-100 fL
Method:Derived from Impedance		
Mean Corpuscular Hemoglobin	28.8	27-32 pg
Method:Derived from Impedance		
Mean Corpuscular Hemoglobin Concentration	32.9	31.5-36 g/dL
Method:Derived from Impedance		
Red Cell Distribution Width - CV	11.2	11.5-16.0 %
Method:Derived from Impedance		
Red Cell Distribution Width - SD	39.5	39-46 fL
Method:Derived from Impedance		
Total WBC Count.	10730	4000-11000 cells/cumm
Method:Impedance Variation		
Neutrophils	67.9	40-75 %
Method:Impedance Variation, Flowcytometry		
Lymphocytes	26.2	20-45 %
Method:Microscopy		
Eosinophils	1.0	01-06 %
Method:Impedance Variation,Method_Desc= Flow Cytometry		
Monocytes	4.0	01-10 %
Method:Impedance Variation, Flowcytometry		



Name	: MR.RAJU M	TID/SID	: UMR1577070/ 27661345
Age / Gender	: 40 Years / Male	Registered on	: 25-May-2024 / 09:57 AM
Ref.By	: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS	Collected on	: 25-May-2024 / 10:00 AM
Req.No	: BIL4292613	Reported on	: 25-May-2024 / 14:33 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

Basophils.	0.9	00-02 %
Method:Impedance Variation,Method_Desc= Flow Cytometry		
Absolute Neutrophils Count.	7286	1500-6600 cells/cumm
Method:Calculated		
Absolute Lymphocyte Count	2811	1500-3500 cells/cumm
Method:Calculated		
Absolute Eosinophils count.	107	40-440 cells/cumm
Method:Calculated		
Absolute Monocytes Count.	429	<1000 cells/cumm
Method:Calculated		
Absolute Basophils count.	97	<200 cells/cumm
Method:Calculated		
Platelet Count.	3.07	1.4-4.4 lakhs/cumm
Method:Impedance Variation		
Mean Platelet Volume.	8.3	7.9-13.7 fL
Method:Derived from Impedance		
Plateletcrit.	0.26	0.18-0.28 %
Method:Derived from Impedance		

Note Kindly correlate clinically

Method: Automated Hematology Analyzer, Microscopy

Reference: Dacie and Lewis Practical Hematology, 12th Edition

Interpretation: A Complete Blood Picture (CBP) is a screening test which can aid in the diagnosis of a variety of conditions and diseases such as anemia, leukemia, bleeding disorders and infections. This test is also useful in monitoring a person's reaction to treatment when a condition which affects blood cells has been diagnosed. All the abnormal results are to be correlated clinically.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Debleena Thakur

Dr Debleena Thakur
Consultant Pathologist





Name	: MR.RAJU M	TID/SID	: UMR1577070/ 27661346
Age / Gender	: 40 Years / Male	Registered on	: 25-May-2024 / 09:57 AM
Ref.By	: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS	Collected on	: 25-May-2024 / 10:00 AM
Req.No	: BIL4292613	Reported on	: 25-May-2024 / 13:20 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF CLINICAL CHEMISTRY I

Blood Urea Nitrogen (BUN), Serum

Investigation	Observed Value	Biological Reference Interval
Blood Urea Nitrogen.	7	6-20 mg/dL
Method:Kinetic, Urease - GLDH, Calculated		

Interpretation: Urea is a waste product formed in the liver when protein is metabolized. Urea is released by the liver into the blood and is carried to the kidneys, where it is filtered out of the blood and released into the urine. Since this is a continuous process, there is usually a small but stable amount of urea nitrogen in the blood. However, when the kidneys cannot filter wastes out of the blood due to disease or damage, then the level of urea in the blood will rise. The blood urea nitrogen (BUN) evaluates kidney function in a wide range of circumstances, to diagnose kidney disease, and to monitor people with acute or chronic kidney dysfunction or failure. It also may be used to evaluate a person's general health status as well.

Reference: Tietz Fundamentals of Clinical Chemistry and Molecular Diagnostics

Creatinine, Serum

Investigation	Observed Value	Biological Reference Interval
Creatinine.	0.70	0.7-1.3 mg/dL
Method:Spectrophotometry, Jaffe - IDMS Traceable		

Interpretation:

Creatinine is a nitrogenous waste product produced by muscles from creatine. Creatinine is majorly filtered from the blood by the kidneys and released into the urine, so serum creatinine levels are usually a good indicator of kidney function. Serum creatinine is more specific and more sensitive indicator of renal function as compared to BUN because it is produced from muscle at a constant rate and its level in blood is not affected by protein catabolism or other exogenous products. It is also not reabsorbed and very little is secreted by tubules making it a reliable marker. Serum creatinine levels are increased in pre renal, renal and post renal azotemia, active acromegaly and gigantism. Decreased serum creatinine levels are seen in pregnancy and increasing age.

Biological reference interval changed; Reference: Tietz Textbook of Clinical Chemistry & Molecular Diagnostics, Fifth Edition.

Bun/Creatinine Ratio, Serum

Investigation	Observed Value
BUN/Creatinine Ratio	10
Method:Calculated	

Reference:

A Manual of Laboratory Diagnostic Tests. Edition 7, Lippincott Williams and Wilkins, By Frances Talaska Fischbach, RN, BSN, MSN, and Marshall Barnett Dunning 111, BS, MS, Ph.D.



PLEASE SCAN QR CODE
TO VERIFY THE REPORT ONLINE



Name	: MR.RAJU M	TID/SID	: UMR1577070/
Age / Gender	: 40 Years / Male	Registered on	: 25-May-2024 / 09:57 AM
Ref.By	: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS	Collected on	:
Req.No	: BIL4292613	Reported on	:
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Kavya SN

Dr.Kavya S N
Consultant Pathologist





Name	: MR.RAJU M	TID/SID	: UMR1577070/ 27661347-F
Age / Gender	: 40 Years / Male	Registered on	: 25-May-2024 / 09:57 AM
Ref.By	: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS	Collected on	: 25-May-2024 / 10:00 AM
Req.No	: BIL4292613	Reported on	: 25-May-2024 / 13:20 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF CLINICAL CHEMISTRY I

Glucose Fasting (FBS), Sodium Fluoride Plasma

Investigation	Observed Value	Biological Reference Interval
Glucose Fasting Method:Hexokinase	266	Normal: 70 -100 mg/dL Impaired FG: 100-125 mg/dL Diabetes mellitus: \geq 126 mg/dL

Interpretation: It measures the Glucose levels in the blood with a prior fasting of 9-12 hours. The test helps screen a symptomatic/ asymptomatic person who is at risk for Diabetes. It is also used for regular monitoring of glucose levels in people with Diabetes.

Reference: American Diabetes Association. Standards of Medical Care in Diabetes-2020.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Kavya SN

Dr.Kavya S N
Consultant Pathologist





Name	: MR.RAJU M	TID/SID	: UMR1577070/ 27661347-P
Age / Gender	: 40 Years / Male	Registered on	: 25-May-2024 / 09:57 AM
Ref.By	: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS	Collected on	: 25-May-2024 / 16:47 PM
Req.No	: BIL4292613	Reported on	: 25-May-2024 / 20:07 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF CLINICAL CHEMISTRY I

Glucose Post Prandial (PPBS), Sodium Fluoride Plasma

Investigation	Observed Value	Biological Reference Interval
Glucose Post Prandial Method:Hexokinase	467	Normal : 90 - 140 mg/dL Impaired PG: 140-199 mg/dL Diabetes mellitus: \geq 200 mg/dL
Note	Kindly correlate clinically	

Interpretation: This test measures the blood sugar levels 2 hours after a normal meal. Abnormally high blood sugars 2 hours after a meal reflect that the body is not producing sufficient insulin which is indicative of Diabetes.

Reference: American Diabetes Association. Standards of Medical Care in Diabetes-2020.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Dr Manjunatha H.K
Consultant Pathologist





Name	: MR.RAJU M	TID/SID	: UMR1577070/ 27661345
Age / Gender	: 40 Years / Male	Registered on	: 25-May-2024 / 09:57 AM
Ref.By	: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS	Collected on	: 25-May-2024 / 10:00 AM
Req.No	: BIL4292613	Reported on	: 25-May-2024 / 17:55 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF CLINICAL CHEMISTRY I

Glycosylated Hemoglobin (HbA1C), EDTA Whole Blood

Investigation	Observed Value	Biological Reference Interval
Glycosylated Hemoglobin (HbA1c) Method:High-Performance Liquid Chromatography	11.9	Non-diabetic: <= 5.6 % Pre-diabetic: 5.7 - 6.4 % Diabetic: >= 6.5 %
Estimated Average Glucose (eAG) Method:High-Performance Liquid Chromatography	295	mg/dL

Interpretation: It is an index of long-term blood glucose concentrations and a measure of the risk for developing microvascular complications in patients with diabetes. Absolute risks of retinopathy and nephropathy are directly proportional to the mean HbA1c concentration. In persons without diabetes, HbA1c is directly related to risk of cardiovascular disease.

In known diabetic patients, HbA1c can be considered as a tool for monitoring the glycemic control.

- Excellent Control - 6 to 7 %,
- Fair to Good Control - 7 to 8 %,
- Unsatisfactory Control - 8 to 10 %
- and Poor Control - More than 10 %.

Reference: American Diabetes Association. Standards of Medical Care in Diabetes-2018.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Kavya SN

Dr.Kavya S N
Consultant Pathologist





Name	: MR.RAJU M	TID/SID	: UMR1577070/ 27661346
Age / Gender	: 40 Years / Male	Registered on	: 25-May-2024 / 09:57 AM
Ref.By	: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS	Collected on	: 25-May-2024 / 10:00 AM
Req.No	: BIL4292613	Reported on	: 25-May-2024 / 13:20 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF CLINICAL CHEMISTRY I

Lipid Profile, Serum

Investigation	Observed Value	Biological Reference Interval
Total Cholesterol Method:Spectrophotometry , CHOD - POD	183	Desirable: < 200 mg/dL Borderline: 200-239 mg/dL High: >= 240 mg/dL
HDL Cholesterol Method:Spectrophotometry , Direct Measurement	35	Optimal : >=60 mg/dL Borderline : 40-59 mg/dL High Risk <40 mg/dL
Non HDL Cholesterol Method:Calculated	148	Optimal : <130 mg/dL Above Optimal : 130-159 mg/dL Borderline : 160-189 mg/dL High Risk : 190-219 mg/dL Very high Risk : >=220 mg/dL
LDL Cholesterol Method:Calculated	113.2	Optimum: <100 mg/dL Near/above optimum: 100-129 mg/dL Borderline: 130-159 mg/dL High: 160-189 mg/dL Very high: >=190 mg/dL
VLDL Cholesterol Method:Calculated	34.80	<30 mg/dL
Total Cholesterol/HDL Ratio Method:Calculated	5.23	Optimal : <3.3 Low Risk : 3.4-4.4 Average Risk : 4.5-7.1 Moderate Risk : 7.2-11.0 High Risk : >11.0
LDL/HDL Ratio Method:Calculated	3.23	Optimal : 0.5-3.0 Borderline : 3.1-6.0 High Risk : >6.0
Triglycerides Method:Spectrophotometry, Enzymatic - GPO/POD	174	Normal:<150 mg/dL Borderline: 150-199 mg/dL High: 200-499 mg/dL Very high: >=500 mg/dL mg/dl #

Interpretation: Lipids are fats and fat-like substances which are important constituents of cells and are rich sources of energy. A lipid profile typically includes total cholesterol, high density lipoproteins (HDL), low density lipoprotein (LDL), chylomicrons, triglycerides, very low density lipoproteins (VLDL), Cholesterol/HDL ratio .The lipid profile is used to assess the risk of developing a heart disease and to monitor its treatment. The results of the lipid profile are evaluated along with other known risk factors associated with heart disease to plan and monitor treatment. Treatment options require clinical correlation.**Reference:** Third Report of the National Cholesterol Education program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III), JAMA 2001.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---



PLEASE SCAN QR CODE
TO VERIFY THE REPORT ONLINE



Name : **MR.RAJU M** TID/SID : UMR1577070/
Age / Gender : 40 Years / Male Registered on : 25-May-2024 / 09:57 AM
Ref.By : ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on :
Req.No : BIL4292613 Reported on :
Reference : Arcofemi Health Care Ltd -

TEST REPORT

Kavya SN

Dr.Kavya S N
Consultant Pathologist





Name	: MR.RAJU M	TID/SID	: UMR1577070/ 27661346
Age / Gender	: 40 Years / Male	Registered on	: 25-May-2024 / 09:57 AM
Ref.By	: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS	Collected on	: 25-May-2024 / 10:00 AM
Req.No	: BIL4292613	Reported on	: 25-May-2024 / 15:37 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF CLINICAL CHEMISTRY I

Liver Function Test (LFT), Serum

Investigation	Observed Value	Biological Reference Interval
Total Bilirubin. Method:Spectrophotometry, Diazo method	0.55	<=1.2 mg/dL
Direct Bilirubin. Method:Spectrophotometry, Diazo method	0.31	<=0.30 mg/dL
Indirect Bilirubin. Method:Calculated	0.24	<=1.0 mg/dL
Alanine Aminotransferase ,(ALT/SGPT) Method: IFCC without pyridoxal phosphate activation	24	<=41 U/L
Aspartate Aminotransferase,(AST/SGOT) Method: IFCC without pyridoxal phosphate activation	10	<=40 U/L
ALP (Alkaline Phosphatase). Method:Spectrophotometry , IFCC	107	40-129 U/L
Gamma GT. Method:Spectrophotometry , IFCC	32	<60 U/L
Total Protein. Method:Spectrophotometry, Biuret	7.6	6.4-8.3 g/dL
Albumin. Method:Spectrophotometry, Bromcresol Green	4.8	3.5-5.2 g/dL
Globulin. Method:Spectrophotometry, Bromcresol Green	2.8	2.0-3.5 g/dL
A/GRatio. Method:Calculated	1.71	1.1-2.5

Interpretation: Liver functions tests help to identify liver disease, its severity, and its type. Generally these tests are performed in combination, are abnormal in liver disease, and the pattern of abnormality is indicative of the nature of liver disease. An isolated abnormality of a single liver function test usually means a non-hepatic cause. If several liver function tests are simultaneously abnormal, then hepatic etiology is likely.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Dr.M.G.Satish
Consultant Pathologist



PLEASE SCAN QR CODE
TO VERIFY THE REPORT ONLINE



Name	: MR.RAJU M	TID/SID	: UMR1577070/ 27661346
Age / Gender	: 40 Years / Male	Registered on	: 25-May-2024 / 09:57 AM
Ref.By	: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS	Collected on	: 25-May-2024 / 10:00 AM
Req.No	: BIL4292613	Reported on	: 25-May-2024 / 14:45 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF CLINICAL CHEMISTRY I

Prostate Specific Antigen (PSA) Total, Serum

Investigation	Observed Value	Biological Reference Interval
Prostate Specific Antigen (PSA) Total Method:ECLIA	0.702	0.0-4.0 ng/mL

Interpretation: PSA is a protein produced by cells in the prostate and is used to screen men for prostate cancer. PSA levels are elevated in Prostate cancer, and other conditions such as benign prostatic hyperplasia (BPH) and inflammation of the prostate. An elevated PSA may be followed by a biopsy and other tests like urinalysis and ultrasound to rule out urinary tract infections and for an accurate diagnosis. PSA levels are vital to determine the effectiveness of treatment and to detect recurrence in diagnosed cases of prostate cancer.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Dr.M.G.Satish
Consultant Pathologist





Name	: MR.RAJU M	TID/SID	: UMR1577070/ 27661346
Age / Gender	: 40 Years / Male	Registered on	: 25-May-2024 / 09:57 AM
Ref.By	: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS	Collected on	: 25-May-2024 / 10:00 AM
Req.No	: BIL4292613	Reported on	: 25-May-2024 / 14:45 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF CLINICAL CHEMISTRY I

Thyroid Profile (T3,T4,TSH), Serum

Investigation	Observed Value	Biological Reference Interval
Triiodothyronine Total (T3) Method:ECLIA	1.09	0.80-2.00 ng/mL Note: Biological Reference Ranges are changed due to change in method of testing.
Thyroxine Total (T4) Method:ECLIA	8.27	4.6-12.0 µg/dL
Thyroid Stimulating Hormone (TSH) Method:ECLIA	1.29	0.27-4.20 µIU/mL

Interpretation: A thyroid profile is used to evaluate thyroid function and/or help diagnose hypothyroidism and hyperthyroidism due to various thyroid disorders. T4 and T3 are hormones produced by the thyroid gland. They help control the rate at which the body uses energy, and are regulated by a feedback system. TSH from the pituitary gland stimulates the production and release of T4 (primarily) and T3 by the thyroid. Most of the T4 and T3 circulate in the blood bound to protein. A small percentage is free (not bound) and is the biologically active form of the hormones.

Reference: Tietz Fundamentals of Clinical Chemistry and Molecular Diagnostics, Carl A. Burtis, David E. Bruns.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Dr.M.G.Satish
Consultant Pathologist





PLEASE SCAN QR CODE
TO VERIFY THE REPORT ONLINE



Name	: MR.RAJU M	TID/SID	: UMR1577070/ 27661346
Age / Gender	: 40 Years / Male	Registered on	: 25-May-2024 / 09:57 AM
Ref.By	: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS	Collected on	: 25-May-2024 / 10:00 AM
Req.No	: BIL4292613	Reported on	: 25-May-2024 / 13:20 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF CLINICAL CHEMISTRY I

Uric Acid, Serum

Investigation	Observed Value	Biological Reference Interval
Uric Acid. Method:Enzymatic	3.3	3.4-7.0 mg/dL

Interpretation: It is the major product of purine catabolism. Hyperuricemia can result due to increased formation or decreased excretion of uric acid which can be due to several causes like metabolic disorders, psoriasis, tissue hypoxia, pre-eclampsia, alcohol, lead poisoning, acute or chronic kidney disease, etc. Hypouricemia may be seen in severe hepato cellular disease and defective renal tubular reabsorption of uric acid.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Kavya SN

Dr.Kavya S N
Consultant Pathologist





PLEASE SCAN QR CODE

Name	: Mr . RAJU M	TID	: UMR1577070
Age/Gender	: 40 Years/Male	Registered On	: 25-May-2024 09:57 AM
Ref By	: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS	Reported On	: 25-May-2024 04:34 PM
Reg.No	: BIL4292613	Reference	: Arcofemi Health Care Ltd - Medi Whe

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in size and shows diffuse mild fatty changes. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER is moderately distended. Two polyps noted, one in the mid body anteriorly measuring 5 mm and the other in the mid body posteriorly measuring 2mm.

No evidence of intraluminal calculi. Gall bladder wall is of normal thickness.

CBD is of normal calibre.

PANCREAS is normal in size and echopattern. No evidence of ductal dilatation or calcification.

SPLEEN is normal in size and echopattern. It measures 8.0cms in long axis and 2.9cms in short axis.

KIDNEYS move well with respiration and are normal in size and echopattern.

Cortico- medullary differentiations are well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney	9.5	1.2
Left Kidney	11.4	1.4

URINARY BLADDER is moderately distended with normal wall thickness. It has clear contents. No evidence of diverticula.

PROSTATE is normal in size and echopattern. It measures 3.8 x 3.0 x 3.3cms (Vol: 20cc).

No evidence of ascites / pleural effusion / para -aortic lymphadenopathy.

IMPRESSION:

- **GRADE I FATTY LIVER.**
- **GALL BLADDER POLYPS.**

*** End Of Report ***

Dr Meera Krishnan
Consultant Radiologist

25 05 2024 11:31:09
tenet
Indiranagar
Bangalore


Male

QRS
QT / QTcBaz 84 ms
338 / 389 ms
PR 172 ms
P 90 ms
RR / PP 750 / 750 ms
P / QRS / T 44 / 25 / 28 degrees

80 bpm
-- / -- mmHg

Technician:
Ordering Ph:
Referring Ph:
Attending Ph:

NORMAL ECG



Name	MR . RAJU M	Req No. 4292613
Age & Gender	40Y/MALE	Registered on:25.05.2024
Ref Doctor	CREDIT CLIENTS	Reported on:25.05.2024

2D ECHOCARDIOGRAPHY & COLOUR DOPPLER REPORT

M-mode:

	Value	Normal range
LA dimension	3.2	(1.9 – 4.0 cm)
Aorta	3.0	(2.5 – 3.7 cm)
IVS (d)	1.0	(0.6 – 1.1 cm)
LV PW (d)	1.0	(0.6- 1.1 cm)
LVID (d)	4.4	(3.5 – 5.5 cm)
LVID (s)	3.0	(2.4 – 4.2 cm)
EDV	92	ml
ESV	34	ml
LV EF	62%	50 – 70 %

CHAMBERS:

LEFT ATRIUM: Normal

RIGHT ATRIUM: Normal

LEFT VENTRICLE: Normal

RIGHT VENTRICLE: Normal

VALVES:

MITRAL VALVE: Normal

AORTIC VALVE: Normal

TRICUSPID VALVE: Normal

PULMONARY VALVE: Normal

GREAT ARTERIES:

AORTA: Normal

PULMONARY ARTERY: Normal

IAS/IVS: Intact

WALL MOTION ABNORMALITIES:

REGIONAL : No RWMA

GLOBAL: Normal

COLOUR DOPPLER:

MITRAL VALVE: TRIVIAL MR, E/A- 1.10

AORTIC VALVE: Normal

TRICUSPID VALVE: TRIVIAL TR, PASP-28mmHg

PULMONARY VALVE: Normal

CLOT/ VEGETATION: Nil

PERICARDIUM: No effusion

IVC : NORMAL & COLLAPSING

CONCLUSION:

- NORMAL CHAMBER AND VALVES
- NO REGIONAL WALL MOTION ABNORMALITIES
- NORMAL LV SYSTOLIC FUNCTION (EF:62%)
- NORMAL PA PRESSURE
- NO CLOT/ VEG / PERICARDIAL EFFUSION


Dr. MAHADEV SWAMY B
MBBS, MD, DM (Cardiology), FSCAI, FICC
Consultant & Interventional Cardiologist
KMC No 75242



PLEASE SCAN QR CODE

Name	: Mr . RAJU M	TID	: UMR1577070
Age/Gender	: 40 Years/Male	Registered On	: 25-May-2024 09:57 AM
Ref By	: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS	Reported On	: 25-May-2024 12:15 PM
Reg.No	: BIL4292613	Reference	: Arcofemi Health Care Ltd - Medi Whe

X-Ray Chest PA View

Lung fields appear normal.

Cardiac size is within normal limits.

Aorta and pulmonary vasculature is normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

IMPRESSION:

Normal study.

*** End Of Report ***

Dr Niharika Gupta
Consultant Radiologist