PID No. : MED410032325 Register On : 25/05/2024 9:12 AM

: 712415799 SID No. Collection On : 25/05/2024 11:41 AM Age / Sex : 42 Year(s) / Male Report On

: 25/05/2024 5:44 PM Type : OP : 28/05/2024 6:37 PM **Printed On**

Ref. Dr : MediWheel



<u>Unit</u> Investigation <u>Observed</u> **Biological** Reference Interval <u>Value</u>

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING

 $({\rm EDTA~Blood} Agglutination)$

Remark: Test to be confirmed by gel method.

'AB' 'Positive'








PID No. : MED410032325 SID No. : 712415799 Collection On : 25/05/2024 11:41 AM Age / Sex : 42 Year(s) / Male Report On

Type : OP

Ref. Dr : MediWheel



: 25/05/2024 5:44 PM : 28/05/2024 6:37 PM **Printed On**



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	<u>Value</u>		Reference Interval

HAEMATOLOGY

Complete Blood Count With - ESR

Haemoglobin	14.4	g/dL	13.5 - 18.0
-------------	------	------	-------------

(EDTA Blood/Spectrophotometry)

INTERPRETATION: Haemoglobin values vary in Men, Women & Children. Low haemoglobin values may be due to nutritional deficiency, blood loss, renal failure etc. Higher values are often due to dehydration, smoking , high altitudes , hypoxia etc.

,	•	0, 0	
PCV (Packed Cell Volume) / Haematocrit (EDTA Blood/Derived)	41.3	%	42 - 52
RBC Count (EDTA Blood/Automated Blood cell Counter)	5.04	mill/cu.mm	4.7 - 6.0
MCV (Mean Corpuscular Volume) (EDTA Blood/Derived from Impedance)	82.0	fL	78 - 100
MCH (Mean Corpuscular Haemoglobin) (EDTA Blood/Derived)	28.6	pg	27 - 32
MCHC (Mean Corpuscular Haemoglobin concentration) (EDTA Blood/Derived)	35.0	g/dL	32 - 36
RDW-CV (Derived)	13.6	%	11.5 - 16.0
RDW-SD (Derived)	39.03	fL	39 - 46
Total WBC Count (TC) (EDTA Blood/Derived from Impedance)	6740	cells/cu.mm	4000 - 11000
Neutrophils (Blood/Impedance Variation & Flow Cytometry)	63	%	40 - 75
Lymphocytes (Blood/Impedance Variation & Flow Cytometry)	27	%	20 - 45







PID No. : MED410032325 Register On : 25/05/2024 9:12 AM : 712415799 SID No. Collection On : 25/05/2024 11:41 AM Age / Sex : 42 Year(s) / Male

Report On : 25/05/2024 5:44 PM

: 28/05/2024 6:37 PM **Printed On**



Ref. Dr : MediWheel

: OP

Type

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Eosinophils (Blood/Impedance Variation & Flow Cytometry)	03	%	01 - 06
Monocytes (Blood/Impedance Variation & Flow Cytometry)	07	%	01 - 10
Basophils (Blood/Impedance Variation & Flow Cytometry)	00	%	00 - 02
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	4.25	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	1.82	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.20	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.47	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.00	10^3 / μl	< 0.2
Platelet Count (EDTA Blood/Derived from Impedance)	192	10^3 / μl	150 - 450
MPV (Blood/ <i>Derived</i>)	12.4	fL	7.9 - 13.7
PCT	0.24	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood/Automated ESR analyser)	10	mm/hr	< 15







 PID No.
 : MED410032325
 Register On
 : 25/05/2024 9:12 AM

 SID No.
 : 712415799
 Collection On
 : 25/05/2024 11:41 AM

 Age / Sex
 : 42 Year(s) / Male
 Report On
 : 25/05/2024 5:44 PM

Printed On : 28/05/2024 6:37 PM

Ref. Dr : MediWheel

: OP

Type



<u>Investigation</u>	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulfanilic Acid)	0.5	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.1	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.40	mg/dL	0.1 - 1.0
Total Protein (Serum/Biuret)	7.2	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.9	gm/dl	3.5 - 5.2
Globulin (Serum/ <i>Derived</i>)	2.30	gm/dL	2.3 - 3.6
A : G Ratio (Serum/Derived)	2.13		1.1 - 2.2
INTERPRETATION: Remark: Electrophoresis is the	preferred method		
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC / Kinetic)	28	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC / Kinetic)	31	U/L	5 - 41
Alkaline Phosphatase (SAP) (Serum/PNPP / Kinetic)	71	U/L	53 - 128
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	35	U/L	< 55







 PID No.
 : MED410032325
 Register On
 : 25/05/2024 9:12 AM

 SID No.
 : 712415799
 Collection On
 : 25/05/2024 11:41 AM

 Age / Sex
 : 42 Year(s) / Male
 Report On
 : 25/05/2024 5:44 PM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
<u>Lipid Profile</u>			
Cholesterol Total (Serum/Oxidase / Peroxidase method)	251	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol phosphate oxidase / peroxidase)	319	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

: 28/05/2024 6:37 PM

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual_circulating level of triglycerides during most part of the day.

r · · · · · · · · · · · · · · · · · · ·			
HDL Cholesterol (Serum/Immunoinhibition)	40	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	147.2	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	63.8	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	211.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220







 PID No.
 : MED410032325
 Register On
 : 25/05/2024 9:12 AM

 SID No.
 : 712415799
 Collection On
 : 25/05/2024 11:41 AM

Age / Sex : 42 Year(s) / Male Report On : 25/05/2024 5:44 PM

Printed On

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	Value		Reference Interval

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2. It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

: 28/05/2024 6:37 PM

Total Cholesterol/HDL Cholesterol Ratio	6.3	Optimal: < 3.3
(Serum/Calculated)		Low Risk: 3.4 - 4.4
		Average Risk: 4.5 - 7.1
		Moderate Risk: 7.2 - 11.0

High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio 8 Optimal: < 2.5

(TG/HDL) Mild to moderate risk: 2.5 - 5.0

(Serum/Calculated) High Risk: > 5.0

LDL/HDL Cholesterol Ratio 3.7 Optimal: 0.5 - 3.0

(Serum/Calculated)
Borderline: 3.1 - 6.0
High Risk: > 6.0









 PID No.
 : MED410032325
 Register On
 : 25/05/2024 9:12 AM

 SID No.
 : 712415799
 Collection On
 : 25/05/2024 11:41 AM

Age / Sex : 42 Year(s) / Male Report On : 25/05/2024 5:44 PM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation Glycosylated Haemoglobin (HbA1c)	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HbA1C (Whole Blood/ <i>HPLC</i>)	5.8	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

: 28/05/2024 6:37 PM

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 119.76 mg/dl

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.







 PID No.
 : MED410032325
 Register On
 : 25/05/2024 9:12 AM

 SID No.
 : 712415799
 Collection On
 : 25/05/2024 11:41 AM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	11.8		
Glucose Fasting (FBS) (Plasma - F/GOD- POD)	98	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

: 28/05/2024 6:37 PM

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Urine sugar, Fasting	Nil		Nil
(Urine - F)			
Glucose Postprandial (PPBS)	97	mg/dL	70 - 140
(Plasma - PP/GOD - POD)			

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Sugar (PP-2 hours) (Urine - PP)	Nil		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	10.7	mg/dL	7.0 - 21
Creatinine (Serum/Jaffe Kinetic)	0.9	mg/dL	0.9 - 1.3

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine etc.

Uric Acid 6.8 mg/dL 3.5 - 7.2

(Serum/Uricase/Peroxidase)







PID No. : MED410032325 Register On : 25/05/2024 9:12 AM

Printed On

: 712415799 SID No. Collection On : 25/05/2024 11:41 AM Age / Sex : 42 Year(s) / Male

Type : OP

Ref. Dr : MediWheel

: 28/05/2024 6:37 PM

Report On : 25/05/2024 5:44 PM



<u>Investigation</u>	<u>Observed</u> <u>U</u>	nit <u>Biological</u>
	Value	Reference Interval

IMMUNOASSAY

THYROID PROFILE / TFT

1.22 0.7 - 2.04T3 (Triiodothyronine) - Total ng/ml

(Serum/Chemiluminescent Immunometric Assay (CLIA))

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

4.2 - 12.0 T4 (Thyroxine) - Total 7.89 Microg/dl

(Serum/Chemiluminescent Immunometric Assay

(CLIA))

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

0.35 - 5.50 TSH (Thyroid Stimulating Hormone) 1.772 uIU/mL

(Serum/Chemiluminescent Immunometric Assay

(CLIA))

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester: 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%,hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values&lt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.







PID No. : MED410032325

SID No. : 712415799

Age / Sex : 42 Year(s) / Male

Type : OP

Ref. Dr : MediWheel

Register On : 25/05/2024 9:12 AM

Collection On : 25/05/2024 11:41 AM

Report On : 25/05/2024 5:44 PM

Printed On : 28/05/2024 6:37 PM



<u>Investigation</u>	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>IMMUNOASSAY</u>			
Total PSA (Serum/Chemiluminescent Immunometric Assay (CLIA))	0.79	ng/ml	Normal: 0.0 - 4.0 Inflammatory & Non Malignant conditions of Prostate & genitourinary system: 4.01 - 10.0

INTERPRETATION: REMARK: PSA alone should not be used as an absolute indicator of malignancy.







Suspicious of Malignant disease of Prostate: > 10.0

 PID No.
 : MED410032325
 Register On
 : 25/05/2024 9:12 AM

 SID No.
 : 712415799
 Collection On
 : 25/05/2024 11:41 AM

 Age / Sex
 : 42 Year(s) / Male
 Report On
 : 25/05/2024 5:44 PM

Type : OP **Printed On** : 28/05/2024 6:37 PM

Ref. Dr : MediWheel



i wearwheer		
Investigation	<u>Observed</u> <u>Unit</u> <u>Value</u>	<u>Biological</u> <u>Reference Interval</u>
CLINICAL PATHOLOGY		
PHYSICAL EXAMINATION		
Colour (Urine/Physical examination)	Pale Yellow	Yellow to Amber
Volume (Urine/Physical examination)	20	ml
Appearance (Urine)	Clear	
CHEMICAL EXAMINATION		
pH (Urine)	6.5	4.5 - 8.0
Specific Gravity (Urine/Dip Stick - Reagent strip method)	1.015	1.002 - 1.035
Protein (Urine/Dip Stick - Reagent strip method)	Negative	Negative
Glucose (Urine)	Nil	Nil
Ketone	Nil	Nil

Negative

Nil

Negative

Nil

Leukocytes (Urine)

Nitrite
(Urine/Dip Stick - Reagent strip method)

(Urine/Dip Stick - Reagent strip method)

Bilirubin

(Urine) Blood

(Urine)

Emohin Min

Sr.LabTechnician

VERIFIED BY

Mohan Kumar



leuco/uL

mg/dL



Negative

Nil

Negative

Nil

 PID No.
 : MED410032325
 Register On
 : 25/05/2024 9:12 AM

 SID No.
 : 712415799
 Collection On
 : 25/05/2024 11:41 AM

 $\textbf{Age / Sex} \quad \textbf{:} \quad 42 \; \text{Year(s) / Male} \qquad \qquad \textbf{Report On} \qquad \textbf{:} \quad 25/05/2024 \; 5:44 \; \text{PM}$

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Urobilinogen (Urine/Dip Stick - Reagent strip method) Urine Microscopy Pictures	Normal		Within normal limits
RBCs (Urine/Microscopy)	Nil	/hpf	NIL
Pus Cells (Urine/Microscopy)	2-3	/hpf	< 5
Epithelial Cells (Urine/Microscopy)	2-3	/hpf	No ranges
Others (Urine)	Nil		Nil

: 28/05/2024 6:37 PM







APPROVED BY

-- End of Report --



Name	Mr.MURAHARI VARMA	ID	MED410032325
Age & Gender	42/MALE	Visit Date	25/05/2024
Ref Doctor Name	MediWheel		

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in size and shows slightly increased echotexture.

No evidence of focal lesion or intrahepatic biliary ductal dilatation.

Hepatic and portal vein radicals are normal.

GALL BLADDER show normal shape and has clear contents.

Gall bladder wall is of normal thickness. CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern.

No evidence of ductal dilatation or calcification.

SPLEEN show normal shape, size and echopattern.

No demonstrable Para-aortic lymphadenopathy.

KIDNEYS move well with respiration and have normal shape, size and echopattern.

Cortico- medullary differentiations are well madeout.

No evidence of calculus or hydronephrosis.

	Bipolar length (cms)	Parenchymal thickness	
		(cms)	
Right Kidney	11.0	1.4	
Left Kidney	11.4	1.4	

URINARY BLADDER show normal shape and wall thickness.

It has clear contents. No evidence of diverticula.

PROSTATE shows normal shape, size and echopattern.

No evidence of ascites.

IMPRESSION:

> GRADE I FATTY CHANGES IN LIVER.

CONSULTANT RADIOLOGISTS

REPORT DISCLAIMER

- 1. This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11.Disputes, if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



Name	Mr.MURAHARI VARMA	ID	MED410032325
Age & Gender	42/MALE	Visit Date	25/05/2024
Ref Doctor Name	MediWheel		

DR. ANITHA ADARSH

DR. MOHAN B

REPORT DISCLAIMER

- 1.This is only a radiologincal imperssion.Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11.Disputes,if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



Name	Mr. MURAHARI VARMA	ID	MED410032325
Age & Gender	42Y/M	Visit Date	May 25 2024 9:11AM
Ref Doctor	MediWheel		

X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

Impression: No significant abnormality detected.

Dr. Anitha Adarsh Consultant Radiologist