

Interest International Bank
Service-Cum-Identy Card for Retired Employee



क.कू.सं. पदनाम बन्म तिथि

Name : Mr. Shyam Lai Mali

Employee Code: 37943 Designation: Date of Birth:

CHIEF MANAGER 02.03.1960 सेवा ग्रहण तिथि Date of Joining: 21.03.1984

सेवानिवृत तिथि Date of Retirement : 31.03.2020

Holder's Signature

मिक्ष्यं प्राधिकारी Issuing Authority

Brok.

Address:

255, Shiv Nagar-II, Murlipura Scheme, Jalpur-302039

In case of EMERGENCY

Contact: (+91) 9929458201

Blood Group : B+

Instructions:

1. Holder will be held accountable against loss, theft or damage to this card
2. Loss of card must be reported immediately to police and the bank.

If found please return this card to:

Bank of Baroda Head Office mandvi Vadodara - 390006 Gujarat-India

Call Ph.: (91)265 2562089 Drop the card any ATM of Bank of Baroda

Dr. PIYUSH GOYAL MBBS, DMRD (Radiologist) RMC No.-037041



- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ♦ +91 141 4824885

 p3healthsolutionsllp@gmail.com



General Physical Examination

Date of Examination: 31150194
Name: SHYAM LAL MOLT Age: GUYRDOB: 0210311360 Sex: Male
Referred By: NANK OFBARODA
Photo ID: BANK JD ID#: 37043
Ht: 160 (cm) Wt: 63 (Kg)
Chest (Expiration): <u>3C</u> (cm) Abdomen Circumference:
Blood Pressure: 30/80 mm Hg PR: 39/min RR: 18/min Temp: Achille
вмі
Eye Examination: RIET CIGNIC NICE LIET CIGNIC NICE
Other:
Other.
On examination he/she appears physically and mentally fit: \Yes/ No
Signature Of Examine : Name of Examinee: SHYAM LAL MALT
Signature Medical Examiner DR. PIYUSH GOYAL Name Medical Examiner DR. P. J. YUSH CIOYAL RMC No037041



- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ♦ +91 141 4824885

 p3healthsolutionsllp@gmail.com



Patient ID	12241073 Pa	tient Mob No.9950890717	Registered On	31/08/2024 08:24:50
NAME	Mr. SHYAM L	AL MALI	Collected On	31/08/2024 09:14:42
Age / Sex	Male	64 Yrs 6 Mon 1 Days	Authorized On	31/08/2024 16:24:15
Ref. By	BANK OF BAI	RODA	Printed On	31/08/2024 16:53:44
Lab/Hosp	Mr.MEDIWHEE	L		

HAEMOGARAM

HAEMATOLOGY

Test Name	Value	Unit		Biological Ref Interval		
FULL BODY HEALTH CHECKUP ABOVE 40 MALE						
HAEMOGLOBIN (Hb)	12.7 L	g/dL		13.0 - 17.0		
TOTAL LEUCOCYTE COUNT	6.00	/cumm		4.00 - 10.00		
DIFFERENTIAL LEUCOCYTE COUNT						
NEUTROPHIL	53.8	%		40.0 - 80.0		
LYMPHOCYTE	40.7 H	%		20.0 - 40.0		
EOSINOPHIL	1.3	%		1.0 - 6.0		
MONOCYTE	4.2	%		2.0 - 10.0		
BASOPHIL	0.0	%	i L	0.0 - 2.0		
TOTAL RED BLOOD CELL COUNT (RBC)	4.89	x10^6/uL		4.50 - 5.50		
HEMATOCRIT (HCT)	39.90 L	%		40.00 - 50.00		
MEAN CORP VOLUME (MCV)	82.0 L	fL		83.0 - 101.0		
MEAN CORP HB (MCH)	25.9 L	pg		27.0 - 32.0		
MEAN CORP HB CONC (MCHC)	31.7	g/dL		31.5 - 34.5		
PLATELET COUNT	154	x10^3/uL	NO.	150 - 410		
RDW-CV	15.2 H	%		11.6 - 14.0		

Technologist

DR.TANU RUNGTA MD (Pathology) RMC No. 17226

This Report Is Not Valid For Medico Legal Purpose



- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ♦ +91 141 4824885

 p3healthsolutionsllp@gmail.com



Pat	tient ID	12241073	Patient Mob No.9950890717	Registered On	31/08/2024 08:24:50
NA	ME	Mr. SHYA	M LAL MALI	Collected On	31/08/2024 09:14:42
Age	/ Sex	Male	64 Yrs 6 Mon 1 Days	Authorized On	31/08/2024 16:24:15
Ref	f. By	BANK OF	BARODA	Printed On	31/08/2024 16:53:44
Lab	/Hosp	Mr.MEDIW	HEEL		

HAEMATOLOGY

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
Erythrocyte Sedimentation Rate (ESR) Methord:- Westergreen	16 H	mm in 1st hr	00 - 15

The erythrocyte sedimentation rate (ESR or sed rate) is a relatively simple, inexpensive, non-specific test that has been used for many years to help detect inflammation associated with conditions such as infections, cancers, and autoimmune diseases.ESR is said to be a non-specific test because an elevated result often indicates the presence of inflammation but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other tests, such as C-reactive protein.ESR is used to help diagnose certain specific inflammatory diseases, including temporal arteritis, systemic vasculitis and polymyalgia rheumatica. (For more on these, read the article on Vasculitis.) A significantly elevated ESR is one of the main test results used to support the diagnosis. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as

Technologist



- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ♦ +91 141 4824885

 p3healthsolutionsllp@gmail.com



Patient ID 12241073 Patient Mob No.9950890717 Registered On 31/08/2024 08:24:50

NAME Mr. SHYAM LAL MALI

Age / Sex Male 64 Yrs 6 Mon 1 Days

Ref. By BANK OF BARODA Printed On 31/08/2024 16:53:44

Lab/Hosp Mr.MEDIWHEEL

(CBC): Methodology: TLC,DLC Fluorescent Flow cytometry, HB SLS method,TRBC,PCV,PLT Hydrodynamically focused Impedance. and MCH,MCV,MCHC,MENTZER INDEX are calculated. InstrumentName: Sysmex 6 part fully automatic analyzer XN-L,Japan



Page No: 3 of 17



- O B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ♦ +91 141 4824885

 p3healthsolutionsllp@gmail.com



Patient ID	12241073 Patient Mob No.9950890717
NAME	Mr. SHYAM LAL MALI

Mr. SHYAM LAL MALI Age / Sex

64 Yrs 6 Mon 1 Days

BANK OF BARODA Ref. By Lab/Hosp Mr.MEDIWHEEL

Registered On

31/08/2024 08:24:50

Collected On

31/08/2024 09:14:42

Authorized On Printed On

31/08/2024 16:24:15

31/08/2024 16:53:44

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
FASTING BLOOD SUGAR (Plasma) Methord:- GLUCOSE OXIDASE/PEROXIDASE	165.0 H	mg/dl	70.0 - 115.0
Impaired glucose tolerance (IGT)	111	- 125 mg/dL	
Diabetes Mellitus (DM)	> 1:	26 mg/dL	

Instrument Name: HORIBA CA60 Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic

hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin

therapy or various liver diseases.

BLOOD SUGAR PP (Plasma) Methord:- GLUCOSE OXIDASE/PEROXIDASE

211.9 H

mg/dl

70.0 - 140.0

Instrument Name: HORIBA Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels(hypoglycemia) may result from excessive insulin therapy or various liver diseases.

Technologist

DR.TANU RUNGTA MD (Pathology)

RMC No. 17226



- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ♦ +91 141 4824885

 p3healthsolutionsllp@gmail.com



Patient ID 12241	073	Patient	Mob	No.	9950890717
------------------	-----	---------	-----	-----	------------

NAME Age / Sex

Mr. SHYAM LAL MALI

64 Yrs 6 Mon 1 Days BANK OF BARODA

Ref. By Lab/Hosp Mr.MEDIWHEEL Registered On

31/08/2024 08:24:50

Collected On

31/08/2024 09:14:42

Authorized On

31/08/2024 16:24:15

Printed On

31/08/2024 16:53:44

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
GLYCOSYLATED HEMOGLOBIN (Hb.A. Methord:- CAPILLARY with EDTA	6.9	mg%	Non-Diabetic < 6.0 Good Control 6.0-7.0 Weak Control 7.0-8.0 Poor control > 8.0
MEAN PLASMA GLUCOSE Methord:- Calculated Parameter	151 H	mg/dL	68 - 125

INTERPRETATION

AS PER AMERICAN DIABETES ASSOCIATION (ADA) Reference Group HbA1c in % Non diabetic adults >=18 years < 5.7 At risk (Prediabetes) 5.7 - 6.4 Diagnosing Diabetes >= 6.5

CLINICAL NOTES

In vitro quantitative determination of HbA1c in whole blood is utilized in long term monitoring of glycemia. The HbA1c level correlates with the mean glucose concentration prevailing in the course of the patient's recent history (approx - 6-8 weeks) and therefore provides much more reliable information for glycemia monitoring than do determinations of blood glucose or urinary glucose. It is recommended that the determination of HbA1c be performed at intervals of 4-6 weeks during Diabetes Mellitus therapy. Results of HbA1c should be assessed in conjunction with the patient's medical history, clinical examinations and other findings. Some of the factors that influence HbA1c and its measurement [Adapted from Gallagher et al]

- 1. Erythropoiesis
- Increased HbA1c: iron, vitamin B12 deficiency, decreased erythropoiesis
- Decreased HbA1c: administration of erythropoietin, iron, vitamin B12, reticulocytosis, chronic liver disease
- 2. Altered Haemoglobin-Genetic or chemical alterations in hemoglobin: hemoglobinopathies, HbF, methemoglobin, may increase or decrease HbA1c.
- 3. Glycation
- Increased HbA1c: alcoholism, chronic renal failure, decreased intraerythrocytic pH.
- Decreased HbA1c: certain hemoglobinopathies, increased intra-erythrocyte pH
- 4. Erythrocyte destruction
- Increased HbA1c: increased erythrocyte life span: Splenectomy
- Decreased A1c: decreased RBC life span: hemoglobinopathies, splenomegaly, rheumatoid arthritis or drugs such as antiretrovirals, ribavirin & dapsone.
- 5. Others
- Increased HbA1c: hyperbilirubinemia, carbamylated hemoglobin, alcoholism, large doses of aspirin, chronic opiate use, chronic renal failure

- Decreased HbA1c: hypertriglyceridemia, reticulocytosis, chronic liver disease, aspirin, vitamin C and E, splenomegaly, rheumatoid arthritis or drugs

Technologist

DR.TANU RUNGTA MD (Pathology)

RMC No. 17226



- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- 🕒 +91 141 4824885 🖾 p3healthsolutionsllp@gmail.com



Patient ID 12241073 Patient Mob No.9950890717

NAME Mr. SHYAM LAL MALI

Age / Sex Male 64 Yrs 6 Mon 1 Days

Ref. By BANK OF BARODA

Lab/Hosp Mr.MEDIWHEEL

Registered On

31/08/2024 08:24:50

Collected On

31/08/2024 09:14:42 31/08/2024 16:24:15

Authorized On Printed On

31/08/2024 16:53:44

HAEMATOLOGY

HAEMATOLOGY

Test Name Value Unit

Biological Ref Interval

BLOOD GROUP ABO Methord:- Haemagglutination reaction "A" POSITIVE

Technologist



Ref. By

P3 HEALTH SOLUTIONS LLP

- O B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- 🕒 +91 141 4824885 🖂 p3healthsolutionsllp@gmail.com



Patient ID 12241073 Patient Mob No.9950890717

NAME Age / Sex

Mr. SHYAM LAL MALI

64 Yrs 6 Mon 1 Days BANK OF BARODA

Mr.MEDIWHEEL Lab/Hosp

Registered On

31/08/2024 08:24:50

Collected On

31/08/2024 09:14:42

Authorized On

31/08/2024 16:24:15

Printed On

31/08/2024 16:53:44

BIOCHEMISTRY

Test Name Value Unit Biological Ref Interval

LIPID PROFILE

SERUM TOTAL CHOLESTEROL

Methord:- CHOLESTEROL OXIDASE/PEROXIDASE

219.00

mg/dl

Desirable <200

Borderline 200-239

High> 240

InstrumentName: HORIBA Interpretation: Cholesterol measurements are used in the diagnosis and treatments of lipid lipoprotein metabolism

SERUM TRIGLYCERIDES

Methord:- GLYCEROL PHOSPHATE OXIDASE/PREOXIDASE

atherosclerosis or reduce its progress and to avoid plaque rupture.

98.70

mg/dl

Normal

<150

Borderline high 150-199 High 200-499

High Very high

>500

InstrumentName: Randox Rx Imola Interpretation: Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction.

DIRECT HDL CHOLESTEROL

Methord:- Direct clearance Method

37.00

mg/dl

MALE- 30-70 FEMALE - 30-85

Instrument Name: Rx Daytona plus Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods LDL CHOLESTEROL

Methord:- Calculated Method

165.55 H

mg/dl

Optimal <100

Near Optimal/above optimal

100-129

Borderline High 130-159

High 160-189 Very High > 190

Interpretation: Accurate measurement of LDL-Cholesterol is of vital importance in therapies which focus on lipid reduction to prevent

VLDL CHOLESTEROL Methord:- Calculated

19.74

mg/dl

0.00 - 80.00

Technologist

DR.TANU RUNGTA

MD (Pathology) RMC No. 17226

Janu



Lab/Hosp

Mr.MEDIWHEEL

P3 HEALTH SOLUTIONS LLP

- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- 🕒 +91 141 4824885 🖂 p3healthsolutionsllp@gmail.com



Patient ID	nt ID 12241073 Patient Mob No.9950890717		Registered On	31/08/2024 08:24:50
NAME	ME Mr. SHYAM LAL MALI		Collected On	31/08/2024 09:14:42
Age / Sex	Male	64 Yrs 6 Mon 1 Days	Authorized On	31/08/2024 16:24:15
Ref. By	BANK OF	BARODA	Printed On	31/08/2024 16:53:44

BIOCHEMISTRY

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
T.CHOLESTEROL/HDL CHOLESTEROL RATIO Methord:- Calculated	5.92 H		0.00 - 4.90
LDL / HDL CHOLESTEROL RATIO Methord:- Calculated	4.47 H		0.00 - 3.50
TOTAL LIPID	613.27	mg/dl	400.00 - 1000.00

^{1.} Measurements in the same patient can show physiological& analytical variations. Three serialsamples I week apart are recommended for Total Cholesterol, Triglycerides, HDL& LDL Cholesterol.

Technologist Page No. 8 grist

^{2.} As per NCEP guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended

³ Low HDL levels are associated with Coronary Heart Disease due to insufficient HDL being available to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.



- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ♦ +91 141 4824885

 p3healthsolutionsllp@gmail.com



Patient ID	12241073 Patient Mob No.9950890717	S (500 to 100	A 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
· additib	12241073 T attent WOD 140.9930090717	Registered On	31/08/2024 08:24:50	
NAME	Mr. SHYAM LAL MALI	Collected On	31/08/2024 09:14:42	
Age / Sex	Male 64 Yrs 6 Mon 1 Days	Authorized On	31/08/2024 16:24:15	
Ref. By	BANK OF BARODA	Printed On	31/08/2024 16:53:44	
Lab/Hosp	Mr.MEDIWHEEL	· ····································	01/00/2024 10:55:44	

BIOCHEMISTRY

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
LIVER PROFILE WITH GGT		_ler	
SERUM BILIRUBIN (TOTAL) Methord:- DIAZOTIZED SULFANILIC	0.51	mg/dL	Infants : 0.2-8.0 mg/dL Adult - Up to - 1.2 mg/dL
SERUM BILIRUBIN (DIRECT) Methord:- DIAZOTIZED SULFANILIC	0.16	mg/dL	Up to 0.40 mg/dL
SERUM BILIRUBIN (INDIRECT) Methord:- Calculated	0.35	mg/dl	0.30-0.70
SGOT Methord:- IFCC	15.8	U/L	0.0 - 40.0
SGPT Methord:- IFCC	18.3	U/L	0.0 - 40.0
SERUM ALKALINE PHOSPHATASE Methord:- DGKC - SCE	99.00	U/L	53.00 - 141.00
SERUM GAMMA GT Methord:- Szasz methodology Instrument Name Randox Rx Imola Interpretation: Elevations in GGT levels areseen earlier and more pronounce	26.00 ed than those with other liver en	U/L zymes in cases of obstructive jaundice and	10.00 - 45.00
metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or prhepatic biliary obstruction. Only moderate elevations in the enzyme level (2		with infectious hepatitis.	
SERUM TOTAL PROTEIN Methord:- BIURET	5.07 L	g/dl	6.00 - 8.40
SERUM ALBUMIN Methord:- BROMOCRESOL GREEN	4.19	g/dl	3.50 - 5.50
SERUM GLOBULIN Methord:- CALCULATION	0.88	gm/dl	2.20 - 3.50
A/G RATIO	4.76 H		1.30 - 2.50

Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

Note: These are group of tests that can be used to detect the presence of liver disease, distinguish among different types of liver disorders, gauge the extent of known liver damage, and monitor the response to treatment. Most liver diseases cause only mild symptoms initially, but these diseases must be detected early. Some tests are associated with functionality (e.g.,

Technologist



 B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023

♦ +91 141 4824885

p3healthsolutionsllp@gmail.com



Patient ID 12241073 Patient Mob No.9950890717

NAME Mr. SHYAM LAL MALI

Age / Sex Male 64 Yrs 6 Mon 1 Days

Ref. By BANK OF BARODA

Lab/Hosp Mr.MEDIWHEEL

Registered On 31/08/2024 08:24:50

Collected On 31/08/2024 09:14:42

Authorized On 31/08/2024 16:24:15 Printed On 31/08/2024 16:53:44

BIOCHEMISTRY

BIOCHEMISTRY

Test Name Value Unit Biological Ref Interval

albumin), some with cellular integrity (e.g., transaminase), and some with conditions linked to the biliary tract (gamma-glutamyl transferase and alkaline phosphatase). Conditions with elevated levels of ALT and AST include hepatitis A,B,C, paracetamol toxicity etc. Several biochemical tests are useful in the evaluation and management of patients with hepatic dysfunction. Some or all of these measurements are also carried out (usually about twice a year for routine cases) on those individuals taking certain medications, such as

Technologist 7



- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- 🕲 +91 141 4824885 ⊠ p3healthsolutionsllp@gmail.com



Patient ID	12241073 Patient Mob No.9950890717	Registered On	31/08/2024 08:24:50
NAME	Mr. SHYAM LAL MALI	Collected On	31/08/2024 09:14:42
Age / Sex	Male 64 Yrs 6 Mon 1 Days	Authorized On	31/08/2024 16:24:15
Ref. By	BANK OF BARODA	Printed On	31/08/2024 16:53:44
Lab/Hosp	Mr.MEDIWHEEL		

BIOCHEMISTRY

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
RFT / KFT WITH ELECTROLYTES	and the same of	4	
SERUM UREA Methord:- UREASE / GLUTAMATE DEHYDROGENASE	23.00	mg/dl	10.00 - 50.00
InstrumentName: HORIBA CA 60 Interpretation : diseases.	Urea measurements are	e used in the diagnosis an	d treatment of certain renal and metabolic
SERUM CREATININE Methord:- JAFFE	0.96	mg/dl	Males : 0.6-1.50 mg/dl Females : 0.6 -1.40 mg/dl
Interpretation: Creatinine is measured primarily to assess kidney funct relatively independent of protein ingestion, water intaked clinically significant. SERUM URIC ACID			
Methord:- URICASE/PEROXIDASE InstrumentName:HORIBA YUMIZEN CA60 Dayto	na plus Interpretation	n: Elevated Urate: High p	urine diet, Alcohol. Renal insufficiency, Drugs,
Polycythaemia vera, Malignancies, Hypothyroidism, Rar			rndrome, Pregnancy,Gout.
SODIUM Methord:- Ion-Selective Electrode with Serum	135.0	mmol/L	
			135 - 150
POTASSIUM Methord:- ISE	3.86	mmol/L	3.50 - 5.50
CHLORIDE Methord:- Ion-Selective Electrode with Serum	104.8	mmol/L	
Methora:- Ion-Selective Electrode with Serum			98 - 106
SERUM CALCIUM Methord:- Arsenazo III Method	10.40 H	mg/dL	8.80 - 10.20
InstrumentName:MISPA PLUS Interpretation: S Increases in serum PTH or vitamin D are usually ass			

nephrosis and pancreatitis.

g/dl

SERUM TOTAL PROTEIN Methord:- BIURET

5.07 L

6.00 - 8.40

Technologist7



- O B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- 🕲 +91 141 4824885 🖂 p3healthsolutionsllp@gmail.com



Patient ID	12241073 Patient Mob No.9950890717	Registered On	31/08/2024 08:24:50
NAME	Mr. SHYAM LAL MALI	Collected On	31/08/2024 09:14:42
Age / Sex	Male 64 Yrs 6 Mon 1 Days	Authorized On	31/08/2024 16:24:15
Ref. By	BANK OF BARODA	Printed On	31/08/2024 16:53:44

Ref. By BANK OF BARODA Lab/Hosp Mr.MEDIWHEEL

BIOCHEMISTRY

BIOCHEMISTRY

		A CONTRACT OF THE PROPERTY OF	
Test Name	Value	Unit	Biological Ref Interval
SERUM ALBUMIN Methord:- BROMOCRESOL GREEN	4.19	g/dl	3.50 - 5.50
SERUM GLOBULIN Methord:- CALCULATION	0.88 └	gm/dl	2.20 - 3.50
A/G RATIO	4.76 H		1.30 - 2.50

Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

Kidney function tests are group of tests that can be used to evaluate how well the kidneys are functioning. Creatinine is a waste product that comes from protein in the diet and also comes from the normal wear and tear of muscles of the body. In blood, it is a marker of GFR .in urine, it can remove the need for 24-hourcollections for many analytes or be used as a quality assurance tool to assess the accuracy of a 24-hour collection Higher levels may be a sign that the kidneys are not working properly. As kidney disease progresses, the level of creatinine and urea in the bloodincreases. Certain drugs are nephrotoxic hence KFT is done before and after initiation of treatment with these drugs

Low serum creatinine values are rare; they almost always reflect low muscle mass

Apart from renal failure Blood Urea can increase in dehydration and GI bleed

Technologist 7



- © B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ③ +91 141 4824885 ⊠ p3healthsolutionsllp@gmail.com

64 Yrs 6 Mon 1 Days



Patient ID 12241073 Patient Mob No.9950890717

NAME Mr. S

Mr. SHYAM LAL MALI

Age / Sex Ref. By

BANK OF BARODA

Lab/Hosp

Mr.MEDIWHEEL

Registered On

31/08/2024 08:24:50

Collected On

31/08/2024 09:14:42 31/08/2024 16:24:15

Authorized On Printed On

31/08/2024 16:53:44

CLINICAL PATHOLOGY

CLINICAL PATHOLOGY

Test Name	Value	Unit	Biological Ref Interval
URINE SUGAR (FASTING) Collected Sample Received	Nil		Nil
URINE SUGAR PP Collected Sample Received	TRACE		Nil

Technologist 7



- O B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- 🕲 +91 141 4824885 🖂 p3healthsolutionsllp@gmail.com



Patient ID 12241073	Patient Mob	No.9950890717
---------------------	-------------	---------------

NAME Age / Sex

Mr. SHYAM LAL MALI

64 Yrs 6 Mon 1 Days

BANK OF BARODA Ref. By Lab/Hosp Mr.MEDIWHEEL

Registered On

31/08/2024 08:24:50

Collected On

31/08/2024 09:14:42

Authorized On

31/08/2024 16:24:15

Printed On

31/08/2024 16:53:44

IMMINOASSAV

		0.100.11	
Test Name	Value	Unit	Biological Ref Interval

PSA (PROSTATE SPECIFIC ANTIGEN) -TOTAL

ng/mL

0.00-4.00

CLINICAL NOTES:- Prostate-specific antigen (PSA) is a 34-kD glycoprotein produced almost exclusively by the prostate gland.

PSA is normally present in the blood at very low levels. Increased levels of PSA may suggest the presence of prostate cancer.

1.Immediate PSA testing following digital rectal examination, ejaculation, prostatic massage, indwelling catheterization, ultrasonography and needle biopsy of prostate is not

recommended as they falsely elevate levels

- 2. PSA values regardless of levels should not be interpreted as absolute evidence of the presence or absence of disease. All values should be correlated with clinical findings and other investigations
- 3. Physiological decrease in PSA level by 18% has been observed in sedentary patients either due to supine position or suspended sexual activity

Clinical Use

- · An aid in the early detection of Prostate cancer when used in conjunction with Digital rectal examination in males more than 50 years of age and in those with two or more affected first degree relatives.
- · Follow up and management of Prostate cancer patients
- · Detect metastatic or persistent disease in patients following surgical or medical treatment of Prostate cancer

PSA levels can be also increased by prostatitis, irritation, benign prostatic hyperplasia (BPH), and recent ejaculation, producing a false positive result. Digital rectal examination (DRE) has been shown in several studies to produce an increase in PSA. However, the effect is clinically insignificant, since DRE causes the most substantial increases in patients with PSA levels already elevated over 4.0 ng/mL.

Obesity has been reported to reduce serum PSA levels. Delayed early detection may partially explain worse outcomes in obese men with early prostate cancer. Aftertreatment, higher BMI also correlates to higher risk of recurrence.

Technologist₇

DR.TANU RUNGTA

MD (Pathology) RMC No. 17226



- O B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- 🕲 +91 141 4824885 🖾 p3healthsolutionsllp@gmail.com



Patient ID	12241073 Patient Mob No.9950890717	Registered On	31/08/2024 08:24:50
NAME	Mr. SHYAM LAL MALI	Collected On	31/08/2024 09:14:42
Age / Sex	Male 64 Yrs 6 Mon 1 Days	Authorized On	31/08/2024 16:24:15
Ref. By	BANK OF BARODA	Printed On	31/08/2024 16:53:44
Lab/Hosp	Mr.MEDIWHEEL		

IMMUNOASSAY

IMMUNOASSAY

Test Name	Value	Unit	Biological Ref Interval
TOTAL THYROID PROFILE	1		
THYROID-TRIIODOTHYRONINE T3 Methord:- Chemiluminescence	1.14	ng/m	0.07 4.70
THYROID - THYROXINE (T4) Methord:- Chemiluminescence	7.96	ug/dl	0.87 - 1.78
			4.82 -15.65
TSH Methord:- Chemiluminescence	3.540	uIU/ml	0.380 - 5.330

Interpretation Notes:

4th Generation Assay, Reference ranges vary between laboratories.

PREGNANCY - REFERENCE RANGE for TSH IN ulU/mL (As per American Thyroid Association)

1st Trimester: 0.10-2.50 uIU/mL 2nd Trimester: 0.20-3.00 uIU/mL 3rd Trimester: 0.30-3.00 uIU/mL

The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy.

NOTE-TSH levels are subject to circardian variation reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result.

- 1.Primary hyperthyroidism is accompanied by \uparrow serum T3 & T4 values along with \downarrow TSH level.
- 2.Primary hypothyroidism is accompanied by ↓ " serum T3 and T4 values &↑ 'serum TSH levels
- 3.Normal T4 levels accompanied by † 'T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis
- 4.Normal or" ↓ T3 &↑ 'T4 levels indicate T4 Thyrotoxicosis (problem is conversion of T4 to T3)
- 5.Normal T3 & T4 along with "TSH indicate mild / Subclinical Hyperthyroidism.

COMMENTS: Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test.

Disclaimer-TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age ,and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly.

Reference ranges are from Teitz fundamental of clinical chemistry 8th ed (2018)

Test performed by Instrument: Beckman coulter Dxi 800.

*** End of Report ***

Technologist 7



- O B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ♦ +91 141 4824885
 p3healthsolutionsllp@gmail.com



12241073 Patient Mob No.9950890717 Patient ID

NAME Age / Sex

Mr. SHYAM LAL MALI 64 Yrs 6 Mon 1 Days

Ref. By

BANK OF BARODA

Lab/Hosp

Mr.MEDIWHEEL

Registered On

31/08/2024 08:24:50

Collected On

31/08/2024 09:14:42 31/08/2024 16:24:15

Authorized On Printed On

31/08/2024 16:53:44

CLINICAL PATHOLOGY

Test Name	Value	Unit		Biological Ref Interval
Urine Routine PHYSICAL EXAMINATION	September 1	-		
COLOUR	PALE YEL	LOW		PALE YELLOW
APPEARANCE	Clear			Clear
CHEMICAL EXAMINATION				
REACTION(PH)	6.5			5.0 - 7.5
SPECIFIC GRAVITY	1.015			1.010 - 1.030
PROTEIN	NIL			NIL
SUGAR	NIL			NIL
BILIRUBIN	NEGATIV	E	nec.	NEGATIVE
UROBILINOGEN	NORMAL			NORMAL
KETONES	NEGATIV	E		NEGATIVE
NITRITE	NEGATIV	E		NEGATIVE
MICROSCOPY EXAMINATION				
RBC/HPF	NIL	/HPF		NIL
WBC/HPF	2-3	/HPF		2-3
EPITHELIAL CELLS	0-1	/HPF		2-3
CRYSTALS/HPF	ABSENT			ABSENT
CAST/HPF	ABSENT			ABSENT
AMORPHOUS SEDIMENT	ABSENT			ABSENT
BACTERIAL FLORA	ABSENT			ABSENT
YEAST CELL	ABSENT			ABSENT
OTHER	ABSENT			

Technologist₇







- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ♦ +91 141 4824885 ⊠ maxcarediagnostics1@gmail.com



Mr. SHYAM LAL MALI	64 Yrs./Male		
Registration Date: 31/08/2024	Ref. by: BANK OF BARODA		

ULTRASOUND OF WHOLE ABDOMEN

Liver is of normal size (14.0 cm). **Mild fatty infiltration**. No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is well distended. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape. Echotexture is normal. No focal lesion is seen.

Kidneys are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. Collecting system does not show any calculus or dilatation.

Right kidney is measuring approx. 9.0 x 4.0 cm.

Left kidney is measuring approx. 9.6 x 4.1 cm.

Urinary bladder is well distended and does not show any calculus or mass lesion.

Prostate is normal in size (21.0 cc) with normal echotexture and outline.


No enlarged nodes are visualized. No retro-peritoneal lesion is identified. No significant free fluid is seen in pelvis.

IMPRESSION:-

Grade I fatty liver.

Farm

DR. ROHAN GAUR M.B.B.S, M.D (Radiodiagnosis) RMC no. 17887





- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- 🕲 +91 141 4824885 🖂 maxcarediagnostics1@gmail.com

NORMAL



Mr. SHYAM LAL MALI	64 Yrs./Male			
Registration Date: 31/08/2024	Ref. by: BANK OF BARODA			

TRICUSPID VALVE

2D-ECHOCARDIOGRAPHY M.MODE WITH DOPPLER STUDY:

FAIR TRANSTHORACIC ECHOCARIDIOGRAPHIC WINDOW MORPHOLOGY:

WITHAL VALVE		IVOI	MAL		1100	COSFID VALVE		NONIVIA	L	
AORTIC VALVE N		NOF	NORMAL			PULMONARY VALVE			NORMAL	
				M.MOD	E EXAMITAT	ION:				
AO	3.1	Cm	LA		3.0	cm	IVS-D	0.9	cm	
IVS-S	1.2	cm	LVI	D	4.3	cm	LVSD	2.9	cm	
LVPW-D	0.9	cm	LVF	PW-S	1.3	cm	RV		cm	
RVWT		cm	ED	v	at 1	MI	LVVS		ml	
LVEF	55-60%			ST. ST. ST.	RWM	RWMA ABSENT				
	7		A	<u>C</u>	HAMBERS:	la l				
LA	NORM	1AL	RA			NORM		RMAL		
LV	NORM	1AL	29	RV			NORMAL			
PERICARDIUM		A	7	NORMAI						
				COLC	UR DOPPLE	R:				
		MITRAL	VALVE	. Ipsi	est Vib.	WE AS				
E VELOCITY 0.53		m/sec PEAK (K GRADIENT	RADIENT		Mm/hg			
A VELOCITY 0.89		m/sec MEAN		N GRADIEN	GRADIENT		Mm/hg			
MVA BY PHT		Cm2 MVA		BY PLANIN	Y PLANIMETRY		Cm2			
MITRAL REGUR	RGITATION	ESA		120		MILD /	#		*	
		AORTIC	VALVE				2			
PEAK VELOCITY 0.86		0.86	m/sec		PEAK G	PEAK GRADIENT		mm/hg		
AR VMAX		m/sec		MEAN	MEAN GRADIENT		mm/hg			
AORTIC REGURGITATION			ABSEN			T				
		TRICUSP	ID VAL	VE						
PEAK VELOCITY		7	m/sec	PEAK G	PEAK GRADIENT		n			
MEAN VELOCITY			m/sec	MEAN	MEAN GRADIENT		n	nm/hg		
VMax VELOCIT	ΓΥ			794.00					310.25	
TRICUSPID REG	URGITATION				MILD					
		PULMO	NARY '	VALVE					1	
PEAK VELOCITY		0.83		M/sec.	PEAK GRADI			Mm/hg		
MEAN VALOCITY						MEAN GRADIENT			Mm/hg	
PULMONARY I	REGURGITA	TION				ABSENT				

Impression—

MITRAL VALVE

- NORMAL LV SIZE & CONTRACTILITY.
- NO RWMA, LVEF 55-60%.
- MILD TR/ PAH (RVSP 28 MMHG+ RAP), MILD MR.
- GRADE I DIASTOLIC DYSFUNCTION.
- NO CLOT, NO VEGETATION, NO PERICARDIAL EFFUSION.

(Cardiologist)

NORMAL



- B-14, Vidhyadhar Nagar Enclave-II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur-302 023
- ♦ +91 141 4824885

 p3healthsolutionsllp@gmail.com



NAME:	MR. SHYAM LAL MALI	AGE	64 YRS/M
REF.BY	BANK OF BARODA	DATE	31/08/2024

CHEST X RAY (PA VIEW)

Bilateral lung fields appear clear.

Bilateral costo-phrenic angles appear clear.

Cardiothoracic ratio is normal.

Thoracic soft tissue and skeletal system appear unremarkable.

Soft tissue shadows appear normal.

IMPRESSION: No significant abnormality is detected

(Kgam)

DR. ROHAN GAUR
M.B.B.S, M.D (Radiodiagnosis)
RMC no. 17887

B-14 VIDHYDHAR NAGAR, (JAIPUR) Ref.: BANK OF BARODA Test Date: 31-Aug-2024(11:46:14) Notch: 50Hz 0.05Hz - 100Hz 1225360/Shyam Lal Mali 64Yrs/Male (P3 HEALTH SOLUTION LLP) Vent Rate: 74 hpm; PR Interval: 110 ms; QRS Duration: 138 msQT/QTc Int: 363/405 ms P-QRS-T axis: 58-48-21-(Deg) Comments: FINDINGS: Normal Sinus Rhythm # **V**4 2 RMS Kgs/ Cms BP: avF 10mm/mV 25mm/Sec HR: 74 bpm MAC 5 6 Horak 7 QRS Duration: 138 ms QT/QTc: 363/405ms P-QRS-T Axis: 58 - 48 - 21 (Deg) PR Interval: 110 ms BBS. DIP. CARDIO (ESCORTS)
DENARES WOHKKA aresn Kumar Mohanka

